

Research Article

Roberto Alicandro, Lucia De Luca*, Mariapia Palombaro and Marcello Ponsiglione

Γ -convergence analysis of the nonlinear self-energy induced by edge dislocations in semi-discrete and discrete models in two dimensions

<https://doi.org/10.1515/acv-2023-0053>

Received May 3, 2023; accepted October 2, 2023

Abstract: We propose nonlinear semi-discrete and discrete models for the elastic energy induced by a finite system of edge dislocations in two dimensions. Within the *dilute regime*, we analyze the asymptotic behavior of the nonlinear elastic energy, as the *core-radius* (in the semi-discrete model) and the lattice spacing (in the purely discrete one) vanish. Our analysis passes through a linearization procedure within the rigorous framework of Γ -convergence.

Keywords: Dislocations, nonlinear elasticity, plasticity, discrete-to-continuum limits, Γ -convergence

MSC 2020: 74C05, 58K45, 70G75, 74G65, 49J45

Communicated by: Ulisse Stefanelli

Introduction

We introduce some variational nonlinear models for the elastic energy induced by a finite system of edge dislocations in two dimensions, proposing and analyzing a purely discrete model settled in the regular triangular lattice, together with (and relying on) a variant of the core radius approach in [26].

We start by describing our variational discrete model in absence of defects. We assume to have a finite portion of the regular triangular lattice, with lattice spacing ε (eventually vanishing), occupying a reference configuration $\Omega \subset \mathbb{R}^2$. In this framework, deformations are described through discrete strain fields β which are defined on the nearest neighbor bonds of the lattice. In the defect-free case, these strains are in fact discrete gradients of some implicitly defined (by integration) deformation function defined on the nodes. The underlying compatibility condition is that the discrete circulation of β around each triangular cell is trivial. In the present formulation the discrete energy is thus a function of the deformation strain β and is given by the sum of two contributions: a two-body interaction potential accounting for the elongation of the bonds, and a three-body interaction term penalizing changes of area of the triangular cells. In the thermodynamic limit (as the lattice spacing vanishes), this model gives back nowadays classical continuous nonlinear models, where the energy density behaves like the squared distance of the strain from the set of rotations; in the small strain limit, one gets,

***Corresponding author: Lucia De Luca**, IAC-CNR, Istituto per le Applicazioni del Calcolo “M. Picone”, via dei Taurini 19, 00185 Roma, Italy, e-mail: lucia.deluca@cnr.it. <https://orcid.org/0000-0002-5544-4624>

Roberto Alicandro, DIEI, Università di Cassino e del Lazio meridionale, via Di Biasio 43, 03043 Cassino (FR), Italy, e-mail: alicandr@unicas.it. <https://orcid.org/0000-0002-8978-8548>

Mariapia Palombaro, DISIM, Università dell’Aquila, Via Vetoio, 67100 L’Aquila, Italy, e-mail: mariapia.palombaro@univaq.it. <https://orcid.org/0000-0001-8153-9769>

Marcello Ponsiglione, Dipartimento di Matematica, Università di Roma-I “La Sapienza”, Piazzale Aldo Moro 5, 00185 Roma, Italy, e-mail: ponsigli@mat.uniroma1.it. <https://orcid.org/0000-0002-0839-5575>

by linearization, the classical isotropic (continuous) linearized elasticity (we refer to [5, 8, 27] for discrete-to-continuum and linearization results in the context of short and long-range interaction energies). Remarkably, by tuning the pre-factors in front of the two contributions of the nonlinear discrete energy, the simultaneous discrete-to-continuum limit and linearization process yield in the limit functional all possible Lamé coefficients (see formula (2.6)).

In this framework, edge dislocations can be introduced as topological singularities of the discrete strain field β , namely, on each cell εT , we enforce the circulation of β to belong to (a rotation of) the ε -spaced triangular lattice. If non-zero, such a vector represents the so-called *Burgers vector* $\varepsilon \xi$, which detects and quantifies the presence of an edge dislocation in the triangle εT . We thus identify such a dislocation with the weighted Dirac mass $\varepsilon \xi \delta_{x_{\varepsilon T}}$, where $x_{\varepsilon T}$ denotes the barycenter of the triangle εT . A finite distribution of edge dislocations, centered at the points x^n and having Burgers vectors $\varepsilon \xi^n \in \mathbb{R}^2$, can be then identified with the empirical measure $\mu = \varepsilon \sum_n \xi^n \delta_{x^n}$. For any given configuration of dislocations μ , the class of admissible strains is given by those discrete strain fields whose discrete circulation around each triangle εT is given by $\mu(\varepsilon T)$. The presence of edge dislocations enforces some distortion on any admissible strain: since pure rotations become incompatible, a resulting stored elastic energy is induced. For a single dislocation, such an energy is of order $\varepsilon^2 |\log \varepsilon|$. Our analysis is performed in a dilute regime of edge dislocations, where the total stored energy is bounded by $C\varepsilon^2 |\log \varepsilon|$, which is consistent with the presence of a finite number of dislocations. This energy regime has been widely analyzed in terms of Γ -convergence, both for continuous models, also referred to as *semi-discrete* models, (see [14, 16] for the linear case and [26] for a nonlinear one) and for discrete linear energies [4]. (For analogous results in the linear scalar framework of screw dislocations and of vortices in superconductivity we refer the interested reader to [1–3, 13, 21, 25].)

As the lattice spacing vanishes, the discrepancy between discrete and continuous models should also vanish; similarly, since we are dealing with a vanishing energy regime of order $\varepsilon^2 |\log \varepsilon|$, also the energy gap between nonlinear and linearized models should vanish. The purpose of this paper is to rigorously prove these facts, by proposing a purely discrete nonlinear model (the first to our knowledge) and building a bridge from discrete to continuous and from nonlinear to linear edge dislocation models. For this purpose, we first introduce and analyze a slight variant of the (nonlinear) continuous model studied in [26], which will be instrumental to the discrete problem. Then we introduce our purely discrete nonlinear model, inspired from the linear discrete models in [4, 6, 17], and we show that it behaves like the continuous one, as $\varepsilon \rightarrow 0$.

The nonlinear continuous model we consider is based on the so-called *core-radius approach*: Given a configuration of dislocations $\mu = \varepsilon \sum_n \xi^n \delta_{x^n}$, we drill an ε -disc around each dislocation x^n and set

$$\Omega_\varepsilon(\mu) := \Omega \setminus \bigcup_n \overline{B}_\varepsilon(x^n).$$

Here, ε is (proportional to) the lattice spacing and the discs $B_\varepsilon(x^n)$ represent the so-called *cores* of the dislocations x^n , where plastic effects take place. Removing the cores means to neglect such plastic effects, whose energy contribution is expected to be a lower order perturbation with respect to the elastic energy computed outside the cores, namely, in $\Omega_\varepsilon(\mu)$. The admissible strains, in analogy with the discrete setting, are those fields β defined on $\Omega_\varepsilon(\mu)$ whose circulation around each $B_\varepsilon(x^n)$ is equal to $\varepsilon \xi^n$. The corresponding elastic energy of β (and, in turn, induced by μ) behaves like the integral on $\Omega_\varepsilon(\mu)$ of the squared distance of β from the set of rotations $\text{SO}(2)$.

Now we describe the further kinematic condition that we impose on the admissible strains. To this end, assume for the time being that the distribution of dislocations is given by a single dislocation $\varepsilon \xi^0 \delta_{x^0}$, with $x^0 \in \Omega$ and ξ^0 belonging to the R^0 -rotation of the regular triangular lattice. In such a case, the rigidity estimate [15] together with the energetic bound, guarantees that β is close to a constant rotation $R \in \text{SO}(2)$ plus an ε -multiple of a linear strain β^{lin} satisfying the circulation condition

$$\int_{\partial B_\varepsilon(x^0)} \beta^{\text{lin}} t \, d\mathcal{H}^1 = \xi^0.$$

In general, the rotation R around which the linearization takes place and the rotation R^0 associated to the Burgers vector are decoupled. However, if $R \neq R^0$, the Burgers vector is in general not consistent with the underlying rigidly rotated lattice, providing then a physically unacceptable configuration. This suggests to incorporate

in the model some extra compatibility condition between the Burgers vectors and the admissible strains in order to prevent such unphysical configurations. In this respect which options are best suited is in fact still questionable, and a canonical choice seems to be missing in the present literature. The main reason is that, in contrast with purely linear models, here we are trying to combine the linear circulation condition, which is built upon the additive decomposition of the linear strain, together with the nonlinear energetic framework. In this respect, it would be preferable to adopt a multiplicative decomposition of the strain in a plastic, and an elastic part [22, 23], but at the present it is not clear to us how to implement it in a purely discrete framework. Moreover, while in the limit as $\varepsilon \rightarrow 0$ we obtain a fixed rotation, for positive ε nonlinear deformations make unavailable a clear notion of local orientation for the deformed lattice. In order to overcome this lack of rigidity, in the present formulation we impose that, in the annulus of radii ε and ε^γ (with $\gamma \in (0, 1)$) around each dislocation point, the average of the admissible strain fields coincides with the rotation associated to the corresponding Burgers vector. For positive ε , rigidity arguments show that such a consistency between the orientation of the Burgers vector and the underlying lattice holds, not only in average, but actually pointwise up to a small error. In fact, such an average condition guarantees, in the compactness result, that the limit constant rotation R of the admissible fields β_ε is also the limit of each of the rotations associated with each Burgers vector. Such a condition leads us to work under the *well separation* assumption, namely, to restrict the class of admissible empirical measures to that for which the distance of two dislocations is larger than (a multiple of) ε^γ .

Within the framework described above, we prove that a given sequence $\{\mu_\varepsilon\}_\varepsilon$ of admissible configurations of edge dislocations, satisfying the energy bound $C\varepsilon^2|\log \varepsilon|$, has (once scaled by ε) uniformly bounded mass; therefore, up to subsequences, $\frac{\mu_\varepsilon}{\varepsilon}$ converges (in the weak star topology) to a finite sum $\mu = \sum_k \xi^k \delta_{x^k}$ of \mathbb{R}^2 -weighted Dirac deltas. Furthermore, a sequence of admissible strains $\{\beta_\varepsilon\}_\varepsilon$ compatible with $\{\mu_\varepsilon\}_\varepsilon$, converges, up to subsequences, to a constant rotation R that is compatible with μ , i.e., the weights ξ^k lie in the R -rotated regular triangular lattice. Eventually, we show that $\beta_\varepsilon \sim R + \varepsilon \sqrt{|\log \varepsilon|} \beta^{\text{cf}}$, for some far-field linear strain β^{cf} , which is curl-free. This is the content of Theorem 1.1. The corresponding Γ -convergence result, Theorem 1.7, shows that the Γ -limit of the energy functional with respect to the convergence described above, is given by the sum of a self-energy plus the linearized (around R) elastic energy of the limit field β^{cf} . The self-energy for such a nonlinear model takes the form of $\sum_k \varphi(R^T \xi^k)$, where φ is a positively 1-homogeneous function that is obtained by a cell-formula through a relaxation procedure accounting for the underlying lattice structure. We highlight that such a density φ depends only on the corresponding linearized elasticity tensor and on the lattice structure and hence it coincides with the one computed in linear models [16]. Loosely speaking, our result shows that the nonlinear elastic energy outside of the cores in the energy regime $\varepsilon^2|\log \varepsilon|$ can be linearized thus obtaining the same Γ -limit as in the corresponding linear model.

Some comments are in order. The proof of our result follows the lines of that in [26], where the authors focus on a finite system of fixed (i.e., independent of ε) edge dislocations, whose Burgers vectors lie in the unrotated lattice. As mentioned above, our varying measures satisfy a uniform bound on the total variation; therefore, from the point of view of compactness properties, our case does not differ much from the case of a fixed system of singularities. Nevertheless, the situation slightly changes when looking at the Γ -limit of the elastic energy. Indeed, whereas for a fixed measure no relaxation takes place in the computation of the self-energy, in our model two or more different singularities for positive ε may converge to the same limiting singularity, thus lowering the value of the self-energy. This means that our Γ -limit could be smaller than that computed in [26]. Furthermore, we stress that the average condition on the strain fields described above is not present in [26], where a possible discrepancy between the (limit) rotation around which the linearization is performed and the rotation (that is actually the identity matrix) of the lattice where the Burgers vectors lie, may occur. However, as explained in [20] (see also [10]), such a gap is not unphysical if one interprets Ω as the deformed configuration instead of the reference one. Having in mind such an interpretation, an admissible strain field β can be locally understood as an inverse deformation gradient, so that its circulation turns out to be a vector of the reference lattice (without any rotation). In fact, while it is quite accepted that the Burgers circuit should be drawn in the deformed configuration, the correct notion of Burgers vector is rather questionable. The one corresponding to the circulation of the (local) inverse of the deformation gradient on a closed path in the deformed configuration is usually referred to as the *true* Burgers vector [18, formula (1-5)] and, as already explained, it has the advantage to have a topological nature, being quantized on the reference lattice, without being effected by any sort of

small elastic fluctuations. On the other hand, the so-called *local* Burgers vector, defined as the circulation of the deformation gradient on a closed path in the reference configuration, besides being very popular, has a direct constitutive relation with the deformation field. Our proposal of notion of Burgers vector represents a kind of compromise between the two, being defined on the reference configuration but belonging to a rotation of the lattice. In fact, such a procedure is borrowed from linear models [6, 9, 14, 16], where the (local) Burgers vector is (somehow tacitly [18, p. 21]) assumed to belong to the (unrotated) reference lattice. Adopting the notion of true Burgers vector within a purely discrete model seems to be a challenging task; in this respect, it would be interesting (also in the linear framework) to introduce discrete models defined directly on the deformed configuration, combining the discrete formalism of [19] with the analysis in [10].

Summarizing, our model is meant to be a first attempt to combine sound and efficient techniques available in the linear framework together with fundamental discrete nonlinear models. Specifically, it relies on the presence of a reference configuration, it is implicitly based on an additive decomposition of the strain, although the energetic setting is nonlinear. The resulting modeling choice has a flavor of compromise: we have shown that canonical arguments in linear elasticity theories of dislocations can be pushed to and exploited in a nonlinear framework as well, at the price of some extra care needed to relieve the conflict generated by such an unusual combination.

Our contribution represents a first attempt to rigorously analyze in terms of Γ -convergence a purely discrete nonlinear model for edge dislocations in two dimensions.

Notation. Let X be an Euclidean space. For every $\rho > 0$ and for every $x \in X$, $B_\rho(x)$ denotes the open ball centered at x with radius equal to ρ (with respect to the Euclidean metric). Furthermore, for every $x \in X$ and for every $0 < r < R$, the symbol $A_{r,R}(x)$ denotes the (open) annulus $B_R(x) \setminus \overline{B_r}(x)$. In what follows, $\{e_1, e_2\}$ denotes the canonical basis of \mathbb{R}^2 , i.e., $e_1 = (1 \ 0)^T$ and $e_2 = (0 \ 1)^T$. Moreover, $\text{SO}(2)$ is the set of rotations of \mathbb{R}^2 , i.e., the set of matrices $R \in \mathbb{R}^{2 \times 2}$ such that $R^T R = \text{Id}$ and $\det R = 1$. Finally, throughout the paper the symbol C denotes a positive constant that may change from line to line. Whenever we want to stress the dependence of C on some parameters p_1, \dots, p_J (with $J \in \mathbb{N}$) we write C_{p_1, \dots, p_J} .

1 The semi-discrete model

In this section we introduce and analyze the nonlinear semi-discrete model for the elastic energy induced by a family of edge dislocations.

1.1 Description of the problem

Here we introduce the main notations that will be used throughout the paper. Let $\Omega \subset \mathbb{R}^2$ be a bounded, open, and simply connected set with Lipschitz continuous boundary. Let $S \subset \mathbb{R}^2$ be a finite subset of \mathbb{R}^2 and set $\mathbb{S} := \text{span}_{\mathbb{Z}} S$. Here \mathbb{S} represents a reference underlying lattice. We set

$$\Pi(\mathbb{S}) := \bigcup_{R \in \text{SO}(2)} R\mathbb{S}$$

and

$$\mathcal{J}(\mathbb{S}) := \{R \in \text{SO}(2) : R\mathbb{S} = \mathbb{S}\}.$$

Let $0 < \gamma < 1$. Denoting by $\mathcal{M}(\mathbb{R}^2; \mathbb{R}^2)$ the class of \mathbb{R}^2 -valued Radon measures on \mathbb{R}^2 , for every $\varepsilon > 0$ (which is a parameter tuning the lattice spacing) we define the class of admissible edge dislocation measures $X_\varepsilon^\gamma(\Omega)$ as

$$X_\varepsilon^\gamma(\Omega) := \left\{ \mu \in \mathcal{M}(\mathbb{R}^2; \mathbb{R}^2) : \mu = \varepsilon \sum_{n=1}^N \xi^n \delta_{x^n}, N \in \mathbb{N}, x^n \in \Omega, \xi^n \in \Pi(\mathbb{S}), \right. \\ \left. |x^{n_1} - x^{n_2}| \geq 4\varepsilon^\gamma \text{ whenever } n_1 \neq n_2, \text{dist}(x^n, \partial\Omega) \geq 2\varepsilon^\gamma \text{ for all } n = 1, \dots, N \right\}. \quad (1.1)$$

Moreover, for every $R \in \text{SO}(2)$, we set

$$\bar{X}^R(\Omega) := \left\{ \mu \in \mathcal{M}(\mathbb{R}^2; \mathbb{R}^2) : \mu = \sum_{k=1}^K \xi^k \delta_{x^k}, K \in \mathbb{N}, x^k \in \Omega, \xi^k \in R\mathbb{S} \right\}.$$

For every $0 < \varepsilon < 1$ and for every $\mu = \varepsilon \sum_{n=1}^N R^n b^n \delta_{x^n} \in X_\varepsilon^\gamma(\Omega)$, with $b^n \in \mathbb{S}$ and $R^n \in \text{SO}(2)$, we set

$$\Omega_\varepsilon(\mu) := \Omega \setminus \bigcup_{n=1}^N \bar{B}_\varepsilon(x^n)$$

and we define the class of *admissible strains* associated to μ as

$$\begin{aligned} \mathcal{AS}_\varepsilon^\gamma(\mu) := \left\{ \beta \in L^2(\Omega; \mathbb{R}^{2 \times 2}) : \beta \equiv 0 \text{ in } \bigcup_{x \in \text{supp } \mu} B_\varepsilon(x), \text{Curl } \beta = 0 \text{ in } \Omega_\varepsilon(\mu), \int_{A_{\varepsilon, \varepsilon^\gamma}(x^n)} \beta \, dx \in R^n \mathcal{J}(\mathbb{S}), \right. \\ \left. \int_{\partial B_\varepsilon(x^n)} \beta t \, d\mathcal{H}^1 = \varepsilon R^n b^n \text{ for every } n = 1, \dots, N \right\}, \end{aligned} \quad (1.2)$$

where, here and below, the operator Curl acts row-wise and is understood in the distributional sense, t denotes the tangent vector field to $\partial B_\varepsilon(x^n)$ and the integrand βt is intended in the sense of traces (see, for instance, [12, Theorem 2, p. 204]). Notice that the weight of any admissible singularity cannot be uniquely decomposed into a product Rb with $R \in \text{SO}(2)$ and $b \in \mathbb{S}$ (as an example, if $\mathbb{S} = \mathbb{Z}^2$, then e_1 can be written also as the clockwise $\frac{\pi}{2}$ -rotation of e_2); this is why the condition on the average of β in (1.2) is written in terms of an inclusion (into $R^n \mathcal{J}(\mathbb{S})$) rather than an identity.

We now introduce the nonlinear energy density. Let $W \in C^0(\mathbb{R}^{2 \times 2}; [0, +\infty))$ satisfy the following assumptions:

$$W(\text{Id}) = 0, \quad (\text{i})$$

$$W(RM) = W(M) \text{ for every } M \in \mathbb{R}^{2 \times 2}, R \in \text{SO}(2), \quad (\text{ii})$$

$$W(MR) = W(M) \text{ for every } M \in \mathbb{R}^{2 \times 2}, R \in \mathcal{J}(\mathbb{S}), \quad (\text{iii})$$

$$\text{there exists a constant } c > 0 \text{ such that: } W(M) \geq c \, \text{dist}^2(M, \text{SO}(2)) \text{ for every } M \in \mathbb{R}^{2 \times 2}, \quad (\text{iv})$$

$$W \in C^2 \text{ in a neighborhood of } \text{SO}(2). \quad (\text{v})$$

Condition (i) states that the reference configuration is stress-free and condition (ii) is the so-called *frame indifference*. Condition (iii) says that the energy density W is invariant with respect to the symmetry group of rotations of the underlying lattice (while, if assumed for all rotations, would be an isotropy condition). Condition (iv) is a coercivity assumption that guarantees compactness properties whereas condition (v) serves to linearize the energy around the equilibria. We highlight that, combining conditions (i), (ii) and (iv), the well of W is $\text{SO}(2)$.

In view of property (v), we can define the $2 \times 2 \times 2 \times 2$ linearized elasticity tensor associated to W as

$$\mathbb{C} := \frac{\partial^2 W}{\partial M^2}(\text{Id}). \quad (1.3)$$

For every $0 < \varepsilon < 1$ we define the energy functional $\mathcal{F}_\varepsilon^\gamma : \mathcal{M}(\mathbb{R}^2; \mathbb{R}^2) \times L^2(\Omega; \mathbb{R}^{2 \times 2}) \rightarrow [0, +\infty]$ as

$$\mathcal{F}_\varepsilon^\gamma(\mu, \beta) := \begin{cases} \int_{\Omega_\varepsilon(\mu)} W(\beta) \, dx & \text{if } \mu \in X_\varepsilon^\gamma(\Omega) \text{ and } \beta \in \mathcal{AS}_\varepsilon^\gamma(\mu), \\ +\infty & \text{otherwise.} \end{cases} \quad (1.4)$$

1.2 Compactness

Here we state and prove the compactness result for the semi-discrete energy in (1.4).

Theorem 1.1. *Let $\{(\mu_\varepsilon; \beta_\varepsilon)\}_\varepsilon \subset \mathcal{M}(\mathbb{R}^2; \mathbb{R}^2) \times L^2(\Omega; \mathbb{R}^{2 \times 2})$ be such that*

$$\mathcal{F}_\varepsilon^\gamma(\mu_\varepsilon, \beta_\varepsilon) \leq C\varepsilon^2 |\log \varepsilon| \quad \text{for every } \varepsilon > 0, \quad (1.5)$$

for some constant $C > 0$. Then, up to a subsequence, there exist $\{R_\varepsilon\}_\varepsilon \subset \text{SO}(2)$, a rotation $R \in \text{SO}(2)$, a measure $\mu \in \overline{X}^R(\Omega)$, and a field $\beta^{\text{cf}} \in L^2(\Omega; \mathbb{R}^{2 \times 2})$ with $\text{Curl } \beta^{\text{cf}} = 0$, such that:

$$R_\varepsilon \rightarrow R, \quad (0)$$

$$\frac{\mu_\varepsilon}{\varepsilon} \xrightarrow{*} \mu, \quad (1)$$

$$\frac{\beta_\varepsilon - R_\varepsilon}{\varepsilon \sqrt{|\log \varepsilon|}} \rightharpoonup \beta^{\text{cf}} \quad \text{in } L^2(\Omega; \mathbb{R}^{2 \times 2}). \quad (2)$$

Definition 1.2. In the following, we write $(\mu_\varepsilon; \beta_\varepsilon; R_\varepsilon) \rightarrow (\mu; \beta^{\text{cf}}; R)$ if conditions (0), (1), and (2) of Theorem 1.1 are satisfied.

In order to prove Theorem 1.1 we follow the lines of the proof of [26, Proposition 3.5]. The following result provides uniform rigidity estimates *à la* Friesecke, James and Müller [15] for a suitable class of domains with holes. It can be proven by arguing verbatim as in [26, Lemma 3.1].

Lemma 1.3. *Let $E \subset \mathbb{R}^2$ be a bounded, open set with Lipschitz continuous boundary. Then there exists a constant $C > 0$ depending only on E such that the following property holds true. Let $r > 0$ and let $\{x^1, \dots, x^N\} \subset E$ be such that $|x^{n_1} - x^{n_2}| > 4r$ for every $n_1, n_2 = 1, \dots, N$ with $n_1 \neq n_2$ and $\text{dist}(x^n, \partial E) > 2r$ for every $n = 1, \dots, N$. Setting $E_r := E \setminus \bigcup_{n=1}^N \overline{B}_r(x^n)$, for every $u \in H^1(E_r; \mathbb{R}^2)$ there exists $R \in \text{SO}(2)$ such that*

$$\|\nabla u - R\|_{L^2(E_r; \mathbb{R}^{2 \times 2})} \leq C \|\text{dist}(\nabla u, \text{SO}(2))\|_{L^2(E_r)}.$$

Proof of Theorem 1.1. By the energetic bound (1.5) we have that $\mu_\varepsilon \in X_\varepsilon^\gamma(\Omega)$ and $\beta_\varepsilon \in \mathcal{AS}_\varepsilon^\gamma(\mu_\varepsilon)$. For every $0 < \varepsilon < 1$ let

$$\mu_\varepsilon := \varepsilon \sum_{n=1}^{N_\varepsilon} R_\varepsilon^n b_\varepsilon^n \delta_{x_\varepsilon^n}.$$

We start by proving the compactness property for the measures $\frac{\mu_\varepsilon}{\varepsilon}$. We set

$$K_\varepsilon^\gamma := \left[(1 - \gamma) \frac{|\log \varepsilon|}{\log 2} - 1 \right].$$

Let $n \in \{1, \dots, N_\varepsilon\}$ be fixed. We have that $A_{\varepsilon, \varepsilon^\gamma}(x_\varepsilon^n) \supset \bigcup_{k=0}^{K_\varepsilon^\gamma} A_{2^k \varepsilon, 2^{k+1} \varepsilon}(x_\varepsilon^n)$. As a consequence of [15], there exists a constant $C > 0$ such that for every $k = 0, 1, \dots, K_\varepsilon^\gamma$,

$$\int_{A_{2^k \varepsilon, 2^{k+1} \varepsilon}(x_\varepsilon^n)} \text{dist}^2(\beta_\varepsilon, \text{SO}(2)) dx \geq C \int_{A_{2^k \varepsilon, 2^{k+1} \varepsilon}(x_\varepsilon^n)} |\beta_\varepsilon - R_k|^2 dx \quad (1.6)$$

for some rotations R_k ; in fact, (1.6) has been proved in [15] for gradient fields defined on sets with Lipschitz continuous boundary; moreover, the constant C is invariant with respect to homotheties of the domain. Actually, in our case β_ε is not a gradient, but one can still deduce (1.6) either covering $A_{2^k \varepsilon, 2^{k+1} \varepsilon}(x_\varepsilon^n)$ with two overlapping simply connected sets with Lipschitz continuous boundary, and applying on each of them [15], or introducing a suitable cut in $A_{2^k \varepsilon, 2^{k+1} \varepsilon}(x_\varepsilon^n)$ and applying directly [15] on the resulting simply connected cut annulus. By summing (1.6) over k and over n , using (1.5) and property (iv), we deduce that

$$\begin{aligned} C \varepsilon^2 |\log \varepsilon| &\geq C \sum_{n=1}^{N_\varepsilon} \sum_{k=0}^{K_\varepsilon^\gamma} \int_{A_{2^k \varepsilon, 2^{k+1} \varepsilon}(x_\varepsilon^n)} |\beta_\varepsilon - R_k|^2 dx \\ &\geq C \sum_{n=1}^{N_\varepsilon} \varepsilon^2 |b_\varepsilon^n|^2 \frac{1}{2\pi} (K_\varepsilon^\gamma + 1) \log 2 \geq C \varepsilon^2 |\log \varepsilon| (1 - \gamma) \left| \frac{\mu_\varepsilon}{\varepsilon} \right|(\Omega), \end{aligned} \quad (1.7)$$

where the last but one inequality follows by applying Fubini's Theorem, Jensen's inequality and using the very definition of $\mathcal{AS}_\varepsilon^\gamma(\mu_\varepsilon)$ combined with Stokes' Theorem. By (1.7), we deduce that $\left| \frac{\mu_\varepsilon}{\varepsilon} \right|(\Omega)$ is uniformly bounded and hence that there exists a measure μ , consisting in a finite sum of vectorial weighted Dirac deltas, such that, up to subsequences, property (1) holds.

Now, we pass to the proof of property (2). To this end, we define the maps $\bar{\beta}_\varepsilon : \Omega \rightarrow \mathbb{R}^{2 \times 2}$ as

$$\bar{\beta}_\varepsilon(x) := \varepsilon \chi_{\Omega_\varepsilon(\mu_\varepsilon)}(x) \sum_{n=1}^{N_\varepsilon} (R_\varepsilon^n b_\varepsilon^n) \otimes \mathbb{J} \frac{x - x_\varepsilon^n}{|x - x_\varepsilon^n|^2},$$

where \mathbb{J} is the counter clockwise $\frac{\pi}{2}$ -rotation. One can check that

$$\int_{\Omega_\varepsilon(\mu_\varepsilon)} |\bar{\beta}_\varepsilon|^2 dx \leq C \varepsilon^2 N_\varepsilon \sum_{n=1}^{N_\varepsilon} |b_\varepsilon^n|^2 |\log \varepsilon| \leq C_\gamma \varepsilon^2 |\log \varepsilon|, \quad (1.8)$$

where the last inequality follows by (1.7). By construction, we have that $\int_{\partial U} (\beta_\varepsilon - \bar{\beta}_\varepsilon) t \, d\mathcal{J}^1 = 0$ for every open set $U \subset \Omega$ with $\partial U \subset \Omega_\varepsilon(\mu_\varepsilon)$ smooth; hence, there exists a map $u_\varepsilon \in H^1(\Omega_\varepsilon(\mu_\varepsilon); \mathbb{R}^2)$ such that $\beta_\varepsilon - \bar{\beta}_\varepsilon = \nabla u_\varepsilon$ in $\Omega_\varepsilon(\mu_\varepsilon)$. By Lemma 1.3 we thus get that there exist a constant $C > 0$ independent of ε and a sequence of rotations $\{R_\varepsilon\}_\varepsilon \subset \text{SO}(2)$ such that

$$\begin{aligned} \int_{\Omega} |\beta_\varepsilon - R_\varepsilon|^2 dx &\leq 2 \int_{\Omega_\varepsilon(\mu_\varepsilon)} |\beta_\varepsilon - \bar{\beta}_\varepsilon - R_\varepsilon|^2 dx + 2 \int_{\Omega_\varepsilon(\mu_\varepsilon)} |\bar{\beta}_\varepsilon|^2 dx + C_\gamma \varepsilon^2 \\ &\leq 2 \int_{\Omega_\varepsilon(\mu_\varepsilon)} |\nabla u_\varepsilon - R_\varepsilon|^2 dx + C_\gamma \varepsilon^2 |\log \varepsilon| \\ &\leq C \int_{\Omega_\varepsilon(\mu_\varepsilon)} \text{dist}^2(\nabla u_\varepsilon, \text{SO}(2)) dx + C_\gamma \varepsilon^2 |\log \varepsilon| \\ &\leq C \int_{\Omega_\varepsilon(\mu_\varepsilon)} \text{dist}^2(\beta_\varepsilon, \text{SO}(2)) dx + C_\gamma \varepsilon^2 |\log \varepsilon| \\ &\leq C_\gamma \varepsilon^2 |\log \varepsilon|, \end{aligned} \quad (1.9)$$

where we have used also (1.7), (1.8), property (iv), and (1.5). By (1.9), up to a subsequence, we get $\frac{\beta_\varepsilon - R_\varepsilon}{\varepsilon \sqrt{|\log \varepsilon|}} \rightharpoonup \beta^{\text{cf}}$ in $L^2(\Omega; \mathbb{R}^{2 \times 2})$ for some $\beta^{\text{cf}} \in L^2(\Omega; \mathbb{R}^{2 \times 2})$, so that property (2) holds true.

Now we prove that $\text{Curl } \beta^{\text{cf}} = 0$ in the distributional sense. To this end, let $\phi \in C_0^1(\Omega)$ and let $\{\phi_\varepsilon\}_\varepsilon \subset H_0^1(\Omega)$ be a family converging to ϕ strongly in $H_0^1(\Omega)$ and such that $\phi_\varepsilon \equiv \phi(x_\varepsilon^n)$ in $B_\varepsilon(x_\varepsilon^n)$ for every $n = 1, \dots, N_\varepsilon$. Then, by (1.7), we get

$$\begin{aligned} \langle \text{Curl } \beta^{\text{cf}}, \phi \rangle &= \lim_{\varepsilon \rightarrow 0} \frac{1}{\sqrt{|\log \varepsilon|}} \left\langle \text{Curl } \frac{\beta_\varepsilon - R_\varepsilon}{\varepsilon}, \phi_\varepsilon \right\rangle = \lim_{\varepsilon \rightarrow 0} \frac{1}{\sqrt{|\log \varepsilon|}} \left\langle \text{Curl } \frac{\beta_\varepsilon}{\varepsilon}, \phi_\varepsilon \right\rangle \\ &= \lim_{\varepsilon \rightarrow 0} \frac{\sum_{n=1}^{N_\varepsilon} R_\varepsilon^n b_\varepsilon^n \phi(x_\varepsilon^n)}{\sqrt{|\log \varepsilon|}} = 0; \end{aligned}$$

by the arbitrariness of ϕ we have that $\text{Curl } \beta^{\text{cf}} = 0$ in the distributional sense.

Furthermore, since $\text{SO}(2)$ is compact, up to a subsequence $R_\varepsilon \rightarrow R$ for some $R \in \text{SO}(2)$. In order to conclude the proof of the theorem it remains to show that $\mu \in \bar{X}^R(\Omega)$. First, for every $n = 1, \dots, N_\varepsilon$, let $\tilde{R}_\varepsilon^n \in \mathcal{J}(\mathbb{S})$ be such that

$$\int_{A_{\varepsilon, \varepsilon^\gamma}(x_\varepsilon^n)} \beta_\varepsilon dx = R_\varepsilon^n \tilde{R}_\varepsilon^n.$$

In view of (1.7), we can assume, up to a subsequence, that $N_\varepsilon \equiv N$ for some $N \in \mathbb{N}$ independent of ε . For every fixed $n = 1, \dots, N$, it holds

$$|R_\varepsilon^n \tilde{R}_\varepsilon^n - R_\varepsilon|^2 = \left| \int_{A_{\varepsilon, \varepsilon^\gamma}(x_\varepsilon^n)} (\beta_\varepsilon - R_\varepsilon) dx \right|^2 \leq C_\gamma \varepsilon^{2(1-\gamma)} |\log \varepsilon|, \quad (1.10)$$

where the last inequality is a consequence of Jensen inequality and of (1.9). Since $\mathcal{J}(\mathbb{S})$ is compact, we have that, up to a further subsequence, $\tilde{R}_\varepsilon^n \rightarrow \tilde{R}^n$ for some $\tilde{R}^n \in \mathcal{J}(\mathbb{S})$. This fact together with (1.10), yields that $R_\varepsilon^n \rightarrow R(\tilde{R}^n)^T$ as $\varepsilon \rightarrow 0$ and hence $\mu \in \bar{X}^R(\Omega)$. \square

Remark 1.4. In this paper, the assumption that Ω is simply connected is used to guarantee that the class of L^2 curl-free matrix fields on Ω coincides with the class of gradients of functions in $H^1(\Omega; \mathbb{R}^2)$. Such an assumption can be dropped if we replace, in the compactness result above and in the Γ -convergence result below, the condition $\beta^{\text{cf}} \in L^2(\Omega; \mathbb{R}^{2 \times 2})$ with $\text{Curl } \beta^{\text{cf}} = 0$ by the condition $\beta^{\text{cf}} = \nabla u$ for some $u \in H^1(\Omega; \mathbb{R}^2)$.

Remark 1.5. As mentioned in the introduction, the condition on the average of the strain field in (1.2) is needed in Theorem 1.1 to guarantee the compatibility condition that R is both the asymptotic rotation around which the linearization is performed and the rotation of the lattice where the weights of the limit singularities lie. Indeed, dropping such an average condition from the definition of (1.2), one could consider $\mu_\varepsilon \equiv \varepsilon b \delta_0$ (with $0 \in \Omega$ and $b \in \mathbb{S}$), R an arbitrary rotation in $\text{SO}(2)$ and the field $\beta_\varepsilon = R + \varepsilon \beta_{\mathbb{R}^2}^{b, \mathbb{C}}$ (where $\beta_{\mathbb{R}^2}^{\zeta, \mathbb{C}}$ is the Green function for the corresponding linearized problem, see (1.11) below); such a field (once dropped the average condition of (1.2)) would be an admissible strain. Now, the weight of the singularity lies on the set \mathbb{S} , while β_ε is linearized around the arbitrarily fixed rotation R , providing an unphysical lack of rigidity in the class of admissible Burgers vectors.

Furthermore, we stress that such an average condition should be required on “thick annuli” with radii $\varepsilon, \varepsilon^\gamma$ and cannot be replaced by a weaker condition on “thin” annuli with radii $\varepsilon, M\varepsilon$ for some $M > 1$ (independent of ε), as shown by the following computation. Let $\vartheta_\varepsilon : [\varepsilon, 1] \rightarrow [0, 1]$ be such that $\vartheta_\varepsilon(t) \equiv 0$ in $[\varepsilon, M\varepsilon]$, $\vartheta_\varepsilon(t) \equiv 1$ in $[2M\varepsilon, 1]$ and $\vartheta_\varepsilon(t) = \frac{t}{M\varepsilon} - 1$ in $[M\varepsilon, 2M\varepsilon]$ and let $u_\varepsilon : A_{\varepsilon, 1}(0) \rightarrow \mathbb{R}^2$ be the function defined by $u_\varepsilon(x) := R(\vartheta_\varepsilon(|x|))x$, where

$$R(\vartheta) := \begin{pmatrix} \cos \vartheta & \sin \vartheta \\ -\sin \vartheta & \cos \vartheta \end{pmatrix}.$$

By direct computations,

$$\nabla u_\varepsilon(x) = R(\vartheta_\varepsilon(|x|)) + \frac{\vartheta'_\varepsilon(|x|)}{|x|} \begin{pmatrix} -\sin \vartheta_\varepsilon(|x|)x_1^2 + \cos \vartheta_\varepsilon(|x|)x_1x_2 & -\sin \vartheta_\varepsilon(|x|)x_1x_2 + \cos \vartheta_\varepsilon(|x|)x_2^2 \\ -\cos \vartheta_\varepsilon(|x|)x_1^2 - \sin \vartheta_\varepsilon(|x|)x_1x_2 & -\sin \vartheta_\varepsilon(|x|)x_2^2 - \cos \vartheta_\varepsilon(|x|)x_1x_2 \end{pmatrix},$$

whence we deduce that

$$\int_{A_{\varepsilon, 1}(0)} \text{dist}^2(\nabla u_\varepsilon, \text{SO}(2)) \, dx \leq \frac{C}{M^2 \varepsilon^2} \int_{A_{M\varepsilon, 2M\varepsilon}(0)} |x|^2 \, dx \leq CM^2 \varepsilon^2.$$

In this example, the average of ∇u_ε on the annulus $A_{\varepsilon, M\varepsilon}(0)$ is the identity matrix $R(0)$, while $\nabla u_\varepsilon \rightarrow R(1)$ (for instance in $L^2(\Omega; \mathbb{R}^{2 \times 2})$). The computation above shows that the fields $\beta_\varepsilon = R(\vartheta_\varepsilon) + \varepsilon \beta_{\mathbb{R}^2}^{b, \mathbb{C}}$ (with $b \in \mathbb{S}$), would satisfy the average assumption with $R = R(0) = \text{Id}$ in the annulus $A_{\varepsilon, M\varepsilon}(0)$, but converge to the rotation $R(1)$. Therefore, prescribing the condition on the average of the admissible fields on the annuli of radii ε and $M\varepsilon$ (for M fixed) does not provide any relationship between the rotation of the lattice where the Burgers vector lies and the rotation around which the energy is linearized.

On the other hand, in order to let our proof (in particular, (1.10)) work, one can see that the average condition in (1.2) can be weakened by requiring that

$$\text{dist} \left(\int_{A_{\varepsilon, \varepsilon^\gamma}(x^n)} \beta \, dx, R^n \mathcal{J}(\mathbb{S}) \right) \leq \delta_\varepsilon,$$

for some $\delta_\varepsilon > 0$ with $\delta_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$.

1.3 Γ -convergence

This subsection is devoted to the Γ -convergence result for the semi-discrete energy in (1.4). In order to define the Γ -limit we first introduce the self-energy of an edge dislocation. Let \mathbb{C} be a given elasticity tensor in linear elasticity. For every $\zeta \in \mathbb{R}^2 \setminus \{0\}$, the corresponding displacement $u_{\mathbb{R}^2}^{\zeta, \mathbb{C}}$ and strain $\beta_{\mathbb{R}^2}^{\zeta, \mathbb{C}}$, induced by the edge dislocation $\zeta \delta_0$ in the whole plane, have been explicitly computed in the literature (see, for instance, [7, formula (4.1.25)]); here, we recall some of their properties we need in our analysis. The strain field $\beta_{\mathbb{R}^2}^{\zeta, \mathbb{C}}$ satisfies the circulation condition

$$\text{Curl } \beta = \zeta \delta_0 \quad \text{in } \mathbb{R}^2$$

and the equilibrium equation

$$\operatorname{Div} \mathbf{C}\beta = 0 \quad \text{in } \mathbb{R}^2.$$

In polar coordinates, $\beta_{\mathbb{R}^2}^{\zeta, \mathbf{C}}$ takes the form

$$\beta_{\mathbb{R}^2}^{\zeta, \mathbf{C}}(\rho, \theta) := \frac{1}{\rho} \Gamma^{\zeta, \mathbf{C}}(\theta), \quad (1.11)$$

where (see [16, Remark 7]) $\Gamma^{\zeta, \mathbf{C}}$ is the unique minimizer of $\int_0^{2\pi} \mathbf{C}\Gamma : \Gamma \, d\theta$ among the functions Γ of the form

$$\Gamma(\theta) := f(\theta) \otimes (-\sin \theta; \cos \theta) + g \otimes (\cos \theta; \sin \theta),$$

with $g \in \mathbb{R}^2$ and $f \in C^0([0, 2\pi]; \mathbb{R}^2)$ satisfying $f(0) = f(2\pi)$ and $\int_0^{2\pi} f(\omega) \, d\omega = \zeta$. The optimal $g^{\zeta, \mathbf{C}}$ and $f^{\zeta, \mathbf{C}}$ are uniquely determined by the vector ζ and the tensor \mathbf{C} .

Remark 1.6. We highlight that $\Gamma^{\zeta, \mathbf{C}} \in C^\infty(\mathbb{R})$ (see [7, formula (4.1.25)]). Moreover,

$$\int_0^{2\pi} \Gamma^{\zeta, \mathbf{C}}(\theta) \, d\theta = 0, \quad (1.12)$$

and hence, for all $0 < r < R$,

$$\int_{A_{r,R}(0)} \beta_{\mathbb{R}^2}^{\zeta, \mathbf{C}} \, dx = 0.$$

Indeed, let $\tilde{f}^{\zeta, \mathbf{C}}$ be the 2π -periodic extension of $f^{\zeta, \mathbf{C}}$ and consider

$$\hat{\Gamma}^{\zeta, \mathbf{C}}(\theta) := -\Gamma^{\zeta, \mathbf{C}}(\theta + \pi) = \tilde{f}^{\zeta, \mathbf{C}}(\theta + \pi) \otimes (-\sin \theta; \cos \theta) + g^{\zeta, \mathbf{C}} \otimes (\cos \theta; \sin \theta).$$

Notice that the circulation of $\hat{\Gamma}^{\zeta, \mathbf{C}}$ coincides with that of $\Gamma^{\zeta, \mathbf{C}}$. Since, by convexity,

$$\int_0^{2\pi} \mathbf{C} \left(\frac{1}{2} \Gamma^{\zeta, \mathbf{C}} + \frac{1}{2} \hat{\Gamma}^{\zeta, \mathbf{C}} \right) : \left(\frac{1}{2} \Gamma^{\zeta, \mathbf{C}} + \frac{1}{2} \hat{\Gamma}^{\zeta, \mathbf{C}} \right) \, d\theta \leq \frac{1}{2} \int_0^{2\pi} \mathbf{C} \Gamma^{\zeta, \mathbf{C}} : \Gamma^{\zeta, \mathbf{C}} \, d\theta + \frac{1}{2} \int_0^{2\pi} \mathbf{C} \hat{\Gamma}^{\zeta, \mathbf{C}} : \hat{\Gamma}^{\zeta, \mathbf{C}} \, d\theta = \int_0^{2\pi} \mathbf{C} \Gamma^{\zeta, \mathbf{C}} : \Gamma^{\zeta, \mathbf{C}} \, d\theta,$$

by strict convexity we deduce that

$$\Gamma^{\zeta, \mathbf{C}}(\cdot) = \hat{\Gamma}^{\zeta, \mathbf{C}}(\cdot) = -\Gamma^{\zeta, \mathbf{C}}(\cdot + \pi); \quad (1.13)$$

therefore, $\Gamma^{\zeta, \mathbf{C}}$ is an odd function and hence (1.12) holds true.

For every $\zeta \in \mathbb{R}^2 \setminus \{0\}$ we set

$$\psi^{\mathbf{C}}(\zeta) := \int_0^{2\pi} \frac{1}{2} \mathbf{C} \Gamma^{\zeta, \mathbf{C}}(\theta) : \Gamma^{\zeta, \mathbf{C}}(\theta) \, d\theta = \frac{1}{|\log r|} \int_{A_{r,1}(0)} \frac{1}{2} \mathbf{C} \beta_{\mathbb{R}^2}^{\zeta, \mathbf{C}} : \beta_{\mathbb{R}^2}^{\zeta, \mathbf{C}} \, dx, \quad 0 < r < 1. \quad (1.14)$$

By [14, Proposition 1] (see also [16, Corollary 6]), we have

$$\psi^{\mathbf{C}}(\zeta) = \lim_{\frac{r_2}{r_1} \rightarrow +\infty} \psi_{r_1, r_2}^{\mathbf{C}}(\zeta), \quad (1.15)$$

where

$$\psi_{r_1, r_2}^{\mathbf{C}}(\zeta) := \frac{1}{\log \frac{r_2}{r_1}} \min_{\substack{\beta \in L^2(A_{r_1, r_2}(0); \mathbb{R}^{2 \times 2}) \\ \operatorname{Curl} \beta = 0 \text{ in } A_{r_1, r_2}(0) \\ \int_{\partial B_{r_1}(0)} \beta t \, d\mathcal{H}^1 = \zeta}} \int \frac{1}{2} \mathbf{C} \beta : \beta \, dx. \quad (1.16)$$

Moreover, for every $\zeta \in \mathbb{R}^2$ and for every $s \in \mathbb{R}$,

$$\psi^{\mathbf{C}}(s\zeta) = s^2 \psi^{\mathbf{C}}(\zeta). \quad (1.17)$$

Finally, for every $b \in \mathbb{S}$ we define

$$\varphi^{\mathbf{C}}(b) := \min \left\{ \sum_{i=1}^I |z_i| \psi^{\mathbf{C}}(b_i) : z_i \in \mathbb{Z}, b_i \in \mathbb{S}, I \in \mathbb{N}, \sum_{i=1}^I z_i b_i = b \right\} \quad (1.18)$$

(with $\varphi^{\mathbf{C}}(b) = 0$ for $b = 0$). We are now in a position to state our Γ -convergence result for the nonlinear energy $\mathcal{F}_\varepsilon^{\mathbf{y}}$ defined in (1.4).

Theorem 1.7. *The following Γ -convergence result holds true.*

(i) (Γ -liminf inequality) *Let $R \in \text{SO}(2)$, $\mu = \sum_{k=1}^K \xi^k \delta_{x^k} \in \bar{X}^R(\Omega)$, $\beta^{\text{cf}} \in L^2(\Omega; \mathbb{R}^{2 \times 2})$ with $\text{Curl } \beta^{\text{cf}} = 0$. For every $\{(\mu_\varepsilon; \beta_\varepsilon; R_\varepsilon)\}_\varepsilon \subset \mathcal{M}(\mathbb{R}^2; \mathbb{R}^2) \times L^2(\Omega; \mathbb{R}^{2 \times 2}) \times \text{SO}(2)$ such that $(\mu_\varepsilon; \beta_\varepsilon; R_\varepsilon) \rightarrow (\mu; \beta^{\text{cf}}; R)$ (in the sense of Definition 1.2), it holds*

$$\sum_{k=1}^K \varphi^{\text{C}}(R^T \xi^k) + \int_{\Omega} \frac{1}{2} \text{C} R^T \beta^{\text{cf}} : R^T \beta^{\text{cf}} \, dx \leq \liminf_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \mathcal{F}_\varepsilon^\gamma(\mu_\varepsilon, \beta_\varepsilon). \quad (1.19)$$

(ii) (Γ -limsup inequality) *Let $R \in \text{SO}(2)$, $\mu = \sum_{k=1}^K \xi^k \delta_{x^k} \in \bar{X}^R(\Omega)$ and $\beta^{\text{cf}} \in L^2(\Omega; \mathbb{R}^{2 \times 2})$ with $\text{Curl } \beta^{\text{cf}} = 0$. Then there exists $\{(\mu_\varepsilon; \beta_\varepsilon; R_\varepsilon)\}_\varepsilon \subset \mathcal{M}(\mathbb{R}^2; \mathbb{R}^2) \times L^2(\Omega; \mathbb{R}^{2 \times 2}) \times \text{SO}(2)$ such that $(\mu_\varepsilon; \beta_\varepsilon; R_\varepsilon) \rightarrow (\mu; \beta^{\text{cf}}; R)$ and*

$$\sum_{k=1}^K \varphi^{\text{C}}(R^T \xi^k) + \int_{\Omega} \frac{1}{2} \text{C} R^T \beta^{\text{cf}} : R^T \beta^{\text{cf}} \, dx \geq \limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \mathcal{F}_\varepsilon^\gamma(\mu_\varepsilon, \beta_\varepsilon). \quad (1.20)$$

Before giving the proof of Theorem 1.7 we state and prove the following linearization lemma for curl-free matrix fields, arguing as in the proof of [26, Proposition 3.11]. We mention that this kind of linearization results in presence of defects generalizes the case of gradient fields originally treated in [11], all relying on the Rigidity Estimate in [15].

Lemma 1.8. *Let $r > 1$. For every $\delta > 0$, let $\zeta_\delta \in \mathbb{R}^2$ and $\beta_\delta \in L^2(A_{1,r}(0); \mathbb{R}^{2 \times 2})$ be such that $\int_{\partial B_1(0)} \beta_\delta t \, d\mathcal{H}^1 = \zeta_\delta$ and $\text{Curl } \beta_\delta = 0$ in $A_{1,r}(0)$. Assume that $\beta_\delta \rightarrow R$ in $L^2(A_{1,r}(0); \mathbb{R}^{2 \times 2})$ and $\frac{\zeta_\delta}{\delta} \rightarrow \zeta$ (as $\delta \rightarrow 0$), for some $\zeta \in \mathbb{R}^2$ and $R \in \text{SO}(2)$. Then*

$$\liminf_{\delta \rightarrow 0} \frac{1}{\delta^2} \int_{A_{1,r}(0)} W(\beta_\delta) \, dx \geq \log r \psi_{1,r}^{\text{C}}(R^T \zeta). \quad (1.21)$$

Proof. In order to prove (1.21) we argue by contradiction. Assume that there exists a vanishing sequence $\{\delta_l\}_{l \in \mathbb{N}} \subset (0, +\infty)$ and a constant $\sigma > 0$ such that

$$\frac{1}{\delta_l^2} \int_{A_{1,r}(0)} W(\beta_{\delta_l}) \, dx < \log r \psi_{1,r}^{\text{C}}(R^T \zeta) - \sigma. \quad (1.22)$$

By (1.22), in view of property (iv) and of the Rigidity Estimate in [15] (see [26, Proposition 3.3] for a specific formulation of such a rigidity result suited for curl-free fields on annuli, as well as Lemma 1.3 above), we get that there exist a constant $C > 0$ and a sequence of constant rotations $\{\bar{R}_{\delta_l}\}_{l \in \mathbb{N}} \subset \text{SO}(2)$ such that

$$\|\beta_{\delta_l} - \bar{R}_{\delta_l}\|_{L^2(A_{1,r}(0); \mathbb{R}^{2 \times 2})}^2 \leq C \log r \psi_{1,r}^{\text{C}}(R^T \zeta) \delta_l^2. \quad (1.23)$$

Clearly, by triangular inequality, $\bar{R}_{\delta_l} \rightarrow R$ as $l \rightarrow +\infty$. By assumption, the field $\frac{\bar{R}_{\delta_l}^T \beta_{\delta_l} - \text{Id}}{\delta_l}$ is curl-free, and hence, setting

$$\text{C} := \{(x_1; 0) : 1 \leq x_1 \leq r\},$$

we have that there exists a sequence $\{v_{\delta_l}\}_{l \in \mathbb{N}} \subset H^1(A_{1,r}(0) \setminus \text{C}; \mathbb{R}^2)$ such that

$$\int_{A_{1,r}(0)} v_{\delta_l} \, dx = 0 \quad \text{and} \quad \frac{\bar{R}_{\delta_l}^T \beta_{\delta_l} - \text{Id}}{\delta_l} = \nabla v_{\delta_l} \quad \text{in } A_{1,r}(0) \setminus \text{C}.$$

By construction,

$$[v_{\delta_l}] = \bar{R}_{\delta_l}^T \frac{\zeta_{\delta_l}}{\delta_l} \quad \text{on } \text{C}, \quad (1.24)$$

where $[v]$ denotes the jump of the function v in the sense of traces. By (1.23), we obtain

$$\int_{A_{1,r}(0)} |\nabla v_{\delta_l}|^2 \, dx \leq C \log r \psi_{1,r}^{\text{C}}(R^T \zeta), \quad (1.25)$$

whence we deduce that, up to a subsequence, $v_{\delta_l} \rightarrow v$ in $H^1(A_{1,r}(0) \setminus \text{C}; \mathbb{R}^2)$ for some $v \in H^1(A_{1,r}(0) \setminus \text{C}; \mathbb{R}^2)$; moreover, by (1.24) and recalling that $\bar{R}_{\delta_l} \rightarrow R$, we get

$$[v] = R^T \zeta \quad \text{on } \text{C}. \quad (1.26)$$

We are now in a position to linearize the energy around the identity. To this end, we define the sequence $\{\chi_{\delta_l}\}_{l \in \mathbb{N}}$ of characteristic functions

$$\chi_{\delta_l}(x) := \begin{cases} 1 & \text{if } |\nabla v_{\delta_l}| < \delta_l^{-\alpha}, \\ 0 & \text{otherwise in } A_{1,r}(0) \setminus C, \end{cases}$$

with $0 < \alpha < 1$ arbitrarily fixed. By (1.25) it follows that $\chi_{\delta_l} \rightarrow 1$ in measure and hence we have $\nabla v_{\delta_l} \chi_{\delta_l} \rightarrow \nabla v$ in $L^2(A_{1,r}(0) \setminus C; \mathbb{R}^{2 \times 2})$. By Taylor expansion, using (ii), we have

$$\begin{aligned} \frac{1}{\delta_l^2} \int_{A_{1,r}(0)} W(\beta_{\delta_l}) \, dx &\geq \frac{1}{\delta_l^2} \int_{A_{1,r}(0)} \chi_{\delta_l} W(\bar{R}_{\delta_l}^T \beta_{\delta_l}) \, dx \\ &= \frac{1}{\delta_l^2} \int_{A_{1,r}(0) \setminus C} \chi_{\delta_l} W(\text{Id} + \delta_l \nabla v_{\delta_l}) \, dx \\ &\geq \int_{A_{1,r}(0) \setminus C} \left(\frac{1}{2} \mathbb{C}(\chi_{\delta_l} \nabla v_{\delta_l}) : (\chi_{\delta_l} \nabla v_{\delta_l}) - \frac{\chi_{\delta_l}}{\delta_l^2} \omega(\delta_l |\nabla v_{\delta_l}|) \right) \, dx, \end{aligned} \quad (1.27)$$

with $\frac{\omega(t)}{t^2} \rightarrow 0$ as $t \rightarrow 0^+$.

Notice that

$$\frac{\chi_{\delta_l}}{\delta_l^2} \omega(\delta_l |\nabla v_{\delta_l}|) = |\nabla v_{\delta_l}|^2 \chi_{\delta_l} \frac{\omega(\delta_l |\nabla v_{\delta_l}|)}{(\delta_l |\nabla v_{\delta_l}|)^2},$$

which is the product of a uniformly bounded sequence $(\{|\nabla v_{\delta_l}|^2\}_{l \in \mathbb{N}})$ in $L^1(A_{1,r}(0))$ and a uniformly vanishing sequence in $L^\infty(A_{1,r}(0))$. Therefore, by (1.27) and by lower semi-continuity, we conclude

$$\liminf_{l \rightarrow +\infty} \frac{1}{\delta_l^2} \int_{A_{1,r}(0)} W(\beta_{\delta_l}) \, dx \geq \int_{A_{1,r}(0) \setminus C} \frac{1}{2} \mathbb{C} \nabla v : \nabla v \, dx \geq \log r \psi_{1,r}^{\mathbb{C}}(R^T \zeta), \quad (1.28)$$

where the last inequality follows by the very definition of $\psi_{1,r}^{\mathbb{C}}$ in (1.16) and by (1.26). Since (1.28) contradicts (1.22), we have that (1.21) holds true. \square

With Lemma 1.8 in hand, we are in a position to prove the Γ -lim inf inequality in Theorem 1.7.

Proof of Theorem 1.7 (i). We can assume without loss of generality that (1.5) is satisfied. For every $\varepsilon > 0$ let $\mu_\varepsilon = \varepsilon \sum_{n=1}^{N_\varepsilon} \xi_\varepsilon^n \delta_{x_\varepsilon^n} \in X_\varepsilon^y(\Omega)$. In view of (1.7), we have that $|\frac{\mu_\varepsilon}{\varepsilon}|(\Omega) \leq C_\gamma$, and hence we may assume that, up to a subsequence, $N_\varepsilon = \hat{N}$ for some \hat{N} independent of ε . Furthermore, up to passing to a further subsequence, we may assume that each of the points x_ε^n converges to some point in a finite set $\{y^j\}_{j=1, \dots, J} \subset \bar{\Omega}$ (with $K \leq J \leq \hat{N}$), where $y^j = x^j$ for $j = 1, \dots, K$. Let $0 < \rho < 1$ be such that the balls $B_{2\rho}(y^j)$ are pairwise disjoint and $\bigcup_{k=1}^K B_{2\rho}(x^k) \subset \Omega$. Then, setting

$$\mathcal{E}_\varepsilon^{\text{far}} := \int_{\Omega \setminus \bigcup_{j=1}^J \bar{B}_\rho(y^j)} W(\beta_\varepsilon) \, dx, \quad \mathcal{E}_\varepsilon^k := \int_{\Omega_\varepsilon(\mu_\varepsilon) \cap B_\rho(x^k)} W(\beta_\varepsilon) \, dx \quad (\text{for every } k = 1, \dots, K),$$

we have that

$$\mathcal{F}_\varepsilon^y(\mu_\varepsilon, \beta_\varepsilon) \geq \mathcal{E}_\varepsilon^{\text{far}} + \sum_{k=1}^K \mathcal{E}_\varepsilon^k. \quad (1.29)$$

By arguing verbatim as in the proof of [26, Proposition 3.11], namely, linearizing W around the limit rotation R and using the lower semi-continuity of the elastic energy with respect to the weak convergence in L^2 , one can check that

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \mathcal{E}_\varepsilon^{\text{far}} \geq \int_{\Omega \setminus \bigcup_{j=1}^J \bar{B}_\rho(y^j)} \mathbb{C} R^T \beta^{\text{cf}} : R^T \beta^{\text{cf}} \, dx. \quad (1.30)$$

Now we claim that, for every $k = 1, \dots, K$,

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \mathcal{E}_\varepsilon^k \geq \varphi^{\mathbb{C}}(R^T \xi^k). \quad (1.31)$$

Notice that (1.29), (1.30) and (1.31) yield

$$\liminf_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \mathcal{F}_\varepsilon^y(\beta_\varepsilon) \geq \sum_{k=1}^K \varphi^{\mathbb{C}}(R^T \xi^k) + \int_{\Omega \setminus \bigcup_{j=1}^J \bar{B}_\rho(y^j)} \mathbb{C} R^T \beta^{\text{cf}} : R^T \beta^{\text{cf}} \, dx,$$

whence (1.19) follows by sending $\rho \rightarrow 0$.

Therefore, in order to conclude the proof it is enough to show that for any fixed $k = 1, \dots, K$, formula (1.31) holds true. To this end, we fix $k \in \{1, \dots, K\}$ and we set

$$\tilde{\mu}_\varepsilon^k := \varepsilon \sum_{x_\varepsilon^n \in B_\rho(x^k)} \xi_\varepsilon^n \delta_{x_\varepsilon^n};$$

trivially, $\frac{\tilde{\mu}_\varepsilon^k}{\varepsilon} \xrightarrow{*} \xi^k \delta_{x^k}$ and

$$\frac{\tilde{\mu}_\varepsilon^k(B_\rho(x^k))}{\varepsilon} \rightarrow \xi^k \quad \text{as } \varepsilon \rightarrow 0. \quad (1.32)$$

Up to a subsequence, we may assume that the cardinality of $\text{supp } \tilde{\mu}_\varepsilon^k$ is given by some $N^k \in \mathbb{N}$ independent of ε . Up to a relabeling, we can also assume that (recall that k is fixed) $\text{supp } \tilde{\mu}_\varepsilon^k = \{x_\varepsilon^1, \dots, x_\varepsilon^{N^k}\}$. Let (for ε small enough) $g_\varepsilon^k : (\frac{|\log \rho|}{|\log \varepsilon|}, 1) \rightarrow \mathbb{N}$ denote the function that associates to any $q \in (\frac{|\log \rho|}{|\log \varepsilon|}, 1)$ the number $g_\varepsilon^k(q)$ of connected components of $\bigcup_{n=1}^{N^k} B_{\varepsilon^q}(x_\varepsilon^n)$. We have that $g_\varepsilon^k \equiv N^k$ in the interval $[\gamma, 1)$ and g_ε^k is monotonically non-decreasing in $(\frac{|\log \rho|}{|\log \varepsilon|}, 1)$; hence, it can have at most $L_\varepsilon^k \leq N^k$ discontinuities. Up to passing to a subsequence, we can assume that $L_\varepsilon^k = L^k$ for some L^k independent of ε . Let

$$\{q_\varepsilon^{k,l}\}_{l=1, \dots, L^k} \subset \left(\frac{|\log \rho|}{|\log \varepsilon|}, \gamma \right)$$

denote the set of discontinuity points of the function g_ε^k with $q_\varepsilon^{k,l} < q_\varepsilon^{k,l+1}$ for every $l = 1, \dots, L^k - 1$. There exists a finite set $\Delta := \{q^{k,1}, \dots, q^{k,\tilde{L}^k}\} \subset (0, \gamma]$ with $q^{k,m} < q^{k,m+1}$ and $\tilde{L}^k \leq L^k$ such that, up to a subsequence, $\{q_\varepsilon^{k,l}\}_\varepsilon$ converges to some point in Δ , as $\varepsilon \rightarrow 0$ for every $l = 1, \dots, L^k$. Moreover, we set $q^{k,0} := 0$ and $q^{k,\tilde{L}^k+1} := 1$. Let $\lambda > 0$ be such that $2\lambda < \min\{q^{k,m+1} - q^{k,m} : m \in \{0, 1, \dots, \tilde{L}^k\}\}$; then, for ε small enough, the function g_ε^k is constant in the interval $[q^{k,m} + \lambda, q^{k,m+1} - \lambda]$, its value being denoted by $I_\varepsilon^{k,m}$.

One can easily see (in fact, the following arguing is a part of the so called ball construction as done for instance in [24], to which we refer the reader for further details) that for every $m = 0, 1, \dots, \tilde{L}^k$ there exists a family of $I_\varepsilon^{k,m}$ annuli

$$C_\varepsilon^{k,m,i} := B_{\varepsilon^{q^{k,m} + \lambda}}(z_\varepsilon^{k,m,i}) \setminus \bar{B}_{\varepsilon^{q^{k,m+1} - \lambda}}(z_\varepsilon^{k,m,i}) \quad (1.33)$$

with $z_\varepsilon^{k,m,i} \in B_\rho(x^k)$ and $i = 1, \dots, I_\varepsilon^{k,m}$ such that the following properties hold true: The annuli $C_\varepsilon^{k,m,i}$ are pairwise disjoint, contained in $B_{2\rho}(x^k)$ and for all $m = 0, 1, \dots, \tilde{L}^k$,

$$\bigcup_{n=1}^{N^k} B_\varepsilon(x_\varepsilon^n) \subset \bigcup_{i=1}^{I_\varepsilon^{k,m}} B_{\varepsilon^{q^{k,m+1} - \lambda}}(z_\varepsilon^{k,m,i}).$$

Setting $\xi_\varepsilon^{k,m,i} := \mu_\varepsilon(B_{\varepsilon^{q^{k,m+1} - \lambda}}(z_\varepsilon^{k,m,i}))$, in view of (1.32), we have that (for every $m = 0, 1, \dots, \tilde{L}^k$)

$$\sum_{i=1}^{I_\varepsilon^{k,m}} \frac{\xi_\varepsilon^{k,m,i}}{\varepsilon} \rightarrow \xi^k \quad \text{as } \varepsilon \rightarrow 0. \quad (1.34)$$

Up to subsequences, we have that (for every $m = 0, 1, \dots, \tilde{L}^k$) $I_\varepsilon^{k,m} = I^{k,m}$ is independent of ε ; we claim that

$$\frac{\xi_\varepsilon^{k,m,i}}{\varepsilon} \rightarrow \hat{\xi}^{k,m,i} \quad \text{for every } i = 1, \dots, I^{k,m}, \quad (1.35)$$

for some vectors

$$\hat{\xi}^{k,m,i} \in \mathbb{R}\mathbb{S} \quad (1.36)$$

with

$$\sum_{i=1}^{I^{k,m}} \hat{\xi}^{k,m,i} = \xi^k. \quad (1.37)$$

Indeed, up to passing to a further subsequence, by construction, $\xi_\varepsilon^{k,m,i} = \varepsilon \sum_{n=1}^{N^{k,m,i}} R_\varepsilon^n b_\varepsilon^n$, for some $N^{k,m,i} \in \mathbb{N}$, $R_\varepsilon^n \in \text{SO}(2)$, and $b_\varepsilon^n \in \mathbb{S}$. By (1.10), we thus have that, for every $n = 1, \dots, N^{k,m,i}$ there exists a rotation $\tilde{R}_\varepsilon^n \in \mathcal{J}(\mathbb{S})$ such that

$$|R_\varepsilon^n b_\varepsilon^n - R_\varepsilon(\tilde{R}_\varepsilon^n)^T b_\varepsilon^n|^2 \leq C_\gamma \varepsilon^{2(1-\gamma)} \sqrt{|\log \varepsilon|}.$$

whence, using that $(\tilde{R}_\varepsilon^n)^T b_\varepsilon^n \in \mathbb{S}$ and that $|R_\varepsilon - R| \rightarrow 0$, we deduce (1.36); finally, (1.37) follows immediately from (1.34) and (1.35).

Fix $r > 1$. For every $m = 0, 1, \dots, \tilde{L}^k$ we set

$$H_\varepsilon^{k,m} := \left[(q^{k,m+1} - q^{k,m} - 2\lambda) \frac{|\log \varepsilon|}{\log r} \right]$$

and, for every $h = 1, \dots, H_\varepsilon^{k,m}$ and for every $i = 1, \dots, I^{k,m}$, we define

$$A_\varepsilon^{k,m,i,h} := A_{r^{h-1} \varepsilon^{q^{k,m+1}-\lambda}, r^h \varepsilon^{q^{k,m+1}-\lambda}}(Z_\varepsilon^{k,m,i}).$$

Recalling the definition of $C_\varepsilon^{k,m,i}$ in (1.33) and setting

$$\hat{\beta}_\varepsilon^{k,m,i,h}(y) := \beta_\varepsilon(r^{h-1} \varepsilon^{q^{k,m+1}-\lambda} y + z_\varepsilon^{k,m,i}) \quad \text{for every } y \in A_{1,r}(0),$$

we get

$$\frac{1}{\varepsilon^2} \int_{C_\varepsilon^{k,m,i}} W(\beta_\varepsilon) dx \geq \sum_{h=1}^{H_\varepsilon^{k,m}} \frac{1}{\varepsilon^2} \int_{A_\varepsilon^{k,m,i,h}} W(\beta_\varepsilon) dx = \sum_{h=1}^{H_\varepsilon^{k,m}} \frac{1}{(r^{1-h} \varepsilon^{1-q^{k,m+1}+\lambda})^2} \int_{A_{1,r}(0)} W(\hat{\beta}_\varepsilon^{k,m,i,h}) dy, \quad (1.38)$$

where we have used the change of variable $y = r^{1-h} \varepsilon^{-q^{k,m+1}+\lambda} (x - z_\varepsilon^{k,m,i})$.

By construction, $\hat{\beta}_\varepsilon^{k,m,i,h} \in L^2(A_{1,r}(0); \mathbb{R}^{2 \times 2})$, $\text{Curl } \hat{\beta}_\varepsilon^{k,m,i,h} = 0$ and

$$\zeta_\varepsilon^{k,m,i,h} := \int_{\partial B_1(0)} \hat{\beta}_\varepsilon^{k,m,i,h} t d\mathcal{H}^1 = r^{1-h} \varepsilon^{-q^{k,m+1}+\lambda} \int_{\partial B_{r^{h-1} \varepsilon^{q^{k,m+1}-\lambda}}(z_\varepsilon^{k,m,i})} \beta_\varepsilon t d\mathcal{H}^1 = r^{1-h} \varepsilon^{-q^{k,m+1}+\lambda} \zeta_\varepsilon^{k,m,i}.$$

By (1.35), we get that

$$\frac{\zeta_\varepsilon^{k,m,i,h}}{r^{1-h} \varepsilon^{1-q^{k,m+1}+\lambda}} \rightarrow \hat{\zeta}^{k,m,i} \quad \text{as } \varepsilon \rightarrow 0.$$

Moreover, by (1.9), we have that

$$\begin{aligned} \int_{A_{1,r}(0)} |\hat{\beta}_\varepsilon^{k,m,i,h} - R_\varepsilon|^2 dy &= (r^{1-h} \varepsilon^{-q^{k,m+1}+\lambda})^2 \int_{A_\varepsilon^{k,m,i,h}} |\beta_\varepsilon - R_\varepsilon|^2 dx \\ &\leq C_\gamma (r^{1-h} \varepsilon^{1-q^{k,m+1}+\lambda})^2 |\log \varepsilon| \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0, \end{aligned}$$

which implies that $\hat{\beta}_\varepsilon^{k,m,i,h} \rightarrow R$ in $L^2(A_{1,r}(0); \mathbb{R}^{2 \times 2})$. Therefore, we can apply Lemma 1.8 with $\delta = r^{1-h} \varepsilon^{1-q^{k,m+1}+\lambda}$, thus obtaining that

$$\frac{1}{(r^{1-h} \varepsilon^{1-q^{k,m+1}+\lambda})^2} \int_{A_{1,r}(0)} W(\hat{\beta}_\varepsilon^{k,m,i,h}) dy \geq \log r \psi_{1,r}^C(R^T \hat{\zeta}^{k,m,i}) - \sigma_\varepsilon \quad (1.39)$$

for some family $\{\sigma_\varepsilon\}_\varepsilon$ (independent of k, m, i and h) with $\sigma_\varepsilon \rightarrow 0$ (as $\varepsilon \rightarrow 0$). Summing (1.39) over $h = 1, \dots, H_\varepsilon^{k,m}$, by (1.38), we have that

$$\begin{aligned} \frac{1}{\varepsilon^2} \int_{C_\varepsilon^{k,m,i}} W(\beta_\varepsilon) dx &\geq H_\varepsilon^{k,m} (\log r \psi_{1,r}^C(R^T \hat{\zeta}^{k,m,i}) - \sigma_\varepsilon) \\ &\geq (q^{k,m+1} - q^{k,m} - 2\lambda) |\log \varepsilon| \left(\psi_{1,r}^C(R^T \hat{\zeta}^{k,m,i}) - \frac{\sigma_\varepsilon}{\log r} \right) - \log r \psi_{1,r}^C(R^T \hat{\zeta}^{k,m,i}) + \sigma_\varepsilon. \end{aligned} \quad (1.40)$$

Notice that by (1.15) and by homogeneity (see (1.17)), $|\psi_{1,r}^C(\zeta) - \psi^C(\zeta)| \leq \omega_r |\zeta|^2$ for some modulus of continuity ω_r with $\omega_r \rightarrow 0$ as $r \rightarrow +\infty$. Now, summing (1.40) first over $i = 1, \dots, I^{k,m}$ and then over $m = 0, 1, \dots, \tilde{L}^k$, and

using (1.36) and (1.37) together with the very definition of $\varphi^{\mathbb{C}}$ in (1.18), we obtain

$$\begin{aligned} \liminf_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \mathcal{E}_\varepsilon^k &\geq \liminf_{\varepsilon \rightarrow 0} \sum_{m=0}^{\tilde{L}^k} (q^{k,m+1} - q^{k,m} - 2\lambda) \sum_{i=1}^{J^{k,m}} \psi_{1,r}^{\mathbb{C}}(R^T \tilde{\xi}^{k,m,i}) \\ &\quad + \liminf_{\varepsilon \rightarrow 0} \sum_{m=0}^{\tilde{L}^k} (q^{k,m+1} - q^{k,m} - 2\lambda) I^{k,m} \left(-\frac{\sigma_\varepsilon}{\log r} \right) \\ &\quad + \liminf_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \sum_{m=0}^{\tilde{L}^k} \sum_{i=1}^{J^{k,m}} (-\log r \psi_{1,r}^{\mathbb{C}}(R^T \tilde{\xi}^{k,m,i}) + \sigma_\varepsilon) \\ &\geq \sum_{m=0}^{\tilde{L}^k} (q^{k,m+1} - q^{k,m} - 2\lambda) \sum_{i=1}^{J^{k,m}} (\psi^{\mathbb{C}}(R^T \tilde{\xi}^{k,m,i}) - \omega_r |\tilde{\xi}^{k,m,i}|^2) \\ &\geq (1 - 2\lambda(\tilde{L}^k + 1)) \varphi^{\mathbb{C}}(R^T \xi^k) - (1 - 2\lambda(\tilde{L}^k + 1)) C \omega_r, \end{aligned}$$

whence (1.31) follows sending $r \rightarrow +\infty$ and $\lambda \rightarrow 0$, thus concluding the proof of (i). \square

We conclude the proof of Theorem 1.7 by constructing the recovery sequence.

Proof of Theorem 1.7 (ii). By a relaxation argument (see [14]), it suffices to prove the claim of the theorem for $\psi^{\mathbb{C}}(R^T \xi^k) = \varphi^{\mathbb{C}}(R^T \xi^k)$ (for every $k = 1, \dots, K$) and $\beta^{\text{cf}} \in L^\infty(\Omega; \mathbb{R}^{2 \times 2})$. Since Ω is simply connected and since $\text{Curl } \beta^{\text{cf}} = 0$, there exists a map $u^{\beta^{\text{cf}}} \in W^{1,\infty}(\Omega; \mathbb{R}^2)$ such that $\beta^{\text{cf}} = \nabla u^{\beta^{\text{cf}}}$.

Setting $C := \{(x_1; 0) : x_1 \geq 0\}$, for every $\zeta \in \mathbb{R}^2$ let $u^{R,\zeta} \in C^\infty(\mathbb{R}^2 \setminus C; \mathbb{R}^2)$ be such that

$$\nabla u^{R,\zeta} = R \beta_{\mathbb{R}^2}^{\zeta, \mathbb{C}}, \quad (1.41)$$

with $\beta_{\mathbb{R}^2}^{\zeta, \mathbb{C}}$ defined in (1.11). For every $k = 1, \dots, K$, setting $C^k := x^k + C$, we define the map $u_\varepsilon^{R,k} \in C^\infty(\mathbb{R}^2 \setminus C^k; \mathbb{R}^2)$ as

$$u_\varepsilon^{R,k}(x) := u^{R,R^T \xi^k}(x - x^k) + \sqrt{|\log \varepsilon|} u^{\beta^{\text{cf}}}(x^k); \quad (1.42)$$

then $\nabla u_\varepsilon^{R,k}(\cdot) = R \beta_{\mathbb{R}^2}^{R^T \xi^k, \mathbb{C}}(\cdot - x^k)$ and

$$[u_\varepsilon^{R,k}] = \xi^k \quad \text{on } C^k \quad (\text{locally in the sense of traces}). \quad (1.43)$$

We define the map $u_\varepsilon^{\text{far}} \in H^1(\Omega_\varepsilon(\mu) \setminus \bigcup_{k=1}^K C^k; \mathbb{R}^2)$ as

$$u_\varepsilon^{\text{far}}(x) := \sum_{k=1}^K u^{R,R^T \xi^k}(x - x^k) + \sqrt{|\log \varepsilon|} u^{\beta^{\text{cf}}}(x). \quad (1.44)$$

Let $\phi \in C^1([1, 2]; [0, 1])$ be such $\phi(1) = 1$ and $\phi(2) = 0$; we define the function $u_\varepsilon \in H^1(\Omega_\varepsilon(\mu) \setminus \bigcup_{k=1}^K C^k; \mathbb{R}^2)$ as

$$u_\varepsilon(x) := \begin{cases} u_\varepsilon^{R,k}(x) & \text{if } x \in A_{\varepsilon, \varepsilon^\nu}(x^k) \text{ for some } k, \\ \phi\left(\frac{|x - x^k|}{\varepsilon^\nu}\right) u_\varepsilon^{R,k}(x) + \left(1 - \phi\left(\frac{|x - x^k|}{\varepsilon^\nu}\right)\right) u_\varepsilon^{\text{far}}(x) & \text{if } x \in A_{\varepsilon^\nu, 2\varepsilon^\nu}(x^k) \text{ for some } k, \\ u_\varepsilon^{\text{far}}(x) & \text{if } x \in \Omega_{2\varepsilon^\nu}(\mu). \end{cases} \quad (1.45)$$

Finally, we set $\beta_\varepsilon := R + \varepsilon \nabla u_\varepsilon$ in $\Omega_\varepsilon(\mu)$ and $\beta_\varepsilon \equiv 0$ elsewhere in Ω , so that $\beta_\varepsilon \in L^2(\Omega; \mathbb{R}^{2 \times 2})$. Notice that, although u_ε jumps across C^k , we have that $\text{Curl } \beta_\varepsilon = 0$ in $\Omega_\varepsilon(\mu)$, since, by (1.43), the tangential derivatives of the traces agree along C^k . Setting $\mu_\varepsilon \equiv \varepsilon \mu$ and $R_\varepsilon \equiv R$ for every $\varepsilon > 0$, we claim that $\{(\mu_\varepsilon; \beta_\varepsilon; R_\varepsilon)\}_\varepsilon$ is a recovery sequence. First, in view of Remark 1.6, $\beta_\varepsilon \in \mathcal{AS}_\varepsilon^\nu(\mu_\varepsilon)$. Second, by the very definition of R_ε and μ_ε , properties (0) and (1) are trivially satisfied. We now show that this is the case also for condition (2). To this end, we first notice that, by (1.11),

$$\|\nabla u^{R,R^T \xi^k}\|_{L^2(\Omega \setminus \bar{B}_\varepsilon(0); \mathbb{R}^{2 \times 2})} \leq C \sqrt{|\log \varepsilon|}, \quad \frac{1}{\sqrt{|\log \varepsilon|}} \|\nabla u^{R,R^T \xi^k}\|_{L^1(\Omega \setminus \bar{B}_\varepsilon(0); \mathbb{R}^{2 \times 2})} \rightarrow 0 \quad \text{for every } k = 1, \dots, K, \quad (1.46)$$

which implies, in particular,

$$\frac{1}{\varepsilon \sqrt{|\log \varepsilon|}} (\beta_\varepsilon - R_\varepsilon) \chi_{A_{\varepsilon, \varepsilon^\nu}(x^k)} = \frac{\nabla u^{R,R^T \xi^k}(\cdot - x^k)}{\sqrt{|\log \varepsilon|}} \chi_{A_{\varepsilon, \varepsilon^\nu}(x^k)} \rightarrow 0 \quad \text{in } L^2(\Omega; \mathbb{R}^{2 \times 2}). \quad (1.47)$$

Moreover, recalling (1.42) and (1.44), for every $k = 1, \dots, K$ and for every $x \in A_{\varepsilon^\nu, 2\varepsilon^\nu}(x^k)$, we have that

$$u_\varepsilon(x) = u^{R, R^T \xi^k}(x - x^k) + \left(1 - \phi\left(\frac{|x - x^k|}{\varepsilon^\nu}\right)\right) \left(\sum_{j \neq k} u^{R, R^T \xi^j}(x - x^j) + \sqrt{|\log \varepsilon|} u^{\beta^{\text{cf}}}(x)\right) + \phi\left(\frac{|x - x^k|}{\varepsilon^\nu}\right) \sqrt{|\log \varepsilon|} u^{\beta^{\text{cf}}}(x^k);$$

hence, for ε small enough and for $x \in A_{\varepsilon^\nu, 2\varepsilon^\nu}(x^k)$, we have

$$\begin{aligned} |\nabla u_\varepsilon(x)| &\leq |\nabla u^{R, R^T \xi^k}(x - x^k)| + \left|1 - \phi\left(\frac{|x - x^k|}{\varepsilon^\nu}\right)\right| \left(\sum_{j \neq k} |\nabla u^{R, R^T \xi^j}(x - x^j)| + \sqrt{|\log \varepsilon|} |\beta^{\text{cf}}(x)|\right) \\ &\quad + \left|\nabla\left(\phi\left(\frac{|x - x^k|}{\varepsilon^\nu}\right)\right)\right| \left(\sum_{j \neq k} |u^{R, R^T \xi^j}(x - x^j)| + \sqrt{|\log \varepsilon|} |u^{\beta^{\text{cf}}}(x) - u^{\beta^{\text{cf}}}(x^k)|\right) \\ &\leq C\varepsilon^{-\nu} + C(1 + \sqrt{|\log \varepsilon|}) + C\varepsilon^{-\nu}(1 + \varepsilon^\nu \sqrt{|\log \varepsilon|}) \leq C\varepsilon^{-\nu}, \end{aligned} \quad (1.48)$$

where we have used that $|\nabla u^{R, R^T \xi^j}(x)| \leq \frac{C}{|x|}$, that $\beta^{\text{cf}} \in L^\infty(\Omega; \mathbb{R}^{2 \times 2})$, and that

$$|u^{\beta^{\text{cf}}}(x) - u^{\beta^{\text{cf}}}(x^k)| \leq \|\beta^{\text{cf}}\|_{L^\infty(\Omega; \mathbb{R}^{2 \times 2})} |x - x^k|.$$

By (1.48) we have immediately that

$$\frac{1}{\varepsilon \sqrt{|\log \varepsilon|}} \|\beta_\varepsilon - R_\varepsilon\|_{L^2(A_{\varepsilon^\nu, 2\varepsilon^\nu}(x^k); \mathbb{R}^{2 \times 2})} = \frac{1}{\sqrt{|\log \varepsilon|}} \|\nabla u_\varepsilon\|_{L^2(A_{\varepsilon^\nu, 2\varepsilon^\nu}(x^k); \mathbb{R}^{2 \times 2})} \leq \frac{C}{\sqrt{|\log \varepsilon|}} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \quad (1.49)$$

Therefore, by (1.47) and (1.49), using again (1.46), we have that

$$\frac{\beta_\varepsilon - R_\varepsilon}{\varepsilon \sqrt{|\log \varepsilon|}} = -\frac{R_\varepsilon}{\varepsilon \sqrt{|\log \varepsilon|}} \chi_{\Omega \setminus \Omega_\varepsilon(\mu_\varepsilon)} + \sum_{k=1}^K \frac{1}{\varepsilon \sqrt{|\log \varepsilon|}} (\beta_\varepsilon - R_\varepsilon) \chi_{A_{\varepsilon, 2\varepsilon^\nu}(x^k)} + \frac{1}{\sqrt{|\log \varepsilon|}} \nabla u_\varepsilon^{\text{far}} \chi_{\Omega_{2\varepsilon^\nu}(\mu_\varepsilon)}, \quad (1.50)$$

i.e., (2).

Now we conclude by proving that the sequence $\{(\mu_\varepsilon; \beta_\varepsilon)\}_\varepsilon$ satisfies estimate (1.20). Since $W \in C^0(\mathbb{R}^{2 \times 2})$ and $\|\beta_{\mathbb{R}^2}^{R^T \xi^k, \mathbb{C}}\|_{L^\infty(\mathbb{R}^2 \setminus \bar{B}_\varepsilon(0); \mathbb{R}^{2 \times 2})} \leq \frac{C}{\varepsilon}$ for some $C > 0$ (independent of ε), for every $\lambda > 1$ we have

$$\frac{1}{\varepsilon^2 |\log \varepsilon|} \int_{A_{\varepsilon, \lambda \varepsilon}(x^k)} W(\beta_\varepsilon) \, dx \leq \frac{C\lambda^2}{|\log \varepsilon|} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \quad (1.51)$$

Moreover, note that, by assumption (v), there exist $\rho > 0$ and $c_\rho > 0$ such that

$$W(M) \leq c_\rho \text{dist}^2(M, \text{SO}(2)) \quad \text{for every } M \in \text{SO}(2) + B_\rho(0). \quad (1.52)$$

Let $\rho > 0$ be such that (1.52) holds true, let $\lambda > 1$ be such that $\|\beta_{\mathbb{R}^2}^{R^T \xi^k, \mathbb{C}}\|_{L^\infty(\mathbb{R}^2 \setminus \bar{B}_{\lambda \varepsilon}(0); \mathbb{R}^{2 \times 2})} \leq \frac{\rho}{\varepsilon}$ and let $\alpha \in (\gamma, 1)$. By the very definition of β_ε , using (ii) and (1.52), one can check that for every $k = 1, \dots, K$,

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \int_{A_{\lambda \varepsilon, \varepsilon^\alpha}(x^k)} W(\beta_\varepsilon) \, dx &= \limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \int_{A_{\lambda \varepsilon, \varepsilon^\alpha}(x^k)} W(R_\varepsilon(\text{Id} + \varepsilon \beta_{\mathbb{R}^2}^{R^T \xi^k, \mathbb{C}}(x - x^k))) \, dx \\ &\leq \limsup_{\varepsilon \rightarrow 0} \frac{c_\rho}{|\log \varepsilon|} \int_{A_{\lambda \varepsilon, \varepsilon^\alpha}(0)} |\beta_{\mathbb{R}^2}^{R^T \xi^k, \mathbb{C}}|^2 \, dx \leq C(1 - \alpha). \end{aligned} \quad (1.53)$$

Moreover, since $\|\varepsilon \beta_{\mathbb{R}^2}^{R^T \xi^k, \mathbb{C}}\|_{L^\infty} \leq C\varepsilon^{1-\alpha}$ in $A_{\varepsilon^\alpha, \varepsilon^\nu}(0)$, by considering the Taylor expansion of W around the identity, for any $k = 1, \dots, K$ we have

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \int_{A_{\varepsilon^\alpha, \varepsilon^\nu}(x^k)} W(\beta_\varepsilon) \, dx &= \limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \int_{A_{\varepsilon^\alpha, \varepsilon^\nu}(x^k)} W(R_\varepsilon(\text{Id} + \varepsilon \beta_{\mathbb{R}^2}^{R^T \xi^k, \mathbb{C}}(x - x^k))) \, dx \\ &= \limsup_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \int_{A_{\varepsilon^\alpha, \varepsilon^\nu}(0)} \frac{1}{2} \mathbb{C} \beta_{\mathbb{R}^2}^{R^T \xi^k, \mathbb{C}} : \beta_{\mathbb{R}^2}^{R^T \xi^k, \mathbb{C}} + \frac{1}{\varepsilon^2} \sigma(\varepsilon \beta_{\mathbb{R}^2}^{R^T \xi^k, \mathbb{C}}) \, dx \end{aligned}$$

with $\lim_{|M| \rightarrow 0} \frac{\sigma(M)}{|M|^2} = 0$. Therefore, in view of (1.11) and (1.14), we get

$$\limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \int_{A_{\varepsilon \alpha, \varepsilon^\gamma}(x^k)} W(\beta_\varepsilon) \, dx \leq (\alpha - \gamma) \psi^{\mathbb{C}}(R^T \xi^k). \quad (1.54)$$

Furthermore, similar computations (or arguing verbatim as in the proof of [26, Proposition 3.12] for all details) show that

$$\limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \int_{\Omega_{2\varepsilon^\gamma}(\mu)} W(\beta_\varepsilon) \, dx \leq \gamma \sum_{k=1}^K \psi^{\mathbb{C}}(R^T \xi^k) + \frac{1}{2} \int_{\Omega} \mathbb{C} R^T \beta^{\text{cf}} : R^T \beta^{\text{cf}} \, dx. \quad (1.55)$$

Finally, by (1.49) and (1.52), for every $k = 1, \dots, K$ we have

$$\frac{1}{\varepsilon^2 |\log \varepsilon|} \int_{A_{\varepsilon^\gamma, 2\varepsilon^\gamma}(x^k)} W(\beta_\varepsilon) \, dx \leq \frac{c}{\varepsilon^2 |\log \varepsilon|} \|\beta_\varepsilon - R_\varepsilon\|_{L^2(A_{\varepsilon^\gamma, 2\varepsilon^\gamma}(x^k); \mathbb{R}^{2 \times 2})}^2 \rightarrow 0,$$

which, combined with (1.51), (1.53), (1.54) and (1.55), implies (1.20) (sending α to 1). \square

Remark 1.9. Let $0 < \gamma \leq \gamma' < 1$. For every $\varepsilon > 0$, we can define the functional

$$\mathcal{F}_\varepsilon^{\gamma, \gamma'} : \mathcal{M}(\mathbb{R}^2; \mathbb{R}^2) \times L^2(\Omega; \mathbb{R}^{2 \times 2}) \rightarrow [0, +\infty]$$

as

$$\mathcal{F}_\varepsilon^{\gamma, \gamma'}(\mu, \beta) := \begin{cases} \int_{\Omega_\varepsilon(\mu)} W(\beta) \, dx & \text{if } \mu \in X_\varepsilon^\gamma(\Omega) \text{ and } \beta \in \mathcal{AS}_\varepsilon^{\gamma'}(\mu), \\ +\infty & \text{otherwise.} \end{cases}$$

Clearly, $\mathcal{F}_\varepsilon^{\gamma, \gamma} \equiv \mathcal{F}_\varepsilon^\gamma$. By following verbatim the proofs of Theorem 1.1 and Theorem 1.7, one can check that the same compactness and Γ -convergence statements hold true also for the functional $\mathcal{F}_\varepsilon^{\gamma, \gamma'}$.

Furthermore, for every $\varepsilon, \delta_\varepsilon > 0$ with $\delta_\varepsilon < \varepsilon^\gamma$ we can define the class $\mathcal{AS}_{\varepsilon, \delta_\varepsilon}(\mu)$ of admissible strains for a measure $\mu \in X_\varepsilon^\gamma(\Omega)$ as in (1.2), with the condition $\int_{A_{\varepsilon, \varepsilon^\gamma}(x^n)} \beta \, dx \in R^{n\mathcal{J}}(\mathbb{S})$ replaced by $\int_{A_{\varepsilon, \delta_\varepsilon}(x^n)} \beta \, dx \in R^{n\mathcal{J}}(\mathbb{S})$; analogously, we can define the energy functional

$$\mathcal{F}_{\varepsilon, \delta_\varepsilon}^\gamma(\mu, \beta) := \begin{cases} \int_{\Omega_\varepsilon(\mu)} W(\beta) \, dx & \text{if } \mu \in X_\varepsilon^\gamma(\Omega) \text{ and } \beta \in \mathcal{AS}_{\varepsilon, \delta_\varepsilon}(\mu), \\ +\infty & \text{otherwise.} \end{cases}$$

One can check that if $\delta_\varepsilon \gg \varepsilon \sqrt{|\log \varepsilon|}$, the compactness and Γ -convergence results proved for the functional $\mathcal{F}_\varepsilon^\gamma$ still hold true. As observed in Remark 1.5, if $\delta_\varepsilon \sim \varepsilon$ the coherence of the micro-rotations around each dislocation with the macroscopic rotation provided by linearization would fail.

2 The purely discrete model

In this section we introduce and analyze the nonlinear purely discrete model for the elastic energy induced by a family of edge dislocations.

2.1 Description of the problem

Here we introduce the main notation that will be used in this section.

The reference lattice. We set $\nu := \frac{1}{2}e_1 + \frac{\sqrt{3}}{2}e_2$ and $\eta := -\frac{1}{2}e_1 + \frac{\sqrt{3}}{2}e_2$. Let $\mathbb{T} := \text{span}_{\mathbb{Z}}\{e_1, \nu\}$ and set

$$T^+ := \text{conv}\{0, e_1, \nu\} \quad \text{and} \quad T^- := \text{conv}\{0, e_1, -\eta\},$$

where, for every $a, b, c \in \mathbb{R}^2$, the set $\text{conv}\{a, b, c\}$ denotes the convex envelope of the points a, b, c , i.e., the (closed) triangle with vertices at a, b, c . For every $\varepsilon > 0$ we denote by \mathcal{T}_ε the family of the triangles T_ε of the form $i + \varepsilon T^\pm$, with $i \in \varepsilon\mathbb{T}$. Moreover, we set

$$\mathcal{T}_\varepsilon(\Omega) := \{T_\varepsilon \in \mathcal{T}_\varepsilon : T_\varepsilon \subset \Omega\}$$

and we define $\Omega_{\mathcal{T}_\varepsilon} := \bigcup_{T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)} T_\varepsilon$. Furthermore, we set $\Omega_\varepsilon^0 := \Omega_{\mathcal{T}_\varepsilon} \cap \varepsilon\mathbb{T}$ and we denote by Ω_ε^1 the family of nearest neighbor bonds in $\Omega_{\mathcal{T}_\varepsilon}$, i.e., $\Omega_\varepsilon^1 := \{(i, j) \in \Omega_\varepsilon^0 \times \Omega_\varepsilon^0 : |i - j| = \varepsilon\}$. Trivially, $(i, j) \in \Omega_\varepsilon^1$ if and only if $(j, i) \in \Omega_\varepsilon^1$.

In the following we will generalize the notation introduced above to general subsets of \mathbb{R}^2 (not necessarily open). In particular, for every triangle $T_\varepsilon \in \mathcal{T}_\varepsilon$, we have

$$(T_\varepsilon)_\varepsilon^1 = \{(i, j) \in (T_\varepsilon \cap \varepsilon\mathbb{T}) \times (T_\varepsilon \cap \varepsilon\mathbb{T}) : i \neq j\}.$$

For every $T_\varepsilon \in \mathcal{T}_\varepsilon$ and for every map $V : (T_\varepsilon)_\varepsilon^1 \rightarrow \mathbb{R}^2$, we define the *discrete circulation* of V on the “boundary of T_ε ” as

$$\text{d}V(T_\varepsilon) := V(i, j) + V(j, k) + V(k, i),$$

where (i, j, k) is a triple of counter-clockwise oriented vertices of T_ε .

The admissible strains and the energy functional. For every $\varepsilon > 0$ we define the class of discrete strains as

$$\mathcal{S}_\varepsilon^{\text{discr}}(\Omega) := \{\beta : \Omega_\varepsilon^1 \rightarrow \mathbb{R}^2 : \beta(i, j) = -\beta(j, i) \text{ for any } (i, j) \in \Omega_\varepsilon^1\}.$$

In the following, for every $\beta \in \mathcal{S}_\varepsilon^{\text{discr}}(\Omega)$, we set

$$\mu[\beta] := \sum_{T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)} \text{d}\beta(T_\varepsilon) \delta_{x_{T_\varepsilon}},$$

where the point x_{T_ε} denotes the barycenter of the ε -triangle T_ε . For any $\beta \in \mathcal{S}_\varepsilon^{\text{discr}}(\Omega)$ and for any $T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)$ with $\mu[\beta](T_\varepsilon) = 0$ we denote by $\tilde{\beta}^{T_\varepsilon}$ the matrix in $\mathbb{R}^{2 \times 2}$ uniquely defined by the following property:

$$\beta(i, j) = \tilde{\beta}^{T_\varepsilon}(j - i) \quad \text{for every } (i, j) \in (T_\varepsilon)_\varepsilon^1. \quad (2.1)$$

Moreover, we define the field $\tilde{\beta}^{\mathcal{T}_\varepsilon} \in L^2(\Omega; \mathbb{R}^{2 \times 2})$ as

$$\tilde{\beta}^{\mathcal{T}_\varepsilon} := \sum_{\substack{T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega) \\ \mu[\beta](T_\varepsilon) = 0}} \tilde{\beta}^{T_\varepsilon} \chi_{T_\varepsilon \cap \Omega_\varepsilon}(\mu[\beta]); \quad (2.2)$$

notice that every $T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)$ with $\mu[\beta](T_\varepsilon) \neq 0$ is contained in $B_\varepsilon(x_{T_\varepsilon})$.

Let $\psi_1, \psi_2 \in C(\mathbb{R}; [0, +\infty))$ be such that $\psi_1^{-1}(0) = \psi_2^{-1}(0) = \{1\}$, $\psi_1, \psi_2 \in C^2$ in a neighborhood of 1, and $\psi_1''(1), \psi_2''(1) > 0$. Assume moreover that there exist $a, b > 0$ such that $\psi_1(t) \geq at^2 - b$, for every $t \in [0, +\infty)$.

We define the discrete energy functional $E_\varepsilon : \mathcal{S}_\varepsilon^{\text{discr}}(\Omega) \rightarrow [0, +\infty)$ as

$$E_\varepsilon(\beta) := \sum_{(i, j) \in \Omega_\varepsilon^1} \varepsilon^2 \psi_1\left(\frac{|\beta(i, j)|}{\varepsilon}\right) + \sum_{\substack{(i, j), (i, k) \in \Omega_\varepsilon^1 \\ \langle (j-i) \wedge (k-i), e_3 \rangle > 0}} \varepsilon^2 \psi_2\left(\frac{2}{\sqrt{3}\varepsilon^2} \langle \beta(i, j) \wedge \beta(i, k), e_3 \rangle\right),$$

where the symbol \wedge denotes the wedge product between vectors in \mathbb{R}^2 , namely,

$$v \wedge w := (v_1 w_2 - v_2 w_1) e_3 \quad \text{for each } v = (v_1; v_2), w = (w_1; w_2) \in \mathbb{R}^2.$$

We remark that, while the two-body interaction potential ψ_1 accounts for the elongation of the bonds, the three-body interaction term ψ_2 acts as a penalization of changes of area of the triangular cells.

We will consider also localized versions of our energy functional E_ε . More specifically, for every open and bounded set $A \subset \mathbb{R}^2$, we define $E_\varepsilon(\cdot; A) : \mathcal{S}_\varepsilon^{\text{discr}}(A) \rightarrow [0, +\infty)$ as

$$E_\varepsilon(\beta; A) := \sum_{(i, j) \in A_\varepsilon^1} \varepsilon^2 \psi_1\left(\frac{|\beta(i, j)|}{\varepsilon}\right) + \sum_{\substack{(i, j), (i, k) \in A_\varepsilon^1 \\ \langle (j-i) \wedge (k-i), e_3 \rangle > 0}} \varepsilon^2 \psi_2\left(\frac{2}{\sqrt{3}\varepsilon^2} \langle \beta(i, j) \wedge \beta(i, k), e_3 \rangle\right),$$

so that $E_\varepsilon(\beta; \Omega) = E_\varepsilon(\beta)$.

Let $0 < \gamma < 1$ be fixed. We set $S := \{e_1, \nu\}$, so that, following the notation in Section 1, $\mathbb{S} = \mathbb{T}$. We define

$$X_\varepsilon^{\gamma, \text{discr}}(\Omega) := \left\{ \mu \in X_\varepsilon^\gamma(\Omega) : \text{supp } \mu \subset \bigcup_{T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)} \{x_{T_\varepsilon}\} \right\},$$

where the class $X_\varepsilon^\gamma(\Omega)$ is defined in (1.1).

Moreover, for any $\mu = \varepsilon \sum_{n=1}^N R^n b^n \delta_{x^n} \in X_\varepsilon^{\gamma, \text{discr}}(\Omega)$, with $N \in \mathbb{N}$, $R^n \in \text{SO}(2)$, $b^n \in \mathbb{T}$, and $x^n \in \Omega$, we define the class of admissible discrete strains associated to μ as

$$\mathcal{AS}_\varepsilon^{\gamma, \text{discr}}(\mu) := \left\{ \beta \in \mathcal{S}_\varepsilon^{\text{discr}}(\Omega) : \int_{A_{\varepsilon, \nu}(x^n)} \tilde{\beta}^{\mathcal{J}} dx \in R^{\mathcal{J}}(\mathbb{T}), \mu[\beta] = \mu \right\},$$

where $\tilde{\beta}^{\mathcal{J}}$ is the map defined in (2.2) and $\mathcal{J}(\mathbb{T})$ is the group of rotations generated by the $\frac{\pi}{3}$ clockwise rotation $R(\frac{\pi}{3})$.

For every $0 < \varepsilon < 1$ we define the discrete energy $\mathcal{E}_\varepsilon^\gamma : \mathcal{M}(\mathbb{R}^2; \mathbb{R}^2) \times \mathcal{S}_\varepsilon^{\text{discr}}(\Omega) \rightarrow [0, +\infty]$ as

$$\mathcal{E}_\varepsilon^\gamma(\mu, \beta) := \begin{cases} E_\varepsilon(\beta) & \text{if } \mu \in X_\varepsilon^{\gamma, \text{discr}}(\Omega) \text{ and } \beta \in \mathcal{AS}_\varepsilon^{\gamma, \text{discr}}(\mu), \\ +\infty & \text{otherwise.} \end{cases}$$

Before stating and proving our Γ -convergence result for the functional $\mathcal{E}_\varepsilon^\gamma$, we first derive the continuous nonlinear elastic energy density associated to the discrete functional E_ε as well as the corresponding linearized elasticity tensor.

Remark 2.1. Let $\beta \in \mathcal{S}_\varepsilon^{\text{discr}}(\Omega)$; for every $T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)$ we define

$$\tilde{E}_\varepsilon(\beta; T_\varepsilon) := \frac{1}{2} \sum_{(i,j) \in (T_\varepsilon)_\varepsilon^1} \varepsilon^2 \psi_1 \left(\frac{|\beta(i,j)|}{\varepsilon} \right) + \sum_{\substack{(i,j), (i,k) \in (T_\varepsilon)_\varepsilon^1 \\ \langle (j-i) \wedge (k-i), e_3 \rangle > 0}} \varepsilon^2 \psi_2 \left(\frac{2}{\sqrt{3} \varepsilon^2} \langle \beta(i,j) \wedge \beta(i,k), e_3 \rangle \right).$$

Note that, for every $A \subset \Omega$,

$$E_\varepsilon(\beta; A) = \sum_{\substack{T_\varepsilon \in \mathcal{T}_\varepsilon \\ T_\varepsilon \subset A}} \tilde{E}_\varepsilon(\beta; T_\varepsilon) + \frac{1}{2} \sum_{\substack{(i,j) \in A_\varepsilon^1 \\ i,j \in \partial A_{T_\varepsilon}}} \varepsilon^2 \psi_1 \left(\frac{|\beta(i,j)|}{\varepsilon} \right). \quad (2.3)$$

Moreover, for any $T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)$ with $\mu[\beta](T_\varepsilon) = 0$, we easily get that

$$\begin{aligned} \tilde{E}_\varepsilon(\beta; T_\varepsilon) &= \varepsilon^2 \left[\frac{1}{2} \left(\psi_1(|\tilde{\beta}^{T_\varepsilon} e_1|) + \psi_1(|\tilde{\beta}^{T_\varepsilon} \nu|) + \psi_1(|\tilde{\beta}^{T_\varepsilon} \eta|) \right) + 3 \psi_2(\det \tilde{\beta}^{T_\varepsilon}) \right] \\ &= \frac{\sqrt{3}}{4} \varepsilon^2 W(\tilde{\beta}^{T_\varepsilon}) \\ &= \int_{T_\varepsilon} W(\tilde{\beta}^{T_\varepsilon}) dx, \end{aligned} \quad (2.4)$$

where $\tilde{\beta}^{T_\varepsilon}$ is the matrix defined in (2.1) and W is given by

$$W(M) := \frac{4}{\sqrt{3}} \left[\frac{1}{2} \left(\psi_1(|M e_1|) + \psi_1(|M \nu|) + \psi_1(|M \eta|) \right) + 3 \psi_2(\det M) \right]. \quad (2.5)$$

The assumptions on ψ_1, ψ_2 yield that the map $W : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}$ satisfies assumptions (i)–(v). Setting $\alpha_1 := \psi_1''(1)$ and $\alpha_2 := \psi_2''(1)$, the corresponding tensor \mathbb{C} in (1.3) is given by

$$\mathbb{C} \delta : \delta := \frac{4}{\sqrt{3}} \left[\frac{1}{2} \alpha_1 (|e_1^* \delta e_1|^2 + |\nu^* \delta \nu|^2 + |\eta^* \delta \eta|^2) + 3 \alpha_2 |\text{tr } \delta|^2 \right] \quad \text{for every } \delta \in \mathbb{R}^{2 \times 2}. \quad (2.6)$$

A straightforward computation shows that

$$\mathbb{C} \delta : \delta = \frac{4}{\sqrt{3}} \left[\frac{1}{2} \alpha_1 \left(\frac{3}{8} |\text{tr } \delta|^2 + \frac{3}{4} |\delta^{\text{sym}}|^2 \right) + 3 \alpha_2 |\text{tr } \delta|^2 \right] = \left(\frac{\sqrt{3}}{4} \alpha_1 + 4 \sqrt{3} \alpha_2 \right) |\text{tr } \delta|^2 + \frac{\sqrt{3}}{2} \alpha_1 |\delta^{\text{sym}}|^2.$$

In particular, \mathbb{C} is isotropic with Lamé moduli $\lambda = \frac{\sqrt{3}}{4}\alpha_1 + 4\sqrt{3}\alpha_2$ and $\mu = \frac{\sqrt{3}}{4}\alpha_1$. In such a case, the function $\psi^{\mathbb{C}}$ defined in (1.14) is given (see, for instance, [9]) by

$$\psi^{\mathbb{C}}(b) = \frac{1}{4\pi} \frac{\mu(\lambda + \mu)}{\lambda + 2\mu} |b|^2 =: C(\alpha_1, \alpha_2) |b|^2,$$

and hence, recalling the definition of $\varphi^{\mathbb{C}}$ in (1.18), we have

$$\varphi^{\mathbb{C}}(b) = C(\alpha_1, \alpha_2) \min \left\{ \sum_{i=1}^3 |z_i| : z_1, z_2, z_3 \in \mathbb{Z}, b = z_1 e_1 + z_2 v + z_3 \eta \right\}. \quad (2.7)$$

By (2.4), (2.2) and (2.3), for each $\mu \in X_\varepsilon^{\gamma, \text{discr}}(\Omega)$ and $\beta \in \mathcal{S}_\varepsilon^{\text{discr}}(\Omega)$ with $\mu[\beta] = \mu$, and for every open set $A \subset\subset \Omega$, for ε sufficiently small it holds

$$E_\varepsilon(\beta) \geq \int_{A_\varepsilon(\mu)} W(\tilde{\beta}^{\mathbb{J}_\varepsilon}) \, dx. \quad (2.8)$$

By a reflection argument (see, for instance, [4, Lemma 4.3]), one can prove the following result.

Lemma 2.2. *Let $\varepsilon > 0$, $\mu \in X_\varepsilon^{\gamma, \text{discr}}(\Omega)$ and $\beta \in \mathcal{A}\mathcal{S}_\varepsilon^{\gamma, \text{discr}}(\mu)$ and let $\tilde{\beta}^{\mathbb{J}_\varepsilon}$ be the map defined in (2.2). Then there exists a field $\hat{\beta}^{\mathbb{J}_\varepsilon} \in L^2(\Omega; \mathbb{R}^{2 \times 2})$ such that*

- (i) $\hat{\beta}^{\mathbb{J}_\varepsilon} = \tilde{\beta}^{\mathbb{J}_\varepsilon}$ in $\Omega_{\mathbb{J}_\varepsilon}$,
- (ii) $\text{Curl } \hat{\beta}^{\mathbb{J}_\varepsilon} = 0$ in $\Omega_\varepsilon(\mu)$ (in the sense of distributions),
- (iii) $\int_{\Omega_\varepsilon(\mu)} W(\hat{\beta}^{\mathbb{J}_\varepsilon}) \, dx \leq C E_\varepsilon(\beta)$, for some constant C independent of ε .

2.2 The main result for the discrete energy

We are now in a position to state and prove our Γ -convergence result for the functionals $\mathcal{E}_\varepsilon^\gamma$ as $\varepsilon \rightarrow 0$ in the $|\log \varepsilon|$ regime.

Theorem 2.3. *Let \mathbb{C} be the elasticity tensor defined by (2.6) and let $\varphi^{\mathbb{C}}$ be the function defined in (2.7). The following Γ -convergence result holds true.*

- (i) *Let $\{(\mu_\varepsilon; \beta_\varepsilon)\}_\varepsilon$, with $\mu_\varepsilon \in \mathcal{M}(\mathbb{R}^2; \mathbb{R}^2)$ and $\beta_\varepsilon \in \mathcal{S}_\varepsilon^{\text{discr}}(\Omega)$ (for every $\varepsilon > 0$) satisfy*

$$\sup_{\varepsilon > 0} \mathcal{E}_\varepsilon^\gamma(\mu_\varepsilon, \beta_\varepsilon) \leq C \varepsilon^2 |\log \varepsilon| \quad (2.9)$$

for some constant $C > 0$. Then there exist a sequence of rotations $\{R_\varepsilon\}_\varepsilon \subset \text{SO}(2)$, a rotation $R \in \text{SO}(2)$, a measure $\mu \in \overline{X}^R(\Omega)$, and a field $\beta^{\text{cf}} \in L^2(\Omega; \mathbb{R}^{2 \times 2})$ with $\text{Curl } \beta^{\text{cf}} = 0$, such that, up to a subsequence,

$$(\mu_\varepsilon; \tilde{\beta}_\varepsilon^{\mathbb{J}_\varepsilon}; R_\varepsilon) \rightarrow (\mu; \beta^{\text{cf}}; R) \quad \text{as } \varepsilon \rightarrow 0$$

in the sense of Definition 1.2.

- (ii) (Γ -liminf inequality) *Let $R \in \text{SO}(2)$, $\mu = \sum_{k=1}^K \xi^k \delta_{x^k} \in \overline{X}^R(\Omega)$, $\beta^{\text{cf}} \in L^2(\Omega; \mathbb{R}^{2 \times 2})$ with $\text{Curl } \beta^{\text{cf}} = 0$. For every $\{(\mu_\varepsilon; \beta_\varepsilon; R_\varepsilon)\}_\varepsilon$, with $\mu_\varepsilon \in X_\varepsilon^{\gamma, \text{discr}}(\Omega)$, $\beta_\varepsilon \in \mathcal{S}_\varepsilon^{\text{discr}}(\Omega)$ and $R_\varepsilon \in \text{SO}(2)$ (for every $\varepsilon > 0$), satisfying $(\mu_\varepsilon; \tilde{\beta}_\varepsilon^{\mathbb{J}_\varepsilon}; R_\varepsilon) \rightarrow (\mu; \beta^{\text{cf}}; R)$ as $\varepsilon \rightarrow 0$, it holds*

$$\sum_{k=1}^K \varphi^{\mathbb{C}}(R^T \xi^k) + \int_{\Omega} \frac{1}{2} \mathbb{C} R^T \beta^{\text{cf}} : R^T \beta^{\text{cf}} \, dx \leq \liminf_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \mathcal{E}_\varepsilon^\gamma(\mu_\varepsilon, \beta_\varepsilon). \quad (2.10)$$

- (iii) (Γ -limsup inequality) *Let $R \in \text{SO}(2)$, $\mu = \sum_{k=1}^K \xi^k \delta_{x^k} \in \overline{X}^R(\Omega)$ and $\beta^{\text{cf}} \in L^2(\Omega; \mathbb{R}^{2 \times 2})$ with $\text{Curl } \beta^{\text{cf}} = 0$. Then there exists $\{(\mu_\varepsilon; \beta_\varepsilon; R_\varepsilon)\}_\varepsilon$, with $\mu_\varepsilon \in X_\varepsilon^{\gamma, \text{discr}}(\Omega)$, $\beta_\varepsilon \in \mathcal{S}_\varepsilon^{\text{discr}}(\Omega)$ and $R_\varepsilon \in \text{SO}(2)$ (for every $\varepsilon > 0$), such that $(\mu_\varepsilon; \tilde{\beta}_\varepsilon^{\mathbb{J}_\varepsilon}; R_\varepsilon) \rightarrow (\mu; \beta^{\text{cf}}; R)$ as $\varepsilon \rightarrow 0$ and*

$$\sum_{k=1}^K \varphi^{\mathbb{C}}(R^T \xi^k) + \int_{\Omega} \frac{1}{2} \mathbb{C} R^T \beta^{\text{cf}} : R^T \beta^{\text{cf}} \, dx \geq \limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \mathcal{E}_\varepsilon^\gamma(\mu_\varepsilon, \beta_\varepsilon). \quad (2.11)$$

Proof. We divide the proof into three parts.

Proof of (i). In view of (2.9), the maps $\hat{\beta}_\varepsilon^{\mathcal{J}_\varepsilon}$ provided by Lemma 2.2 belong to $\mathcal{AS}_\varepsilon^\gamma(\mu_\varepsilon)$ and satisfy

$$\mathcal{F}_\varepsilon^\gamma(\mu_\varepsilon, \hat{\beta}_\varepsilon^{\mathcal{J}_\varepsilon}) \leq C\varepsilon^2 |\log \varepsilon|,$$

where the functional $\mathcal{F}_\varepsilon^\gamma$ is defined in (1.4) for the choice of W in (2.5). Therefore, the conclusion follows by Theorem 1.1 and Lemma 2.2 (i), once noticed that $\|R_\varepsilon\|_{L^2(\Omega \setminus \Omega_{\mathcal{J}_\varepsilon})} \leq C\varepsilon$.

Proof of (ii). Without loss of generality we may assume that (2.9) is satisfied. In view of (1) we have that $|\frac{\mu_\varepsilon}{\varepsilon}|(\Omega) \leq C$ for some C independent of ε , so that, up to passing to a subsequence, $|\frac{\mu_\varepsilon}{\varepsilon}| \xrightarrow{*} \bar{\mu}$ for some measure $\bar{\mu} \in \mathcal{M}(\mathbb{R}^2; \mathbb{R}^2)$ with finite support contained in Ω . Let $A \subset\subset \Omega$ be an open set with Lipschitz continuous boundary such that $\text{supp } \bar{\mu} \subset A$. By construction, for sufficiently small $\varepsilon > 0$ we have that $\mu_\varepsilon \llcorner A \in \mathcal{X}_\varepsilon^\gamma(A)$ and $\hat{\beta}_\varepsilon^{\mathcal{J}_\varepsilon} \llcorner A \in \mathcal{AS}_\varepsilon^\gamma(A)$. Then (2.10) follows by (2.8), Theorem 1.7 (i) applied with $\Omega = A$, and by the arbitrariness of A .

Proof of (iii). By standard density arguments in Γ -convergence we can assume that $\varphi^{\mathbb{C}}(R^T \xi^k) = \psi^{\mathbb{C}}(R^T \xi^k)$ for every $k = 1, \dots, K$ and that $\beta^{\text{cf}} \in W^{1,\infty}(\Omega; \mathbb{R}^{2 \times 2})$. Let furthermore $u^{\beta^{\text{cf}}} \in W^{2,\infty}(\Omega; \mathbb{R}^2)$ be such that $\beta^{\text{cf}} = \nabla u^{\beta^{\text{cf}}}$.

For every $k = 1, \dots, K$ let

$$x^{k,\varepsilon} \in \text{argmin}\{|x^k - x_{T_\varepsilon}| : T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)\}$$

and set $\mu_\varepsilon := \varepsilon \sum_{k=1}^K \xi^k \delta_{x^{k,\varepsilon}}$. Then, for ε small enough, $\mu_\varepsilon \in \mathcal{X}_\varepsilon^{\gamma, \text{discr}}(\Omega)$. Moreover, we set $R_\varepsilon \equiv R$. In order to construct β_ε , we proceed as in the proof of Theorem 1.7 (ii). We first construct u_ε as in (1.45), replacing x^k with $x^{k,\varepsilon}$ and suitably modifying the radii of the annuli.

More precisely, for every $k = 1, \dots, K$ we define the map $u_\varepsilon^{R,k} \in C^\infty(\mathbb{R}^2 \setminus C^{k,\varepsilon}; \mathbb{R}^2)$ as

$$u_\varepsilon^{R,k}(x) := u^{R, R^T \xi^k}(x - x^{k,\varepsilon}) + \sqrt{|\log \varepsilon|} u^{\beta^{\text{cf}}}(x^{k,\varepsilon}),$$

with $C^{k,\varepsilon} := x^{k,\varepsilon} + C$ (with $C := \{(x_1; 0) : x_1 \geq 0\}$) and $u^{R,\zeta}$ satisfying (1.41). Thus, $\nabla u_\varepsilon^{R,k}(\cdot) = R \beta_{\mathbb{R}^2}^{R^T \xi^k, \mathbb{C}}(\cdot - x^{k,\varepsilon})$ and $[u_\varepsilon^{R,k}] = \xi^k$ on $C^{k,\varepsilon}$ (locally in the sense of traces).

We define the map $u_\varepsilon^{\text{far}} : \Omega_\varepsilon(\mu_\varepsilon) \rightarrow \mathbb{R}^2$ as

$$u_\varepsilon^{\text{far}}(x) := \sum_{k=1}^K u^{R, R^T \xi^k}(x - x^{k,\varepsilon}) + \sqrt{|\log \varepsilon|} u^{\beta^{\text{cf}}}(x).$$

Let $\phi \in C^1([0, 3]; [0, 1])$ be such that $\phi \equiv 1$ in $[0, 1]$ and $\phi \equiv 0$ in $[2, 3]$. We define the function $u_\varepsilon : \Omega_{\frac{\varepsilon}{2}}(\mu_\varepsilon) \rightarrow \mathbb{R}^2$ as

$$u_\varepsilon(x) := \begin{cases} u_\varepsilon^{R,k}(x) & \text{if } x \in A_{\frac{\varepsilon}{2}, \varepsilon^\gamma + \varepsilon}(x^{k,\varepsilon}) \text{ for some } k, \\ \phi\left(\frac{|x - x^{k,\varepsilon}|}{\varepsilon^\gamma + \varepsilon}\right) u_\varepsilon^{R,k}(x) + \left(1 - \phi\left(\frac{|x - x^{k,\varepsilon}|}{\varepsilon^\gamma + \varepsilon}\right)\right) u_\varepsilon^{\text{far}}(x) & \text{if } x \in A_{\varepsilon^\gamma + \varepsilon, 2(\varepsilon^\gamma + \varepsilon)}(x^{k,\varepsilon}) \text{ for some } k, \\ u_\varepsilon^{\text{far}}(x) & \text{if } x \in \Omega_{2(\varepsilon^\gamma + \varepsilon)}(\mu). \end{cases}$$

Furthermore, for every $k = 1, \dots, K$ we define the slip variable $\sigma^{k,\varepsilon} : \Omega_\varepsilon^1 \rightarrow \varepsilon \mathbb{T}$ as

$$\sigma^{k,\varepsilon}(i, j) := \begin{cases} -\xi^k & \text{if } C^{k,\varepsilon} \cap [i, j] \neq \emptyset \text{ and } \langle i - x^{k,\varepsilon}, e_2 \rangle < 0, \\ +\xi^k & \text{if } C^{k,\varepsilon} \cap [i, j] \neq \emptyset \text{ and } \langle i - x^{k,\varepsilon}, e_2 \rangle > 0, \\ 0 & \text{elsewhere,} \end{cases}$$

and we set $\sigma_\varepsilon := \sum_{k=1}^K \sigma^{k,\varepsilon}$. Finally, we define the map $\beta_\varepsilon \in \mathcal{S}_\varepsilon^{\text{discr}}(\Omega)$ as

$$\beta_\varepsilon(i, j) := R(j - i) + \varepsilon(u_\varepsilon(j) - u_\varepsilon(i) - \sigma_\varepsilon(i, j)).$$

We claim that $\{(\mu_\varepsilon; \beta_\varepsilon; R_\varepsilon)\}_\varepsilon$ is a recovery sequence. By construction, $\mu_\varepsilon = \mu[\beta_\varepsilon]$ and, by Remark 1.6 (in particular, by (1.13)), we have also that $\beta_\varepsilon \in \mathcal{AS}_\varepsilon^{\gamma, \text{discr}}(\mu_\varepsilon)$ for every $\varepsilon > 0$. Moreover, properties (0) and (1) are trivially satisfied. We show that this is the case also for (2). By the very definition of $\beta_{\mathbb{R}^2}^{\zeta, \mathbb{C}}$ in (1.11), there exists a universal constant $C > 0$ such that for every $\zeta \in \mathbb{T}$,

$$|\beta_{\mathbb{R}^2}^{\zeta, \mathbb{C}}(\rho, \theta)| \leq \frac{C}{\rho} |\zeta|^2, \quad |\nabla \beta_{\mathbb{R}^2}^{\zeta, \mathbb{C}}(\rho, \theta)| \leq \frac{C}{\rho^2} |\zeta|^2. \quad (2.12)$$

Therefore, recalling that $\beta^{\text{cf}} \in W^{1,\infty}(\Omega; \mathbb{R}^{2 \times 2})$, for every $T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega)$ with $T_\varepsilon \subset \bigcup_{k=1}^K A_{\frac{\varepsilon}{2}, \varepsilon^\gamma + \varepsilon}(x^{k,\varepsilon}) \cup \Omega_{2(\varepsilon^\gamma + \varepsilon)}(\mu_\varepsilon)$ and for every $x \in T_\varepsilon$ it holds

$$\begin{aligned} \left| \frac{\tilde{\beta}_\varepsilon^{T_\varepsilon} - R_\varepsilon}{\varepsilon} - \nabla u_\varepsilon(x) \right|^2 &\leq C\varepsilon^2 \|\nabla^2 u_\varepsilon\|_{L^\infty(T_\varepsilon; \mathbb{R}^{2 \times 2})}^2 \\ &\leq C\varepsilon^2 \left(\sum_{k=1}^K \|\nabla \beta_{\mathbb{R}^2}^{R^T \xi^k, C}\|_{L^\infty(T_\varepsilon; \mathbb{R}^{2 \times 2})}^2 + |\log \varepsilon| \|\nabla \beta^{\text{cf}}\|_{L^\infty(T_\varepsilon; \mathbb{R}^{2 \times 2})}^2 \right) \\ &\leq C\varepsilon^2 \left(\sum_{k=1}^K \frac{1}{|x - x^{k,\varepsilon}|^4} + |\log \varepsilon| \right) \end{aligned} \quad (2.13)$$

for some constant $C > 0$ independent of ε . Furthermore, by arguing verbatim as in the proof of (1.48), we have that, for every $k = 1, \dots, K$,

$$\|\nabla u_\varepsilon\|_{L^\infty(A_{\frac{\varepsilon^\gamma}{2}, 3\varepsilon^\gamma}(x^{k,\varepsilon}); \mathbb{R}^{2 \times 2})} + \left\| \frac{\tilde{\beta}_\varepsilon^{T_\varepsilon} - R_\varepsilon}{\varepsilon} \right\|_{L^\infty(A_{\frac{\varepsilon^\gamma}{2}, 3\varepsilon^\gamma}(x^{k,\varepsilon}); \mathbb{R}^{2 \times 2})} \leq \frac{C}{\varepsilon^\gamma}. \quad (2.14)$$

Therefore, by (2.13)–(2.14) and the fact that $|\Omega \setminus \Omega_{\mathcal{T}_\varepsilon}| \leq C\varepsilon$, we readily see that

$$\frac{1}{|\log \varepsilon|} \left\| \frac{\tilde{\beta}_\varepsilon^{T_\varepsilon} - R_\varepsilon}{\varepsilon} - \nabla u_\varepsilon \chi_{\Omega_\varepsilon(\mu_\varepsilon)} \right\|_{L^2(\Omega; \mathbb{R}^{2 \times 2})}^2 \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0. \quad (2.15)$$

Moreover, by arguing verbatim as in (1.47), (1.49) and (1.50), we have

$$\frac{\nabla u_\varepsilon \chi_{\Omega_\varepsilon(\mu_\varepsilon)}}{\sqrt{|\log \varepsilon|}} \rightharpoonup \beta^{\text{cf}} \quad \text{in } L^2(\Omega; \mathbb{R}^{2 \times 2}). \quad (2.16)$$

Therefore, by (2.15) and (2.16), we deduce that $(\mu_\varepsilon; \tilde{\beta}_\varepsilon^{T_\varepsilon}; R_\varepsilon) \rightarrow (\mu; \beta^{\text{cf}}; R)$.

Next we show that also (2.11) holds true. By construction we have

$$\tilde{E}_\varepsilon(\beta_\varepsilon; T_\varepsilon) \leq C\varepsilon^2 \quad \text{for every } T_\varepsilon \in \mathcal{T}_\varepsilon(\Omega) \text{ with } \min_{k=1, \dots, K} \text{dist}(x^{k,\varepsilon}, T_\varepsilon) \leq \varepsilon, \quad (2.17)$$

and that for every $(i, j) \in \Omega_\varepsilon^1$ with $i, j \in \partial\Omega_{\mathcal{T}_\varepsilon}$

$$\varepsilon^2 \psi_1\left(\frac{|\beta_\varepsilon(i, j)|^2}{\varepsilon^2}\right) \leq C\varepsilon^2. \quad (2.18)$$

Moreover, by (2.14) and (1.52), for every $k = 1, \dots, K$, we have

$$\frac{1}{\varepsilon^2 |\log \varepsilon|} \int_{A_{\frac{\varepsilon^\gamma}{2}, 3\varepsilon^\gamma}(x^{k,\varepsilon})} W(\tilde{\beta}_\varepsilon^{T_\varepsilon}) \, dx \leq \frac{1}{|\log \varepsilon|} \left\| \frac{\tilde{\beta}_\varepsilon^{T_\varepsilon} - R_\varepsilon}{\varepsilon} \right\|_{L^2(A_{\frac{\varepsilon^\gamma}{2}, 3\varepsilon^\gamma}(x^{k,\varepsilon}); \mathbb{R}^{2 \times 2})}^2 \leq \frac{C}{|\log \varepsilon|}. \quad (2.19)$$

Let $\alpha \in (\gamma, 1)$. By arguing as in the proof of (1.51) and (1.53), one can show that

$$\limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \int_{A_{\varepsilon^\alpha, \varepsilon^\gamma}(x^{k,\varepsilon})} W(\tilde{\beta}_\varepsilon^{T_\varepsilon}) \, dx = C(1 - \alpha) \quad (2.20)$$

for each $k = 1, \dots, K$. Moreover, since $\|\tilde{\beta}_\varepsilon^{T_\varepsilon} - R_\varepsilon\|_{L^\infty} \leq C\varepsilon^{1-\alpha}$ in $A_{\varepsilon^\alpha, \varepsilon^\gamma}(x^{k,\varepsilon})$, by (ii), writing the Taylor expansion of W around the identity, we find

$$\begin{aligned} \limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \int_{A_{\varepsilon^\alpha, \varepsilon^\gamma}(x^{k,\varepsilon})} W(\tilde{\beta}_\varepsilon^{T_\varepsilon}) \, dx &\leq \limsup_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \int_{A_{\varepsilon^\alpha, \varepsilon^\gamma}(x^{k,\varepsilon})} \frac{1}{2} \mathbf{C} \left(R^T \frac{\tilde{\beta}_\varepsilon^{T_\varepsilon} - R}{\varepsilon} \right) : \left(R^T \frac{\tilde{\beta}_\varepsilon^{T_\varepsilon} - R}{\varepsilon} \right) \, dx \\ &\quad + \limsup_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \int_{A_{\varepsilon^\alpha, \varepsilon^\gamma}(x^{k,\varepsilon})} \frac{\sigma(\tilde{\beta}_\varepsilon^{T_\varepsilon} - R)}{|\tilde{\beta}_\varepsilon^{T_\varepsilon} - R|^2} \left| \frac{\tilde{\beta}_\varepsilon^{T_\varepsilon} - R}{\varepsilon} \right|^2 \, dx \end{aligned} \quad (2.21)$$

with $\lim_{|M| \rightarrow 0} \frac{\sigma(M)}{|M|^2} = 0$. By (2.13) and (2.12), we have that

$$\left| \frac{\tilde{\beta}_\varepsilon^{\mathcal{J}_\varepsilon} - R}{\varepsilon} \right| \leq \frac{C}{|x - x^{k,\varepsilon}|} \quad \text{for every } x \in A_{\varepsilon^\alpha, \varepsilon^\gamma}(x^{k,\varepsilon}),$$

whence we deduce that the last lim sup in (2.21) is equal to zero. Therefore, by (2.21), using again (2.13), we get

$$\begin{aligned} & \limsup \frac{1}{\varepsilon^2 |\log \varepsilon|} \int_{A_{\varepsilon^\alpha, \varepsilon^\gamma}(x^{k,\varepsilon})} W(\tilde{\beta}_\varepsilon^{\mathcal{J}_\varepsilon}) \, dx \\ & \leq \limsup_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \int_{A_{\varepsilon^\alpha, \varepsilon^\gamma}(x^{k,\varepsilon})} \frac{1}{2} C(R^T \nabla u_\varepsilon) : (R^T \nabla u_\varepsilon) + C\varepsilon^2 \left(\frac{1}{|x - x^{k,\varepsilon}|^4} + |\log \varepsilon| \right) \, dx \\ & = \limsup_{\varepsilon \rightarrow 0} \frac{1}{|\log \varepsilon|} \int_{A_{\varepsilon^\alpha, \varepsilon^\gamma}(0)} \frac{1}{2} C \beta_{\mathbb{R}^2}^{R^T \xi^k, C} : \beta_{\mathbb{R}^2}^{R^T \xi^k, C} \, dx \\ & = (\alpha - \gamma) \psi^C(R^T \xi^k), \end{aligned} \tag{2.22}$$

where the last equality follows by (1.14). Furthermore, similar arguments yield

$$\limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2 |\log \varepsilon|} \int_{\Omega_{2\varepsilon^\gamma}(\mu_\varepsilon)} W(\tilde{\beta}_\varepsilon^{\mathcal{J}_\varepsilon}) \, dx \leq \gamma \sum_{k=1}^K \psi^C(R^T \xi^k) + \frac{1}{2} \int_{\Omega} C R^T \beta^{\text{cf}} : R^T \beta^{\text{cf}} \, dx. \tag{2.23}$$

Finally, combining (2.3), (2.17)–(2.20), (2.22)–(2.23), and sending $\alpha \rightarrow 1$, we obtain (2.11). \square

Acknowledgment: The authors thank G. Lazzaroni for interesting and fruitful discussions on discrete nonlinear elasticity models.

Funding: Roberto Alicandro, Lucia De Luca, and Mariapia Palombaro are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

References

- [1] R. Alicandro and M. Cicalese, Variational analysis of the asymptotics of the XY model, *Arch. Ration. Mech. Anal.* **192** (2009), no. 3, 501–536.
- [2] R. Alicandro, M. Cicalese and M. Ponsiglione, Variational equivalence between Ginzburg–Landau, XY spin systems and screw dislocations energies, *Indiana Univ. Math. J.* **60** (2011), no. 1, 171–208.
- [3] R. Alicandro, L. De Luca, A. Garroni and M. Ponsiglione, Metastability and dynamics of discrete topological singularities in two dimensions: A Γ -convergence approach, *Arch. Ration. Mech. Anal.* **214** (2014), no. 1, 269–330.
- [4] R. Alicandro, L. De Luca, G. Lazzaroni, M. Palombaro and M. Ponsiglione, Coarse-graining of a discrete model for edge dislocations in the regular triangular lattice, *J. Nonlinear Sci.* **33** (2023), no. 2, Paper No. 33.
- [5] R. Alicandro, G. Lazzaroni and M. Palombaro, Derivation of linear elasticity for a general class of atomistic energies, *SIAM J. Math. Anal.* **53** (2021), no. 5, 5060–5093.
- [6] M. P. Ariza and M. Ortiz, Discrete crystal elasticity and discrete dislocations in crystals, *Arch. Ration. Mech. Anal.* **178** (2005), 149–226.
- [7] D. J. Bacon, D. M. Barnett and R. O. Scattergood, Anisotropic continuum theory of lattice defects, *Progr. Mater. Sci.* **23** (1978), 51–262.
- [8] A. Braides, M. Solci and E. Vitali, A derivation of linear elastic energies from pair-interaction atomistic systems, *Netw. Heterog. Media* **2** (2007), no. 3, 551–567.
- [9] P. Cermelli and G. Leoni, Renormalized energy and forces on dislocations, *SIAM J. Math. Anal.* **37** (2005), no. 4, 1131–1160.
- [10] S. Conti, A. Garroni and R. Marziani, Line-tension limits for line singularities and application to the mixed-growth case, *Calc. Var. Partial Differential Equations* **62** (2023), no. 8, Paper No. 228.
- [11] G. Dal Maso, M. Negri and D. Percivale, Linearized elasticity as Γ -limit of finite elasticity, *Set-Valued Anal.* **10** (2002), 165–183.
- [12] R. Dautray and J.-L. Lions, *Mathematical Analysis and Numerical Methods for Science and Technology. Vol. 3*, Springer, Berlin, 1988.
- [13] L. De Luca, Γ -convergence analysis for discrete topological singularities: The anisotropic triangular lattice and the long range interaction energy, *Asymptot. Anal.* **96** (2016), no. 3–4, 185–221.

- [14] L. De Luca, A. Garroni and M. Ponsiglione, Γ -convergence analysis of systems of edge dislocations: The self energy regime, *Arch. Ration. Mech. Anal.* **206** (2012), no. 3, 885–910.
- [15] G. Friesecke, R. D. James and S. Müller, A theorem on geometric rigidity and the derivation of nonlinear plate theory from three-dimensional elasticity, *Comm. Pure Appl. Math.* **55** (2002), no. 11, 1461–1506.
- [16] A. Garroni, G. Leoni and M. Ponsiglione, Gradient theory for plasticity via homogenization of discrete dislocations, *J. Eur. Math. Soc. (JEMS)* **12** (2010), no. 5, 1231–1266.
- [17] A. Giuliani and F. Theil, Long range order in atomistic models for solids, *J. Eur. Math. Soc. (JEMS)* **24** (2022), no. 10, 3505–3555.
- [18] J. P. Hirth and J. Lothe, *Theory of Dislocations*, Krieger Publishing, Malabar, 1982.
- [19] S. Luckhaus and L. Mugnai, On a mesoscopic many-body Hamiltonian describing elastic shears and dislocations, *Contin. Mech. Thermodyn.* **22** (2010), no. 4, 251–290.
- [20] S. Müller, L. Scardia and C. I. Zeppieri, Gradient theory for geometrically nonlinear plasticity via the homogenization of dislocations, in: *Analysis and Computation of Microstructure in Finite Plasticity*, Lect. Notes Appl. Comput. Mech. 78, Springer, Cham (2015), 175–204.
- [21] M. Ponsiglione, Elastic energy stored in a crystal induced by screw dislocations: From discrete to continuous, *SIAM J. Math. Anal.* **39** (2007), no. 2, 449–469.
- [22] C. Reina and S. Conti, Kinematic description of crystal plasticity in the finite kinematic framework: a micromechanical understanding of $\mathbf{F} = \mathbf{F}^e \mathbf{F}^p$, *J. Mech. Phys. Solids* **67** (2014), 40–61.
- [23] C. Reina, A. Schlömerkemper and S. Conti, Derivation of $\mathbf{F} = \mathbf{F}^e \mathbf{F}^p$ as the continuum limit of crystalline slip, *J. Mech. Phys. Solids* **89** (2016), 231–254.
- [24] E. Sandier, Lower bounds for the energy of unit vector fields and applications, *J. Funct. Anal.* **152** (1998), no. 2, 379–403.
- [25] E. Sandier and S. Serfaty, *Vortices in the magnetic Ginzburg–Landau model*, Progr. Nonlinear Differential Equations Appl. 70, Birkhäuser, Boston, 2007.
- [26] L. Scardia and C. I. Zeppieri, Line-tension model for plasticity as the Γ -limit of a nonlinear dislocation energy, *SIAM J. Math. Anal.* **44** (2012), no. 4, 2372–2400.
- [27] B. Schmidt, On the derivation of linear elasticity from atomistic models, *Netw. Heterog. Media* **4** (2009), no. 4, 789–812.