

Increased Carotid Intima-Media Thickness and Stiffness in Obese Children

ARCANGELO IANNUZZI, MD¹
 MARIA ROSARIA LICENZIATI, MD²
 CIRO ACAMPORA, MD³
 VITTORIO SALVATORE, MD¹
 LUCIA AURIEMMA²

MARIA LUGIA ROMANO, MD³
 SALVATORE PANICO, MD⁴
 PAOLO RUBBA, MD⁴
 MAURIZIO TREVISAN, MD⁵

Obesity in childhood increases the risk of atherosclerotic disease and death in adulthood (1). A dramatic increase in overweight among children and adolescents during the past 2 decades has been documented (2). Moreover, overweight children and adolescents have an increased risk of adult obesity (3). A clustering of factors typical of the insulin resistance syndrome has been identified in 5- to 10-year-old overweight/obese African-American children (4) and in pre-adolescent obese children (5). The pediatric obesity epidemic accounts for most new diagnoses of type 2 diabetes, a disease once virtually confined to adulthood, in adolescence (6).

High-resolution B-mode ultrasound measurements of carotid intima-media thickness (IMT) and stiffness are markers of early, preclinical atherosclerosis. Previous observations show significantly increased IMT in familial hypercholesterolemic children (7,8) and in children with type 1 diabetes (9) and hypertension (10). Recent evidence suggests an increased arterial stiffness in familial hypercholesterolemic children (11) and in children with severe obesity (12).

However, in the study by Tounian et al. (12) there was no evidence of statistically significant differences in carotid IMT between severely obese children and control subjects; moreover, there was no sub-

group analyses for boys versus girls. The present study aimed to verify whether obesity in childhood may cause premature vascular alterations in both sexes.

RESEARCH DESIGN AND METHODS

— We studied 100 children with obesity consecutively recruited from the outpatient clinic of the Department of Pediatrics, A. Cardarelli Hospital, Naples, Italy, among patients evaluated for overweight or obesity and 47 healthy age-matched control subjects. The inclusion criteria for both obese children and control subjects were age 6–14 years; no personal history of diabetes or impaired fasting glucose, hypercholesterolemia, or hypertension; absence of any pharmacological therapy; and absence of cardiovascular disease in their parents. Obesity was defined as BMI >95th percentile of the reference values stated in the Centers for Disease Control and Prevention growth chart (13).

All of the children had measurements of fasting serum total cholesterol, triglycerides, apolipoproteins A and B, HDL cholesterol, plasma glucose and insulin, C-reactive protein, and glycated HbA_{1c}. An oral glucose tolerance test was performed in the obese children. The estimate of insulin resistance was calculated

by a homeostasis model assessment (HOMA) score, as described by Matthews et al. (14). Quantitative B-mode ultrasound measurements of common carotid IMT and stiffness were calculated for all children following a standardized protocol. Carotid stiffness was calculated using the following formula: $\beta = (\text{natural logarithm systolic blood pressure} - \text{natural logarithm diastolic blood pressure}) / (\text{systolic diameter} - \text{diastolic diameter}) / \text{diastolic diameter}$.

Statistical analysis

Statistical analysis was performed using the Statistical Package for Social Sciences (SPSS 11). Univariate, unadjusted analyses between obese and control subjects were performed with the independent samples *t* test. Stiffness measures were not normally distributed, and a Kruskal-Wallis test was used. ANCOVA was used to evaluate the presence of confounding variables in the relationship between obesity status and vascular parameters. Models were adjusted for several confounding variables, including age, sex, systolic blood pressure, total cholesterol, HDL cholesterol, and triglycerides. Adjusted means and CIs were estimated with the use of the Bonferroni method. The coefficient of variation for each pair of IMT and diameter measurements was 3.9% for IMT measurements, 1.8% for lumen systolic diameters, and 2.5% for lumen diastolic diameters.

RESULTS— Table 1 shows anthropometrical, biochemical, and ultrasonic parameters of the arterial carotid wall in obese and nonobese children. Obese children had significantly higher blood pressure and plasma concentrations of triglycerides, cholesterol, glucose, insulin, HOMA, and C-reactive protein than control subjects. Carotid thickness and stiffness were significantly different between obese and nonobese children and in both boys and girls.

For carotid IMT, when we added traditional cardiovascular risk factors (except for blood pressure and HOMA) in

From the ¹Department of Medicine, Cava de' Tirreni Hospital, Salerno, Italy; the ²Department of Pediatrics, A. Cardarelli Hospital, Naples, Italy; the ³Department of Radiology, A. Cardarelli Hospital, Naples, Italy; the ⁴Department of Clinical and Experimental Medicine, School of Medicine, Federico II University, Naples, Italy; and the ⁵Department of Social and Preventive Medicine, School of Public Health and Health Professions, State University of New York, Buffalo, New York.

Address correspondence and reprint requests to Dr. Arcangelo Iannuzzi, MD, Cava de' Tirreni Hospital, Department of Medicine, Via de Marinis, SS18, 84013 Cava de' Tirreni, Salerno, Italy. E-mail: lelliann@libero.it.

Received for publication 15 May 2004 and accepted in revised form 30 June 2004.

Abbreviations: HOMA, homeostasis model assessment; IMT, intima-media thickness.

A table elsewhere in this issue shows conventional and Système International (SI) units and conversion factors for many substances.

© 2004 by the American Diabetes Association.

Table 1—Physical characteristics and biochemical and ultrasonographic parameters

	Entire group			Boys			Girls		
	Obese	Control subjects	P	Obese	Control subjects	P	Obese	Control subjects	P
n	100	47	—	61	26	—	39	21	—
Age (years)	10.0 ± 2.7	10.0 ± 2.5	0.82	10.3 ± 2.8	11.1 ± 1.9	0.13	9.6 ± 2.4	8.8 ± 2.6	0.22
Weight (kg)	58.6 ± 21.4	34.8 ± 10.0	—	62.4 ± 24.7	38.5 ± 9.3	—	52.7 ± 13.2	30.2 ± 9.1	—
Height (cm)	143.2 ± 14.9	138.7 ± 16.2	0.097	145.5 ± 16.2	146.1 ± 11.8	0.86	139.6 ± 11.9	129.4 ± 16.4	0.02
BMI (kg/m ²)	27.8 ± 5.3	17.8 ± 2.6	—	28.5 ± 5.9	17.9 ± 2.8	—	26.6 ± 3.9	17.6 ± 2.3	—
Systolic blood pressure (mmHg)	120.6 ± 15.8	98.8 ± 13.8	0.001	122.4 ± 16.9	103.4 ± 10.6	0.001	117.9 ± 13.6	93.0 ± 15.4	0.001
Diastolic blood pressure (mmHg)	76.4 ± 8.0	65.4 ± 8.3	0.001	77.1 ± 8.8	69.4 ± 6.2	0.001	75.4 ± 6.6	60.5 ± 8.0	0.001
Fasting cholesterol (mmol/l)	4.25 ± 0.74	3.96 ± 0.52	0.008	4.14 ± 0.78	3.99 ± 0.61	0.38	4.42 ± 0.65	3.93 ± 0.39	0.001
Fasting triglycerides (mmol/l)	0.91 ± 0.41	0.67 ± 0.22	0.001	0.93 ± 0.46	0.73 ± 0.24	0.008	0.87 ± 0.30	0.61 ± 0.18	0.001
Plasma fasting glucose (mmol/l)	4.55 ± 0.51	4.37 ± 0.45	0.036	4.62 ± 0.46	4.37 ± 0.52	0.030	4.45 ± 0.56	4.36 ± 0.38	0.50
Insulin (μU/ml)	16.1 ± 11.4	7.1 ± 2.1	0.001	16.4 ± 13.3	7.5 ± 2.3	0.001	15.6 ± 7.6	6.7 ± 1.8	0.001
HOMA	3.2 ± 2.1	1.4 ± 0.5	0.001	3.3 ± 2.3	1.5 ± 0.5	0.003	3.1 ± 1.6	1.3 ± 0.4	0.001
C-reactive protein	1.45 ± 0.78	1.26 ± 0.08	0.001	1.49 ± 0.45	1.26 ± 0.09	0.001	1.39 ± 0.06	1.26 ± 0.06	0.004
IMT (mm)	0.55 ± 0.08	0.49 ± 0.09	0.001	0.56 ± 0.08	0.48 ± 0.10	0.001	0.53 ± 0.09	0.49 ± 0.07	0.045
Stiffness	3.48 ± 1.26	2.38 ± 0.61	0.001	3.65 ± 1.3	2.43 ± 0.64	0.001	3.19 ± 1.17	2.32 ± 0.59	0.004

Data are means ± SD.

addition to age and sex as covariates, we still found evidence of statistically significant differences between obese children and control subjects. IMT in obese children was 0.55 mm (95% CI 0.54–0.57) and in healthy control subjects was 0.48 mm (0.46–0.51, $P = 0.001$). The further addition of C-reactive protein did not materially modify the outcome of our analyses. Adding systolic blood pressure as a covariate reduced the P value, but the statistical significance was retained. However, when glucose level was substituted by HOMA, the statistical significance was further reduced to a level that did not reach the formal level of statistical significance. IMT in obese children was 0.54 mm (0.52–0.56), and IMT in healthy control subjects was 0.51 mm (0.48–0.54, $P = 0.099$).

CONCLUSIONS— Obese children have significantly increased carotid IMT and stiffness compared with healthy control children. A previous study demonstrated an increased stiffness of the carotid artery in obese children (12) (a finding confirmed in the present study) but concluded that no significant differ-

ence was present in IMT between obese and nonobese children. The apparent discrepancy with the present observation could be due to the limited statistical power available for the study by Tounian et al. (12), which focuses on much smaller sample sizes (48 obese children and 27 control subjects). Increased carotid IMT and stiffness in obese boys and girls could reflect structural changes of large arteries that occur very early in atherosclerosis. The increased vascular IMT and stiffness in obese children compared with healthy control subjects suggests that obesity in children represents a powerful determinant of early manifestations of atherosclerosis and affects structural and mechanical properties of major vessels. The effect on structural changes appears to be mediated, at least in part, by the risk factors considered, especially systolic blood pressure and, most importantly, insulin resistance. Based on these findings, children affected by the insulin resistance syndrome are at particularly higher risk of developing premature cardiovascular complications. Our observation confirms that obesity and its associated comorbidities, particularly hypertension and insulin resistance, even in this young age should

be regarded as a disease with vascular implications. We emphasize that weight reduction and risk-factor control in obese children for the long-term prevention of atherosclerosis and its sequelae should begin in childhood.

References

1. Must A, Jacques PF, Dallal GE, Bajema CJ, Dietz WH: Long-term morbidity and mortality of overweight adolescents: a follow-up of the Harvard Growth Study of 1922 to 1935. *N Engl J Med* 327:1350–1355, 1992
2. Freedman DS, Srinivasan SR, Valdez RA, Williamson DF, Berenson GS: Secular increases in relative weight and adiposity among children over two decades: the Bogalusa Heart Study. *Pediatrics* 99:420–426, 1997
3. Serdula MK, Ivery D, Coates RJ, Freedman DS, Williamson DF, Byers T: Do obese children become obese adults? A review of literature. *Prev Med* 22:167–177, 1993
4. Young-Hyman D, Schlundt DG, Herman L, De Luca F, Counts D: Evaluation of the insulin resistance syndrome in 5- to 10-year-old overweight/obese African-American children. *Diabetes Care* 24:1359–1364, 2001
5. Caprio S, Bronson M, Sherwin RS, Rife F,

- Tamborlane WV: Coexistence of severe insulin resistance and hyperinsulinaemia in pre-adolescent obese children. *Diabetologia* 39:1489–1497, 1996
- Ludwig DS, Ebbeling CB: Type 2 diabetes mellitus in children: primary care and public health considerations. *JAMA* 286:1427–1430, 2001
 - Pauciullo P, Iannuzzi A, Sartorio R, Irace C, Covetti G, Di Costanzo A, Rubba P: Increased intima-media thickness of the common carotid artery in hypercholesterolemic children. *Arterioscler Thromb* 14:1075–1079, 1994
 - Tonstad S, Joakimsen O, Stensland-Bugge E, Leren TP, Ose L, Russell D, Bonna KH: Risk factors related to carotid intima-media thickness and plaque in children with familial hypercholesterolemia and control subjects. *Arterioscler Thromb Vasc Biol* 16:984–991, 1996
 - Jarvisalo MJ, Jartti L, Nanto-Salonen K, Irjala K, Ronnema T, Hartiala JJ, Celermajer DS, Raitakari OT: Increased aortic intima-media thickness: a marker of pre-clinical atherosclerosis in high-risk children. *Circulation* 104:2943–2947, 2001
 - Sorof JM, Alexandrov AV, Cardwell G, Portman RJ: Carotid artery IMT and LVH in children with elevated blood pressure. *Pediatrics* 111:61–66, 2003
 - Aggoun Y, Bonnet D, Sidi D, Girardet JP, Brucker E, Polak M, Safar ME, Levy BI: Arterial mechanical changes in children with familial hypercholesterolemia. *Arterioscler Thromb Vasc Biol* 20:2070–2075, 2000
 - Tounian P, Aggoun Y, Dubem B, Varille V, Guy-Grand B, Sidi D, Girardet JP, Bonnet D: Presence of increased stiffness of the common carotid artery and endothelial dysfunction in severely obese children: a prospective study. *Lancet* 358:1400–1404, 2001
 - Kuczmarski RJ, Ogden CL, Guo SS, Grummer-Strawn LM, Flegal KM, Mei Z, Wei R, Curtin LR, Roche AF, Johnson CL: 2000 CDC growth charts for the United States: methods and development. *Vital Health Stat* 11:1–190, 2002
 - Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC: Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28:412–419, 1985