

# On the homology of the universal Steenrod algebra at odd primes

A. Ciampella

Received: 19 May 2014 / Accepted: 21 June 2014 / Published online: 30 July 2014  
© Università degli Studi di Napoli “Federico II” 2014

**Abstract** We give an explicit description of the homology  $H_*(\mathcal{Q})$  of the universal Steenrod algebra  $\mathcal{Q}$  for any odd prime  $p$ , extending the work done for the  $p = 2$  case. We also exhibit an isomorphism with a certain coalgebra of invariants  $\Gamma$ .

**Keywords** Universal Steenrod algebra · Invariant theory · Koszul algebras · Bar resolutions · Homology of algebras · Modular invariants

**Mathematics Subject Classification** 13A50 · 55S10

## 1 Introduction

Let  $p$  be any prime and  $\mathbb{F}_p$  the field with  $p$  elements. In [14] May introduced the universal Steenrod algebra  $\mathcal{Q}$  over the field  $\mathbb{F}_p$  as the algebra of all cohomology operations in the category of  $H_\infty$ -ring spectra. This algebra is also known as the *algebra of all generalized Steenrod operations* [13] or the *extended Steenrod algebra* [6]. It is closely related to other well known algebras, such as the opposite  $\Lambda^{opp}$  of the  $\Lambda$  algebra introduced in [1], the Steenrod algebra  $\mathcal{A}$  in [15] and the Steenrod algebra for simplicial restricted Lie algebras  $A_L$  in [14]. The algebra  $\mathcal{Q}$  has extensively been studied in [2–5, 7, 8] and [12]. In particular the papers [7] and [12] contain an invariant-theoretic description of  $\mathcal{Q}$  and a computation of the diagonal cohomology  $D^*(\mathcal{Q}) = \bigoplus \text{Ext}_{\mathcal{Q}}^{q,q}(\mathbb{F}_p, \mathbb{F}_p)$  for  $p = 2$  and for any odd prime  $p$ , respectively. In [4] the authors prove that  $\mathcal{Q}$ , which is a non-locally finite homogeneous quadratic algebra, is an example of a *good PBW*-algebra, hence it is *koszul*, i.e. the cohomology  $H^{**}(\mathcal{Q}) =$

---

Communicated by Salvatore Rionero.

---

A. Ciampella (✉)  
Dipartimento di Matematica e applicazioni, Università di Napoli “Federico II”,  
Piazzale Tecchio 80, 80125 Napoli, Italy  
e-mail: ciampell@unina.it

$\oplus Ext_{\mathcal{Q}}^{s,t}(\mathbb{F}, \mathbb{F})$  is purely diagonal. The analogous for the homology  $H_{*,*}(\mathcal{Q})$  holds:  $Tor_{\mathcal{Q}}^{s,t}(\mathbb{F}_p, \mathbb{F}_p) = 0$  if  $s \neq t$ . In [8] there is an explicit description of the non trivial part of the homology of  $\mathcal{Q}$ ,  $D_*(\mathcal{Q}) = \oplus H_{q,q}(\mathcal{Q})$ , for the  $p = 2$  case. The classical bar construction is used to get our main results: the mod  $p$  homology of  $\mathcal{Q}$  and its description in terms of invariant theory. In Sect. 2 we recall the basic results of invariant theory. Section 3 is devoted to the computation of the diagonal homology  $D_*(\mathcal{Q})$  of  $\mathcal{Q}$ . In Sect. 4 we prove that  $D_*(\mathcal{Q})$  is isomorphic to the coalgebra  $\Gamma$  of certain invariants with respect to the action of the general linear group  $GL(n, \mathbb{F}_p)$ .

### 2 Preliminaries on invariant theory

Let  $\Phi_n = (E(x_1, \dots, x_n) \otimes \mathbb{F}_p[y_1, \dots, y_n])[L_n^{-1}]$  be the localization out of the Euler class  $L_n$  of  $H^*(B(\mathbb{Z}_p)^n)$ . The general linear group  $GL_n = GL(n, \mathbb{F}_p)$  acts on  $\Phi_n$ . Let  $B_n$  be the Borel subgroup of  $GL_n$  of all upper triangular matrices and  $T_n$  the subgroup of  $B_n$  of all matrices with 1 on the main diagonal. We are interested on the invariant rings

$$\Delta_n = \Phi_n^{T_n}, \quad \bar{\Delta}_n = \Phi_n^{B_n}, \quad \Gamma_n = \Phi_n^{GL_n}.$$

We know from [9] (Corollary 1.3) that

$$\Delta_n = E(u_1, \dots, u_n) \otimes \mathbb{F}_p[v_1^{\pm 1}, \dots, v_n^{\pm 1}],$$

where  $u_i$  and  $v_i$  have degree  $|u_i| = 1, |v_i| = 2$  for  $1 \leq i \leq n$ .

Combining results from [11] and [7] (Proposition 1.2), we get

$$\bar{\Delta}_n = E(\bar{u}_1, \dots, \bar{u}_n) \otimes \mathbb{F}_p[w_1^{\pm 1}, \dots, w_n^{\pm 1}],$$

where

$$\begin{aligned} \bar{u}_i &= (-1)^i u_i v_i^{-1}, & w_i &= v_i^{p-1}, \\ |\bar{u}_i| &= -1, & |w_i| &= 2(p-1), \end{aligned} \tag{2.1}$$

for  $1 \leq i \leq n$ .

In [9] (Corollary 1.2, item (ii)), we find

$$\Gamma_n = E(R_{n,0}, \dots, R_{n,n-1}) \otimes \mathbb{F}_p[Q_{n,0}^{\pm 1}, Q_{n,1}, \dots, Q_{n,n-1}],$$

$|R_{n,i}| = 2(p^n - p^i) - 1, |Q_{n,i}| = 2(p^n - p^i)$ . Moreover, the following proposition gives recursive formulas for the generators of  $\Gamma_n$ .

**Proposition 2.1** (1)  $Q_{n,i} = Q_{n-1,i-1}^p + Q_{n-1,0}^{p-1} Q_{n-1,i} v_n^{p-1} = Q_{n-1,i-1}^p + Q_{n-1,0}^{p-1} Q_{n-1,i} w_n$ .

(2)  $R_{n,i} = Q_{n-1,0}^{p-1}(R_{n-1,i}v_n^{p-1} + Q_{n-1,i}u_nv_n^{p-2})$ ; in terms of  $B_n$  invariants,  $R_{n,i} = Q_{n-1,0}^{p-1}(R_{n-1,i} + (-1)^n Q_{n-1,i}\bar{u}_n)w_n$ .

*Proof* See Proposition 1.4 in [9]. □

By convention,  $Q_{n,i} = 0$  for either  $i < 0$  or  $n < i$ ,  $Q_{n,n} = 1$  and  $R_{n,i} = 0$  for either  $i < 0$  or  $i \geq n$ .

In particular we have

$$Q_{1,0} = v_1^{p-1} = w_1, \quad R_{1,0} = u_1v_1^{p-2} = -\bar{u}_1w_1, \tag{2.2}$$

$$Q_{2,0} = Q_{1,0}^pv_2^{p-1} = w_1^pw_2,$$

$$Q_{2,1} = Q_{1,0}^p + Q_{1,0}^{p-1}v_2^{p-1} = w_1^{p-1}(w_1 + w_2),$$

$$R_{2,0} = Q_{1,0}^{p-1}(R_{1,0}v_2^{p-1} + Q_{1,0}u_2v_2^{p-2}) = (\bar{u}_2 - \bar{u}_1)w_1^pw_2, \tag{2.3}$$

$$R_{2,1} = Q_{1,0}^{p-1}u_2v_2^{p-2} = \bar{u}_2w_1^{p-1}w_2.$$

Set

$$\Delta = \bigoplus_{n \geq 0} \Delta_n, \quad \bar{\Delta} = \bigoplus_{n \geq 0} \bar{\Delta}_n, \quad \Gamma = \bigoplus_{n \geq 0} \Gamma_n.$$

Here,  $\Delta_0 = \bar{\Delta}_0 = \Gamma_0 = \mathbb{F}_p$ .

For any non-negative integers  $n, q, t$  such that  $q + t = n$ , we define

$$\psi_{q,t} : \Delta_n \rightarrow \Delta_q \otimes \Delta_t$$

by setting

$$\psi_{q,t}(u_i) = \begin{cases} u_i \otimes 1, & 1 \leq i \leq q \\ 1 \otimes u_{i-q}, & q < i \leq n, \end{cases} \quad \psi_{q,t}(v_i) = \begin{cases} v_i \otimes 1, & 1 \leq i \leq q \\ 1 \otimes v_{i-q}, & q < i \leq n. \end{cases} \tag{2.4}$$

The map  $\psi_{q,t}$  turns out to be an isomorphism of algebras and  $\Delta$  turns out to be a coalgebra with comultiplication  $\psi : \Delta \rightarrow \Delta \otimes \Delta$  induced by the maps  $\psi_{q,t}$  and defined by

$$\psi(\delta) = \sum_{q+t=n} \psi_{q,t}(\delta)$$

for any  $\delta \in \Delta_n$ .

**Proposition 2.2** *For any  $q, t, n$  such that  $q + t = n$ ,  $\psi_{q,t}(\bar{\Delta}_n) \subset \bar{\Delta}_q \otimes \bar{\Delta}_t$ , so  $\bar{\Delta}$  is a subcoalgebra of  $\Delta$ .*

*Proof* According to (2.1),

$$\psi_{q,t}(\bar{u}_i) = \begin{cases} \bar{u}_i \otimes 1, & 1 \leq i \leq q \\ 1 \otimes \bar{u}_{i-q}, & q < i \leq n, \end{cases} \quad \psi_{q,t}(w_i) = \begin{cases} w_i \otimes 1, & 1 \leq i \leq q \\ 1 \otimes w_{i-q}, & q < i \leq n. \end{cases} \tag{2.5}$$

□

**Proposition 2.3** For any  $q, t, n$  such that  $q + t = n$ , the following relations hold

- (1)  $\psi_{q,t}(Q_{n,i}) = \sum_{j \geq 0} Q_{q,0}^{p^t - p^j} Q_{q,i-j}^{p^j} \otimes Q_{t,j};$
- (2)  $\psi_{q,t}(R_{n,i}) = Q_{q,0}^{p^t - 1} R_{q,i} \otimes Q_{t,0} + \sum_{j \geq 0} Q_{q,0}^{p^t - p^j} Q_{q,i-j}^{p^j} \otimes R_{t,j}.$

*Proof* See Proposition 3.3 in [9]. □

**Corollary 2.4**  $\psi(\Gamma) \subset \Gamma \otimes \Gamma$ , so  $\Gamma$  is a subcoalgebra of  $\overline{\Delta}$  and of  $\Delta$ .

*Proof* For any  $f \in \Gamma_n$ ,  $\psi(f) = \sum_{q+t=n} \psi_{q,t}(f)$  belongs to  $\Gamma_q \otimes \Gamma_t$ , so the restriction to  $\Gamma$  of the comultiplication  $\psi : \Delta \rightarrow \Delta \otimes \Delta$  defines a comultiplication on  $\Gamma$ . □

### 3 The homology of $\mathcal{Q}$ over $\mathbb{F}_p$

The universal Steenrod algebra  $\mathcal{Q}$  at odd primes is generated as an  $\mathbb{F}_p$ -algebra by

$$\mathcal{F} = \{z_{\varepsilon,i} \mid \varepsilon \in \{0, 1\}, i \in \mathbb{Z}\} \cup \{1\} \quad \text{with} \quad \deg z_{\varepsilon,i} = 2i(p - 1) + \varepsilon,$$

subject to the following generalized Adem relations:

$$z_{\varepsilon,pk-1-n}z_{0,k} = \sum_j A(n,j) z_{\varepsilon,pk-1-j}z_{0,k-n+j}, \tag{3.1}$$

$$z_{1-\varepsilon,pk-n}z_{1,k} = \sum_j A(n,j) z_{1-\varepsilon,pk-j}z_{1,k-n+j} + \varepsilon \sum_j B(n,j) z_{1,pk-j}z_{0,k-n+j}, \tag{3.2}$$

for each  $(k, n) \in \mathbb{Z} \times \mathbb{N}_0$ , where  $A(n, j)$  and  $B(n, j)$  are respectively equal to

$$(-1)^{j+1} \binom{(p-1)(n-j)-1}{j} \quad \text{and} \quad (-1)^j \binom{(p-1)(n-j)}{j}.$$

Such presentation already appeared in [7] where the authors also proved that

$$\mathcal{B} = \{z_{\varepsilon_1,i_1} \dots z_{\varepsilon_h,i_h} \mid i_j \geq pi_{j+1} + \varepsilon_{j+1} \text{ for each } j = 1, \dots, h-1\} \cup \{1\}$$

is a basis of  $\mathcal{Q}$ , called the basis of *admissible* monomials.

Let  $T$  denote the associative algebra freely generated by the  $\mathbb{F}_p$ -module with basis  $\mathcal{F}$ . We consider the map  $d : T \rightarrow T$  given by  $d(z_{\varepsilon,i}) = z_{\varepsilon,i-1}$ , such that  $d(\tau_1 \tau_2) = d(\tau_1) \tau_2 + \tau_1 d(\tau_2)$  for any  $\tau_1, \tau_2 \in T$ . Then  $d$  is a derivation in  $T$ . We write  $d^s$  for the  $s$ -iterated of  $d$ . Let  $L$  be the two-sided ideal generated by the set

$$\{d^s(z_{0,ph-1}z_{0,h}), d^s(z_{1,ph-1}z_{0,h}), d^s(z_{0,ph}z_{1,h} - z_{1,ph}z_{0,h}), d^s(z_{1,ph}z_{1,h})\}$$

for all  $s \in \mathbb{N}_0, h \in \mathbb{Z}$ .

**Proposition 3.1** *The algebras  $\mathcal{Q}$  and  $T/L$  are isomorphic.*

*Proof*  $\mathcal{Q}$  and  $T/L$  are both isomorphic to  $\overline{\Delta}/(\Gamma_2)$ ; the isomorphisms are established by Proposition 2.5 and Proposition 2.3 of [7], respectively.  $\square$

If  $z_{\varepsilon_1, i_1} \dots z_{\varepsilon_h, i_h} \in \mathcal{B}$ , the string  $I = ((\varepsilon_1, i_1), (\varepsilon_2, i_2), \dots, (\varepsilon_h, i_h)) \in (\{0, 1\} \times \mathbb{Z})^h$  will be called the *label* of  $z_{\varepsilon_1, i_1} \dots z_{\varepsilon_h, i_h}$  and we write  $z_I$  instead of  $z_{\varepsilon_1, i_1} \dots z_{\varepsilon_h, i_h}$ . We say that  $z_I$  has *length*  $h$  and *total degree*  $\varepsilon_1 + \dots + \varepsilon_h + 2(p - 1)(i_1 + \dots + i_h)$ ; hence  $\mathcal{Q}$  is a bigraded algebra. It is also an augmented algebra through the map  $\varepsilon : \mathcal{Q} \rightarrow \mathbb{F}_p$  which vanishes on the monomials of positive length and is the identity over  $\mathbb{F}_p \subset \mathcal{Q}$ . Let us denote by  $J$  the augmentation ideal  $J = \ker(\varepsilon)$ .

Let  $\overline{B}(\mathcal{Q}) = T(J) = \bigoplus_{s \in \mathbb{N}_0} J \otimes \dots \otimes J$ . Thus  $\overline{B}(\mathcal{Q})$  is generated by elements of the form  $z_{I_1} \otimes \dots \otimes z_{I_s}$  where  $z_{I_j} \in J$ . Such elements are written simply as

$$[z_{I_1} | \dots | z_{I_s}] = [z_{\varepsilon_1, i_1} \dots z_{\varepsilon_{t_1}, i_{t_1}} | z_{\varepsilon_{t_1+1}, i_{t_1+1}} \dots z_{\varepsilon_{t_2}, i_{t_2}} | \dots | z_{\varepsilon_{t_{s-1}+1}, i_{t_{s-1}+1}} \dots z_{\varepsilon_{t_s}, i_{t_s}}]$$

and are trigraded:  $s$  is the *homological degree*,  $t = t_s$  is the *length* and  $d = 2(p - 1) \sum_{k=1}^{t_s} i_k + \sum_{k=1}^{t_s} \varepsilon_k$  is the *total degree*, which we usually disregard in notations. Let  $\overline{B}_s(\mathcal{Q})_t$  be the submodule generated by elements of bidegree  $(s, t)$ . Given a generator  $z = [z_{I_1} | \dots | z_{I_s}]$  of  $\overline{B}_s(\mathcal{Q})$ , for any  $j = 1, \dots, s - 1$ , let

$$\partial_{s,j} : \overline{B}_s(\mathcal{Q}) \rightarrow \overline{B}_{s-1}(\mathcal{Q})$$

be the map defined by

$$\partial_{s,j}(z) = [z_{I_1} | \dots | z_{I_j} z_{I_{j+1}} | \dots | z_{I_s}].$$

Then we consider the following differential  $\partial$  for  $\overline{B}(\mathcal{Q})$ :

$$\partial_s : \overline{B}_s(\mathcal{Q}) \rightarrow \overline{B}_{s-1}(\mathcal{Q})$$

defined by

$$\partial_s(z) = \sum_{j=1}^{s-1} (-1)^{e_{I_j}} \partial_{s,j}(z),$$

where  $e_{I_j} = j + \sum_{k=1}^j |z_{I_k}|$ , being  $|z_{I_k}|$  the total degree of  $z_{I_k}$ . The chain complex  $(\overline{B}(\mathcal{Q}), \partial)$ , known as the *reduced bar construction*, computes the homology of  $\mathcal{Q}$ ,  $H_{s,t}(\mathcal{Q}) = \text{Tor}_{s,t}^{\mathcal{Q}}(\mathbb{F}_p, \mathbb{F}_p)$ . We know by [4] that  $\text{Tor}_{s,t}^{\mathcal{Q}}(\mathbb{F}_p, \mathbb{F}_p) = 0$  when  $s \neq t$ , so we are only interested on the diagonal part of the homology:

$$D_*(\mathcal{Q}) = \bigoplus_{k \geq 0} D_k(\mathcal{Q}) = \bigoplus_{k \geq 0} H_{k,k}(\mathcal{Q}).$$

The group  $D_k(\mathcal{Q})$  turns out simply to be  $\ker(\partial_k) : \overline{B}_k(\mathcal{Q})_k \rightarrow \overline{B}_{k-1}(\mathcal{Q})_k$ , since there exist no non-zero  $(k + 1)$ -chains of length  $k$ . The following Theorem helps to identify the elements

$$z = \sum_I f_I [z_{\epsilon_1, i_1} | \cdots | z_{\epsilon_k, i_k}] \tag{3.3}$$

in  $\overline{B}_k(Q)_k$  belonging to  $D_k(Q) = \ker(\partial_k)$ . Note that only a finite number of  $\mathbb{F}_p$ -coefficients  $f_I$  in (3.3) is non-zero.

**Theorem 3.2** *The element  $z = \sum_I f_I [z_{\epsilon_1, i_1} | \cdots | z_{\epsilon_k, i_k}] \in \overline{B}_k(Q)_k$  is a cycle if and only if for each  $j (1 \leq j \leq k - 1)$  and each  $((\epsilon_1, s_1), \dots, (\epsilon_{j-1}, s_{j-1})) \in (\{0, 1\} \times \mathbb{Z})^{j-1}, ((\epsilon_{j+2}, s_{j+2}), \dots, (\epsilon_k, s_k)) \in (\{0, 1\} \times \mathbb{Z})^{k-j-1}$ , the following condition holds:*

$$\sum_I f_I z_{e_j, i_j} z_{e_{j+1}, i_{j+1}} = 0,$$

where the summation runs over all  $I$ 's such that

$$((e_1, i_1), \dots, (e_{j-1}, i_{j-1})) = ((\epsilon_1, s_1), \dots, (\epsilon_{j-1}, s_{j-1}))$$

and

$$((e_{j+2}, i_{j+2}), \dots, (e_k, i_k)) = ((\epsilon_{j+2}, s_{j+2}), \dots, (\epsilon_k, s_k)).$$

*Proof* One can follow the same argument used to prove Theorem 1 in [8]. □

As a consequence of this result we have the following Corollary. It can be proved by an argument similar to that for  $p = 2$  in Corollary 2 of [8].

**Corollary 3.3** *Suppose that  $z = \sum_I f_I [z_{\epsilon_1, i_1} | \cdots | z_{\epsilon_k, i_k}] \in \overline{B}_k(Q)_k$  is a cycle. For each  $S = ((e_1, s_1), \dots, (e_q, s_q)) \in (\{0, 1\} \times \mathbb{Z})^q$  and  $S' = ((e_{q+1}, s_{q+1}), \dots, (e_k, s_k)) \in (\{0, 1\} \times \mathbb{Z})^{k-q}$ , let  $z_S$  equal to*

$$\sum_I f_I [z_{\epsilon_{q+1}, i_{q+1}} | \cdots | z_{\epsilon_k, i_k}],$$

where the summation runs over the labels  $I$  such that

$$(\epsilon_1, i_1) = (e_1, s_1), \dots, (\epsilon_q, i_q) = (e_q, s_q),$$

and  $z_{S'}$  equal to

$$\sum_I f_I [z_{\epsilon_1, i_1} | \cdots | z_{\epsilon_q, i_q}],$$

where the summation runs over the labels  $I$  such that

$$(\epsilon_{q+1}, i_{q+1}) = (e_{q+1}, s_{q+1}), \dots, (\epsilon_k, i_k) = (e_k, s_k).$$

Then  $z_S$  is a cycle of  $\overline{B}_{k-q}(Q)_{k-q}$  and  $z_{S'}$  is a cycle of  $\overline{B}_q(Q)_q$ .

**Proposition 3.4** *Set*

$$R(k, n, \varepsilon) = [z_{\varepsilon, pk-1-n} | z_{0,k}] - \sum_j A_{(n,j)} [z_{\varepsilon, pk-1-j} | z_{0, k-n+j}]$$

and

$$S(k, n, \varepsilon) = [z_{1-\varepsilon, pk-n} | z_{1,k}] - \sum_j A_{(n,j)} [z_{1-\varepsilon, pk-j} | z_{1, k-n+j}] + \varepsilon \sum_j B_{(n,j)} [z_{1, pk-j} | z_{0, k-n+j}].$$

Then  $D_2(\mathcal{Q})$  has  $\{R(k, n, \varepsilon), S(k, n, \varepsilon)\}_{k \in \mathbb{Z}, n \in \mathbb{N}_0, \varepsilon \in \{0,1\}}$  as a linear  $\mathbb{F}_p$ -basis.

*Proof* To see this, observe that

$$\partial_2([z_{\varepsilon_1, i_1} | z_{\varepsilon_2, i_2}]) = (-1)^{1+\varepsilon_1+2i_1(p-1)} [z_{\varepsilon_1, i_1} z_{\varepsilon_2, i_2}] = (-1)^{1+\varepsilon_1} [z_{\varepsilon_1, i_1} z_{\varepsilon_2, i_2}].$$

Then  $\partial_2(R(k, n, \varepsilon))$  and  $\partial_2(S(k, n, \varepsilon))$  vanish since they correspond to the generating relations (3.1) and (3.2) of  $\mathcal{Q}$ . □

Now we give some examples of cycles constructed by iterating the following generalized Adem relations:

$$z_{0, pk-1} z_{0,k} = 0, \quad z_{1, pk} z_{1,k} = 0, \quad z_{1, pk-1} z_{0,k} = 0.$$

They are

$$z = [z_{0, p^{m-1}k - \frac{p^{m-1}-1}{p-1}} | z_{0, p^{m-2}k - \frac{p^{m-2}-1}{p-1}} | \cdots | z_{0, pk-1} | z_{0,k}]$$

and

$$z' = [z_{1, p^{m-1}k} | [z_{1, p^{m-2}k} | \cdots | z_{1, pk} | z_{1,k}],$$

both elements of  $D_m(\mathcal{Q})$ .

Further, for any  $1 \leq j < m$ , we get another element of  $D_m(\mathcal{Q})$  given by the following chain:

$$z''_j = [z_{1, \alpha_{m-1}} | \cdots | z_{1, \alpha_{j+1}} | z_{1, \alpha_j} | z_{0, \alpha_{j-1}} | \cdots | z_{0, \alpha_1} | z_{0, \alpha_0}],$$

where

$$\alpha_t = \begin{cases} p^t k - \frac{p^t-1}{p-1} & \text{if } 0 \leq t \leq j-1 \\ p^t k - p^{t-j+1} \frac{p^{j-1}-1}{p-1} & \text{if } j \leq t \leq m-1 \end{cases} \tag{3.4}$$

Then, using the Adem relation  $z_{0, pk} z_{1,k} = z_{1, pk} z_{0,k}$  in addition to the others above, we get the following cycle of  $D_3(\mathcal{Q})$ :

$$z_3 = [z_{0,p^2k}|z_{1,pk}|z_{1,k}] + [z_{1,p^2k}|z_{0,pk}|z_{1,k}] + [z_{1,p^2k}|z_{1,pk}|z_{0,k}]$$

and two examples of cycles in  $D_4(\mathcal{Q})$ :

$$\begin{aligned} z_4 &= [z_{\epsilon,p^3k-1}|z_{0,p^2k}|z_{1,pk}|z_{1,k}] + [z_{1,p^3k}|z_{1,p^2k}|z_{0,pk}|z_{1,k}] \\ &\quad + [z_{1,p^3k}|z_{1,p^2k}|z_{1,pk}|z_{0,k}], \\ z'_4 &= [z_{1,p^3k}|z_{0,p^2k}|z_{1,pk}|z_{1,k}] + [z_{1,p^3k}|z_{1,p^2k}|z_{0,pk}|z_{1,k}] \\ &\quad + [z_{1,p^3k}|z_{1,p^2k}|z_{1,pk}|z_{0,k}] + [z_{0,p^3k}|z_{1,p^2k}|z_{1,pk}|z_{1,k}]. \end{aligned}$$

**Theorem 3.5** *The diagonal homology  $D_*(\mathcal{Q})$  has a coalgebra structure given by*

$$\begin{aligned} \psi_n : D_n(\mathcal{Q}) &\rightarrow \bigoplus_{q+t=n} (D_q(\mathcal{Q}) \otimes D_t(\mathcal{Q})), \\ z &= \sum_I f_I [z_{\epsilon_1,i_1}|z_{\epsilon_2,i_2}|\dots|z_{\epsilon_n,i_n}] \mapsto z \otimes 1 + 1 \otimes z + \sum z' \otimes z'', \end{aligned}$$

where the cycles  $z'$  and  $z''$  are obtained by splitting all the summands of  $z$  and suitably grouping the common terms.

*Proof* According to Corollary 3.3, the elements  $z'$  and  $z''$ , coming from the procedure described in the statement above, are cycles. □

### 4 The isomorphism between $D_*(\mathcal{Q})$ and $\Gamma$

We want to show that the diagonal homology of  $\mathcal{Q}$  is isomorphic to  $\Gamma$ . To this purpose, let us consider the  $\mathbb{F}_p$ -linear maps  $\pi_{n,q} : \overline{\Delta}_n \rightarrow \overline{B}_{n-1}(\mathcal{Q})_n$ , for  $n \geq 2$  and  $q = 1, \dots, n - 1$ , defined as follows: given  $\overline{u}^{\mathcal{E}} w^I = \overline{u}_1^{\epsilon_1} \dots \overline{u}_n^{\epsilon_n} w_1^{i_1} \dots w_n^{i_n} \in \overline{\Delta}_n$ ,

$$\pi_{n,q}(\overline{u}^{\mathcal{E}} w^I) = [z_{1-\epsilon_1,i_1}|\dots|z_{1-\epsilon_{q-1},i_{q-1}}|z_{1-\epsilon_q,i_q}z_{1-\epsilon_{q+1},i_{q+1}}|\dots|z_{1-\epsilon_n,i_n}].$$

We begin by looking at the map  $\pi_{2,1}$ .

**Proposition 4.1** *ker  $\pi_{2,1} = \Gamma_2$ .*

*Proof* The map  $\pi_{2,1} : \overline{\Delta}_2 \rightarrow \overline{B}_1(\mathcal{Q})_2$  acts as follows:

$$\pi_{2,1}(\overline{u}_1^{\epsilon_1} \overline{u}_2^{\epsilon_2} w_1^{i_1} w_2^{i_2}) = [z_{1-\epsilon_1,i_1}z_{1-\epsilon_2,i_2}].$$

Using relations (2.3) and the linearity of  $\pi_{2,1}$ , we get

$$\begin{aligned} \pi_{2,1}(\mathcal{Q}_{2,0}^k \mathcal{Q}_{2,1}^s) &= \pi_{2,1}(w_1^{pk+(p-1)s} w_2^k (w_1 + w_2)^s) \\ &= \pi_{2,1} \left( \sum_{j=0}^s \binom{s}{j} w_1^{p(k+s)-s+j} w_2^{k+s-j} \right) \end{aligned}$$



$$\begin{aligned}
 &= \sum_{j=0}^s \binom{s}{j} \pi_{2,1} \left( w_1^{p(k+s)-s+j} w_2^{k+s-j} \right) \\
 &= \sum_{j=0}^s \binom{s}{j} [z_{1,p(k+s)-(s-j)} z_{1,k+s-j}] \\
 &= [d^s(z_{1,p(k+s)} z_{1,k+s})],
 \end{aligned}$$

and  $d^s(z_{1,p(k+s)} z_{1,k+s}) = 0$  according to Proposition 3.1. In a similar way, one can also prove that

$$\begin{aligned}
 \pi_{2,1}(R_{2,0} R_{2,1} Q_{2,0}^k Q_{2,1}^s) &= -[d^s(z_{0,p(k+s+2)-1} z_{0,k+s+2})] = 0, \\
 \pi_{2,1}(R_{2,1} Q_{2,0}^k Q_{2,1}^s) &= [d^s(z_{1,p(k+s+1)-1} z_{0,k+s+1})] = 0, \\
 \pi_{2,1}(R_{2,0} Q_{2,0}^k Q_{2,1}^s) &= [d^s(z_{1,p(k+s+1)} z_{0,k+s+1} - z_{0,p(k+s+1)} z_{1,k+s+1})] = 0,
 \end{aligned}$$

that is the elements of  $\Gamma_2 \subset \overline{\Delta}_2$  correspond to the defining relations of  $\mathcal{Q}$  in terms of  $d$ . Hence  $\Gamma_2$  is the kernel of  $\pi_{2,1}$ . □

**Lemma 4.2** For any  $n \geq 2$  and  $q = 1, \dots, n - 1$ :

$$ker \pi_{n,q} = \overline{\Delta}_{q-1} \otimes \Gamma_2 \otimes \overline{\Delta}_{n-q-1}.$$

*Proof* For any  $t \in \mathbb{N}$ , we define  $\pi_t : \overline{\Delta}_t \rightarrow \overline{B}_t(\mathcal{Q})_t$  as

$$\pi_t(\overline{u}_1^{\varepsilon_1} \dots \overline{u}_t^{\varepsilon_t} w_1^{i_1} \dots w_t^{i_t}) = [z_{1-\varepsilon_1, i_1} | \dots | z_{1-\varepsilon_t, i_t}].$$

We write  $f$  for the composition  $(\psi_{q-1,2} \otimes 1) \circ \psi_{q+1,n-q-1}$ :

$$f : \overline{\Delta}_n \rightarrow \overline{\Delta}_{q-1} \otimes \overline{\Delta}_2 \otimes \overline{\Delta}_{n-q-1}.$$

Then

$$\pi_{n,q} = (\pi_{q-1} \otimes \pi_{2,1} \otimes \pi_{n-q-1}) \circ f.$$

Our result follows from Proposition 4.1 and the fact that  $f, \pi_{q-1}$  and  $\pi_{n-q-1}$  are  $\mathbb{F}_p$ -linear isomorphisms. □

**Lemma 4.3** The general linear group  $GL_n(\mathbb{F}_p)$  is generated by all matrices of the form

$$M = \begin{pmatrix} I_{q-1} & O & O \\ O & A & O \\ O & O & I_{n-q-1} \end{pmatrix},$$

where  $A \in GL_2(\mathbb{F}_p)$ .

*Proof* This result follows from the fact that every invertible matrix admits an *elementary bidiagonal factorization* (see [10]). □

According to the previous Lemma,

$$\Gamma_n = \cap_{q=1}^{n-1} \overline{\Delta}_{q-1} \otimes \overline{\Gamma}_2 \otimes \overline{\Delta}_{n-q-1}.$$

Combining with Lemma 4.2, we arrive at

$$\Gamma_n = \cap_{q=1}^{n-1} \ker \pi_{n,q}. \tag{4.1}$$

We write  $h_n$  for the  $\mathbb{F}_p$ -linear isomorphism inverse to  $\pi_n$ ,

$$h_n = \pi_n^{-1} : \overline{B}_n(\mathcal{Q})_n \rightarrow \overline{\Delta}_n, \quad h_n(|z_{\varepsilon_1, i_1} \cdots |z_{\varepsilon_n, i_n}) = \overline{u}_1^{1-\varepsilon_1} \cdots \overline{u}_n^{1-\varepsilon_n} w_1^{i_1} \cdots w_n^{i_n}.$$

We observe that, for any  $q = 1, \dots, n - 1$ ,  $\partial_{n,q} : \overline{B}_n(\mathcal{Q})_n \rightarrow \overline{B}_{n-1}(\mathcal{Q})_n$  is the result of the composition  $\pi_{n,q} \circ h_n$ . We are going to use this fact in the proof of our main result.

**Theorem 4.4**  $\Gamma$  and  $D_*(\mathcal{Q})$  are isomorphic as coalgebras.

*Proof* The maps  $\{h_n\}_{n \in \mathbb{N}}$  establish a map of coalgebras

$$h : \oplus_{n \in \mathbb{N}} \overline{B}_n(\mathcal{Q})_n \rightarrow \overline{\Delta}.$$

A chain  $z \in \overline{B}_n(\mathcal{Q})_n$  represents a cycle if and only if  $\partial_{n,q}(z) = (\pi_{n,q} \circ h_n)(z) = 0$  for any  $q = 1, \dots, n - 1$ . This holds if and only if  $h_n(z) \in \cap_{q=1}^{n-1} \ker \pi_{n,q}$ , that is  $h_n(z) \in \Gamma_n$  according to (4.1). Then  $h_n$  restricts to an isomorphism of coalgebras

$$\overline{h}_n : D_n(\mathcal{Q}) \rightarrow \Gamma_n.$$

□

**References**

1. Bousfield, A.K., Curtis, E.B., Kan, D.M., Quillen, D.G., Rector, D.L., Schlesinger, J.W.: The mod  $p$  lower central series and the Adams spectral sequence. *Topology* **5**, 331–342 (1966)
2. Brunetti, M., Ciampella, A., Lomonaco, L.A.: The cohomology of the universal Steenrod algebra. *Manuscr. Math.* **118**, 271–282 (2005)
3. Brunetti, M., Ciampella, A., Lomonaco, L.A.: An embedding for the  $E_2$ -term of the Adams spectral sequence at odd primes. *Acta Math. Sin. Engl. Ser.* **22**(6), 1657–1666 (2006)
4. Brunetti, M., Ciampella, A.: A Priddy-type Koszulness criterion for non-locally finite algebras. *Colloq. Math.* **109**(2), 179–192 (2007)
5. Brunetti, M., Ciampella, A., Lomonaco, L.A.: Homology and cohomology operations in terms of differential operators. *Bull. Lond. Math. Soc.* **42**, 53–63 (2010)
6. Chataur, D., Livernet, M.: Adem-Cartan operads. *Commun. Algebra* **33**, 4337–4360 (2005)
7. Ciampella, A., Lomonaco, L.A.: The universal Steenrod algebra at odd primes. *Commun. Algebra* **32**(7), 2589–2607 (2004)

8. Ciampella, A., Lomonaco, L.A.: Homological computations in the universal Steenrod algebra. *Fund. Math.* **183**(3), 245–252 (2004)
9. Hung, N.H.V., Sum, H.: On singer's invariant-theoretic description of the lambda algebra: a mod  $p$  analogue. *J. Pure Appl. Algebra* **99**, 297–329 (1995)
10. Johnson, Charles R., Olesky, D.D., van den Driessche, P.: Elementary bidiagonal factorizations. *Linear Algebra Appl.* **292**, 233–244 (1999)
11. Li, H.H., Singer, W.M.: Resolutions of modules over the Steenrod algebra and the classical theory of invariants. *Mathematische Zeitschrift* **81**, 268–286 (1982)
12. Lomonaco, L.A.: The diagonal cohomology of the universal Steenrod algebra. *J. Pure Appl. Algebra* **121**, 315–323 (1997)
13. Mandell, M.A.:  $E_\infty$  algebras and  $p$ -adic homotopy theory. *Topology* **40**, 43–94 (2001)
14. May, J.P.: *A General Approach to Steenrod Operations*, Lecture Notes in Mathematics. Springer, Berlin (1970)
15. Steenrod, N.E.: *Cohomology Operations*, Lectures Written and Revised by D. B. A. Epstein, *Ann. of Math. Studies* 50. Princeton Univ. Press, Princeton (1962)