

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Medicine

journal homepage: <http://ees.elsevier.com/apjtm>Original research <http://dx.doi.org/10.1016/j.apjtm.2016.06.015>Unlocking the *in vitro* anti-*Trypanosoma cruzi* activity of halophyte plants from the southern PortugalMarta Oliveira¹, Policarpo Ademar Sales Junior², Maria João Rodrigues¹, Marina DellaGreca³, Luísa Barreira¹,
Silvane Maria Fonseca Murta², Alvaro José Romanha², Luísa Custódio^{1*}¹Center of Marine Sciences (CCMAR), University of Algarve, Campus de Gambelas, 8005-139, Faro, Portugal²Centro de Pesquisas René Rachou, FIOCRUZ, Belo Horizonte, Minas Gerais, Brazil³Dipartimento di Chimica Organica e Biochimica, Università Federico II, Complesso Universitario Monte Sant'Angelo, Via Cintia 4, I-80126, Napoli, Italy

ARTICLE INFO

Article history:

Received 15 May 2016

Received in revised form 16 Jun 2016

Accepted 21 Jun 2016

Available online 29 Jun 2016

Keywords:

Chagas disease

Halophytes

Trypanosoma cruzi

Phenanthrenes

Juncunol

ABSTRACT

Objective: To evaluate the *in vitro* anti-*Trypanosoma cruzi* (*T. cruzi*) activity of organic extracts prepared from halophyte species collected in the southern coast of Portugal (Algarve), and chemically characterize the most active samples.**Methods:** Acetone, dichloromethane and methanol extracts were prepared from 31 halophyte species and tested *in vitro* against trypomastigotes and intracellular amastigotes of the Tulahuen strain of *T. cruzi*. The most active extract was fractionated by preparative HPLC-DAD, affording 11 fractions. The most selective fraction was fully characterized by ¹H NMR.**Results:** From 94 samples tested, one was active, namely the root dichloromethane extract of *Juncus acutus* (IC₅₀ < 20 µg/mL). This extract was fractionated by HPLC, affording 11 fractions, one of them containing only a pure compound (juncunol), and tested for anti-parasitic activity. Fraction **8** (IC₅₀ = 4.1 µg/mL) was the most active, and was further characterized by ¹H NMR. The major compounds were phenanthrenes, 9,10-dihydrophenanthrenes and benzocoumarins.**Conclusion:** Our results suggest that the compounds identified in fraction **8** are likely responsible for the observed anti parasitic activity. Further research is in progress aiming to isolate and identify the specific active molecules. To the best of our knowledge, this is the first report on the *in vitro* anti *T. cruzi* activity of halophyte species.

1. Introduction

Chagas disease (CD) is a neglected tropical disease (NTD) caused by the protozoan parasite *Trypanosoma cruzi* (*T. cruzi*), transmitted to humans and animals from the faeces of triatomine bugs (kissing bugs). It is estimated that 20–30% of humans infected with *T. cruzi* suffer with severe cardiopathy or megasophagus – megacolon [1]. About 8 million people are probably infected worldwide, especially in Latin America,

where CD is a significant health and socioeconomic problem [2]. Moreover, CD is becoming increasingly widespread in the southern area of the United States, overlapping with the poorest states [3,4]. Recently, non-vectorial *T. cruzi* infection has been increasingly recognized outside endemic areas. Europe, the United States, Canada, Australia, New Zealand and Japan host millions of at-risk Latin American immigrants [5]. Therefore, the potential of CD becoming a public health issue in that area is considerably high, mainly due to the high number of Latin American immigrants and international travellers, which may contribute for indirect transmission such as blood transfusion, organ transplantation and congenital route; and the presence of potential vectors, triatominae, in this region [5,6].

*Corresponding author: Luísa Custódio, Center of Marine Sciences, University of Algarve, Faculty of Sciences and Technology, Ed. 7, Campus de Gambelas, Faro, Portugal.

Tel: +351 289 800900x7381

Fax: +351 289800051

E-mail: lcustodio@ualg.pt

Peer review under the responsibility of Hainan Medical College.

The available drugs for CD treatment are the 5-nitrofurantoin, nitrofurantoin and the 2-nitroimidazole, benznidazole (BZ). Both drugs present low percentage cure rate, mainly in the chronic phase of the disease, when the majority of cases are diagnosed [7]. Moreover, available chemotherapeutics are highly toxic, with severe systemic side effects [7]. The problems associated with the available drugs highlight the urgent need to develop new strategies for chemotherapy against Chagas disease.

Nature has provided an important number of compounds with anti-parasitic activity. For example, quinine, first isolated from the bark of the cinchona tree (*Cinchona* sp.) was the first effective Western treatment for malaria caused by *Plasmodium falciparum*, while artemisinin from *Artemisia annua* L. is still used in malaria treatment in artemisinin-combination therapies. Noteworthy is the fact that the Nobel Prize of Physiology and Medicine 2015 was awarded to three scientists for their discoveries and development of effective drugs against parasitic infections, namely avermectin isolated from *Streptomyces avermitilis* (and its derivative ivermectin) and artemisinin from *A. annua* L. (Asteraceae) [8]. This award emphasizes that nature present unlimited chemical diversity, and highlights the value of natural products as promising alternative therapeutics towards NTDs.

Halophytes are specialized plants able to survive and thrive in saline soils. Although representing only 2% of terrestrial plant species, they are present in about half the higher plant families and have a high diversity of forms. Halophytes have evolved a complex suite of adaptations in response to the osmotic and ionic deficits of saline environments that contribute to the generation of reactive oxygen species (ROS). In order to manage with excessive toxic ROS, halophytes contain antioxidant systems, including enzymes and bioactive compounds, which give them a significant plethora of other biological activities.

Although there are reports of the traditional use of different halophytic species as anti-parasitic and/or anti-helminthic agents [9], to the best of our knowledge there is no scientific information regarding the potential use of halophytes against NTDs in general, or against CD in particular. Therefore this work evaluated organic extracts made from 31 species of halophytes *in vitro* against trypomastigotes and intracellular amastigotes forms from Tulahuen strain of *T. cruzi*. The most active extract was submitted to a bio-guided fractionation, and the most promising fraction was chemically characterized by ¹H-NMR.

2. Material and methods

2.1. Chemicals

All chemicals used in the experiments were of analytical grade, and were purchased from VWR International (Leuven, Belgium).

2.2. Sample collection

A total of 31 indigenous (Table 1), mostly obligate, halophyte species were collected from different saline habitats of

the southern Portugal (Algarve) at their full flowering time during June of 2013. The researched halophytes belong to 16 plant families and include Aizoaceae (*Mesembryanthemum crystallinum* L. and *Carpobrotus edulis* L.), Amaranthaceae (*Arthrocnemum macrostachyum* L., *Halopeplis amplexicaulis* (Vahl) Ung.-Sternb. ex Ces., Pass. & Gibelli, *Salicornia ramosissima* J. Woods, *Salicornia fragilis* P.W. Ball & Tutin, *Salsola vermiculata* L., *Sarcocornia perennis* (Mill.) A.J. Scott subsp. *alpini* (Lag.) Castrov. and *S. perennis* (Mill.) subsp. *perennis*), Anacardiaceae (*Pistacia lentiscus* L.), Asteraceae (*Aster tripolium* L. and *Inula crithmoides* L.), Caryophyllaceae (*Spergularia rubra* (L.) J. Presl & C. Presl), Convolvulaceae [*Calystegia (Convolvulus) soldanella* (L.) R. Br.], Cyperaceae (*Claudium mariscus* (L.) Pohl), Frankeniaceae (*Frankenia pulverulenta* L. and *Frankenia laevis* L.), Gentianaceae (*Centaurium erythraea* Rafn), Juncaceae [*Juncus acutus* (J. acutus) L., *Juncus inflexus* L. and *Juncus maritimus* Lam.], Lythraceae (*Lythrum salicaria* L.), Plumbaginaceae (*Limonium monoptalum* (L.) Boiss., *Limonium algarvense* Erben and *Limonium lanceolatum* Hoffmanns. & Link), Polygonaceae (*Polygonum maritimum* L.) Poaceae (*Panicum repens* L., *Puccinellia maritima* (Huds.) Parl., *Spartina versicolor* Fabre), Tamaricaceae (*Tamarix africana* Poir) and Typhaceae (*Typha domingensis* Pers).

The taxonomical classification was determined by the botanist Dr Manuel J. Pinto (National Museum of Natural History, University of Lisbon, Botanical Garden, Portugal) and voucher specimens are kept in the herbarium of the MarBiotech laboratory (MBH01–MBH31). Different organs were collected for each species, whenever possible (Table 1). Plant material was oven dried for 3 days at 40 °C, powdered and stored at –20 °C until needed.

2.3. Preparation of the extracts

Dried samples were mixed with 80% aqueous acetone, dichloromethane and methanol (1:10, w/v) (Table 1), and extracted overnight at room temperature (RT), under stirring. Extracts were filtered (Whatman n° 4) and concentrated under reduced pressure and temperature (<40 °C). Dried extracts were dissolved in dimethyl sulfoxide (DMSO) and stored at 4 °C at the concentration of 25 mg/mL until analysis.

2.4. Evaluation of *in vitro* antitrypanosomal activity, cellular toxicity and selectivity

The *in vitro* antitrypanosomal activity was evaluated on L929 cells (mouse fibroblasts) infected with the Tulahuen strain of the parasite expressing the *Escherichia coli* β-galactosidase as reporter gene, according to the method described previously [10]. The extracts were tested at the concentration of 20 µg/mL, for a period of incubation of 96 h. Fractions obtained from the active extract were tested at concentrations ranging from 10 µg/mL to 100 µg/mL, also during a 96 h period. Controls with uninfected cells, untreated infected cells, infected cells treated with benznidazole at the concentration of 1 µg/mL (3.8 µM, positive control) or DMSO (1%, v/v) were used. The results were expressed as the percentage of *T. cruzi* growth inhibition

Table 1

Botanical names, families, plant parts and extracts used of the 31 halophytic species included in this study.

Family	Botanical name	Plant part tested	Extracts tested
Aizoaceae	<i>Mesembryanthemum crystallinum</i> L.	Aerial organs	80% acetone, dichloromethane
	<i>Carpobrotus edulis</i> L.	Leaves	80% acetone, dichloromethane
Amaranthaceae	<i>Artrocnemum macrostachyum</i> L.	Aerial organs	80% acetone, dichloromethane
	<i>Halopeplis amplexicaulis</i> (Vahl) Ung.-Sternb. ex Ces., Pass. & Gibelli	Aerial organs	80% acetone, dichloromethane
	<i>Salicornia ramosissima</i> J. Woods	Aerial organs	80% acetone, dichloromethane
	<i>Salicornia fragilis</i> P.W.Ball & Tutin	Aerial organs	80% acetone, dichloromethane
	<i>Salsola vermiculata</i> L.	Aerial organs	80% acetone, dichloromethane
	<i>Sarcocornia perennis</i> (Mill.) A.J. Scott subsp <i>alpini</i> (Lag.) Castrov.	Aerial organs	80% acetone, dichloromethane
	<i>Sarcocornia perennis</i> (Mill.) A.J. Scott subsp <i>perennis</i>	Aerial organs	80% acetone, dichloromethane
Anacardiaceae	<i>Pistacia lentiscus</i> L.	Leaves	80% acetone, dichloromethane
Asteraceae	<i>Aster tripolium</i> L.	Aerial organs	80% acetone, dichloromethane
	<i>Inula crithmoides</i> L.	Aerial organs	80% acetone, dichloromethane
Caryophyllaceae	<i>Spergularia rubra</i> (L.) J. Presl & C. Presl	Whole plant	80% acetone, dichloromethane
Convolvulaceae	<i>Calystegia (Convolvulus) soldanella</i> (L.) R.Br.	Aerial organs	80% acetone, dichloromethane
Cyperaceae	<i>Cladium mariscus</i> (L.) Pohl	Aerial organs	80% acetone, dichloromethane
Frankeniaceae	<i>Frankenia pulverulenta</i> L.	Whole plant	80% acetone, dichloromethane
	<i>Frankenia laevis</i> L.	Whole plant	80% acetone, methanol, dichloromethane
Gentianaceae	<i>Centaurium erythraea</i> Rafn	Whole plant	80% acetone, dichloromethane
Juncaceae	<i>Juncus acutus</i> L.	Leaves, roots, seeds	methanol, dichloromethane
	<i>Juncus inflexus</i> L.	Leaves, roots	methanol, dichloromethane
	<i>Juncus maritimus</i> Lam.	Leaves, roots, seeds	methanol, dichloromethane
Lythraceae	<i>Lythrum salicaria</i> L.	Aerial organs	80% acetone, dichloromethane
Plumbaginaceae	<i>Limoniastrum monopetalum</i> (L.) Boiss.	Aerial organs	80% acetone, dichloromethane
	<i>Limonium algarvense</i> Erben	Leaves, flowers, flower stalks, roots	methanol, dichloromethane
	<i>Limonium lanceolatum</i> Hoffmanns. & Link	Leaves	80% acetone, dichloromethane
Polygonaceae	<i>Polygonum maritimum</i> L.	Leaves, roots	methanol, dichloromethane
Poaceae	<i>Panicum repens</i> L.	Whole plant	80% acetone, dichloromethane
	<i>Pucinellia maritima</i> (Huds.) Parl.	Whole plant	80% acetone, dichloromethane
	<i>Spartina versicolor</i> Fabre	Whole plant	80% acetone, dichloromethane
Tamaricaceae	<i>Tamarix africana</i> Poir.	Leaves	80% acetone, dichloromethane
Typhaceae	<i>Typha domingensis</i> Pers.	Aerial organs	80% acetone, dichloromethane

in extracts-tested infected cells as compared to the untreated infected cells. Active fractions were evaluated for cytotoxicity and selectivity on uninfected fibroblasts [10]. Results obtained with the fractions were expressed as IC₅₀ values, calculated by linear interpolation and the selectivity index (SI) was determined based on the ratio of the IC₅₀ value in the host cell divided by the IC₅₀ value of the parasite.

2.5. Sample fractionation

The active dichloromethane crude extract of *J. acutus* roots was dissolved in dichloromethane at the concentration of 100 mg/mL, and fractionated by preparative HPLC-DAD (Knauer Smartline, Germany) constituted by the following modules: vacuum degasser (E4320V2), quaternary pump (EA4300V1) and the diode array detector (E4350), using a semi-preparative. Analyses were performed on a Luna 5u C18 (2) 100A, (250 × 10) mm, 5 μm particle size (Phenomenex, Spain). The mobile phase consisted on acetonitrile (solvent A) and mili-Q water (solvent B) with the following gradient: 0–40 min: 80–10% A, 40–45 min: 10–0% A, 45–55 min: 0–0% A, 55–60 min: 0–80% A, using a flow of 3.5 mL/min. The injection volume was 200 μL and the detector was set at 216 nm. 11 fractions were collected for re-testing for anti-parasitic activity.

2.6. Spectral analysis

NMR spectra were acquired on a Bruker-600 DRX (¹H NMR: 600 MHz, ¹³C NMR: 150 MHz) spectrometer equipped with a cryo probe, in CDCl₃ (internal standard, for ¹H: CHCl₃ at δ 7.26 ppm; for ¹³C: CDCl₃ at δ 77.0 ppm).

2.7. Statistical analyses

Results were expressed as Mean ± Standard deviation (SD), of at least three replicates. Significant differences were assessed by analysis of variance (ANOVA) or using Kruskal–Wallis test (*P* < 0.05) when parametricity of data did not prevail (SPSS statistical package for Windows, release 15.0). The IC₅₀ values were calculated with GraphPad Prism V 5.0.

3. Results

From 94 samples tested, only was active, namely the dichloromethane extract of *J. acutus* with an IC₅₀ value lower than 20 μg/mL. This extract was fractionated by HPLC affording 11 fractions, one of them containing a pure compound (juncunol) which were tested for cytotoxic activity against *T. cruzi* and mouse fibroblasts (L929 cells), for the determination of selectivity (Table 2). From these, fraction 8 was the most active and selective (IC₅₀ = 4.1 μg/mL, SI = 1.5), and was further characterized chemically.

Although fraction 8 gave rather complex ¹H NMR spectra a qualitative analysis of the overall spectrum was possible (figure 1). The spectrum show typical signals of phenanthrenes

Table 2

Effect of the application of fractions from the dichloromethane extract of *Juncus acutus* (roots), on the viability of trypomastigotes and intracellular amastigotes of Tulahuen strain of *T. cruzi*.

Fraction/Compound	IC ₅₀ values (μg/mL)		Selectivity index
	Parasites	Mammalian cells	
1 Juncunol	67.7	63.5	0.9
2	>100.0	80.0	nd
3	34.3	40.0	1.2
4	>100.0	22.9	nd
5	>100.0	>100.0	nd
6	>100.0	12.8	nd
7	8.4	10.2	1.2
8	4.1	6.0	1.5
9	19.0	19.4	1.0
10	>100.0	80.0	nd
11	>100.0	>100.0	nd
Benznidazole^a	1.0	625.0	625.0

^a Positive control.

(Phe), 9,10-dihydrophenanthrenes (dPhe) and benzocoumarins (Benz), all isolated previously from *J. acutus* (Figures 2 and 3) [11–15]. Peaks have been assigned by comparison with previously published data [10–14]. In the downfield region, between 9.1 and 8.3 ppm, the doublet signals of H-4 proton from phenanthrenes and benzocoumarins are detected. The region between 8.0 and 6.2 ppm shows all the other aromatic signals and H-12 proton from compounds with vinyl chain at C-5. Between 5.9 and 4.7 ppm H-13 protons of vinyl chain (doublets) and carbinol protons of 1-hydroxyethyl (quartet) or hydroxymethylene (singlet) are observed. Lastly, in the upfield region between 3.2 and 0.70 ppm singlet methyls and multiplets of H-9 and H-10 methylenes of 9,10-dihydrophenanthrenes are present. Trying to identify some components of the mixture, a 2D NMR experiments were performed (COSY and HMBC). The analysis of ¹H–¹H COSY evidences, in the downfield region, correlations of doublet at 9.06 δ with signal at 7.08 δ, doublet at 9.02 δ with signal at 7.20 δ, doublet at 8.99 δ with signal at 7.06 δ, and doublet at 8.98 δ with signal at 6.48 δ. The first three spin systems were assigned at phenanthrenes [12,15], while the last was attributed to a benzocoumarin [13]. The presence in the ¹H NMR of a singlet at δ 3.87 (a methoxyl group), correlation observed in COSY (7.63 with 6.73 δ, 6.80 with 5.54 and 5.25 δ), and methyl singlets at 2.30 and 2.24 δ could be attributed at 8-hydroxy-2-methoxy-1,6-dimethyl-5-vinyl-9,10-dihydrophenanthrene [12]. The presence of this metabolite was confirmed by the long-range hetero-correlations observed in the HMBC spectrum, in fact methoxyl and doublet proton (7.63 δ) gave crosspeak with carbon at 156.0 ppm assigned to C-2. Furthermore, a careful analysis of COSY spectrum evidenced a proton 4.15 δ (dd, *J* = 9.4, 7.8 Hz) correlated to 3.19 and 1.91 δ, and 5.81 δ with 4.98 and 4.44 (dd, *J* = 17.8, 1.3 Hz), these two spin system are reliable with a dimeric phenanthrenoid (Figure 3) [11]. Determinant long-range hetero-correlations of both signal at 4.44 and 4.15 ppm with carbon at 62.0 δ, observed in the HMBC spectrum, supported the identification of these metabolites.




Figure 1. ^1H NMR of *J. acutus* active fraction.




Figure 2. Structure of phenanthrene, benzocoumarin and 9,10-dihydrophenanthrene.




Figure 3. Structure of dimeric 9,10-dihydrophenanthrene.

4. Discussion

There are several reports on the antiparasitic activity of halophyte species. For example, seed kernels of *Caesalpinia crista* (Fabaceae) are used in traditional medicine for the

treatment of malaria [16]. That activity was confirmed by *in vivo* studies using mice infected with *Plasmodium berghei* [17] and was attributed to the presence of cassane- and norcassane-type diterpenes [18]. The whole aerial organs and roots of *Inula cappa* (Asteraceae) are also traditionally used for the treatment of malaria [19,20], although the molecules responsible for that activity were not described yet. To the best of our knowledge the antiprotozoal potential of marine halophytes still remains unexplored, especially as possible candidate against CD. In this context, this work evaluated for the first time the anti-*T. cruzi* activity of organic extracts made from 31 halophyte species abundant in the southern area of Portugal.

Only one extract was able to decrease the growth of parasites, namely the dichloromethane extract of *J. acutus* roots. *Juncus* is the largest genus in the Juncaceae family comprising more than 200 species that usually grow in maritime environments. Several *Juncus* species have medicinal properties, e.g., the medulla of *Juncus effusus* is used as antipyretic and sedative in Japan and China [21]. Also, they are used in traditional medicine for the

treatment of different health problems. For example, the rhizomes of *J. acutus* are used for insomnia and the seed of *Juncus* species for the treatment of stomach disorders [21].

To assess which extract components could be responsible for the antiprotozoal activity, the active extract from *J. acutus* was fractionated by HPLC, affording 11 fractions, one of them containing a pure compound (juncunol), which were also evaluated for antitrypanosomal activity. When applied at the concentration of 67.7 µg/mL juncunol was able to reduce the parasites growth by 50%. However, it was also cytotoxic against mouse fibroblasts L929 cells. Juncunol is a dihydrophenanthrene previously isolated from of different *Juncus* species, including *J. acutus* and *Juncus roemerianus* [15,22–25], and has cytotoxic activity towards the microalga *Selenastrum capricornutum* [15], and for several mammalian tumour cell lines [22]. However, there were no reports until now on its antiparasitic activity.

Fraction 8 had the highest activity towards *T. cruzi*, and contained phenanthrenes, dihydrophenanthrenes and benzocoumarins, which were identified by ¹H NMR spectra analysis. Species belonging to the *Juncus* genus are one of the most prolific sources of phenanthrenes [18], which have several biological activities, including antiproliferative, antioxidant, antimicrobial and cytotoxic [22,26]. Phenanthrenoids and benzocoumarins obtained from the rhizome of *J. acutus* have *in vitro* phytotoxicity, antialgal and anti-inflammatory activities [12,14,15,27]. Recently, phenanthrene and phenanthrenoids, obtained through a bioguided fractionation of the ethanol extract of *Juncus effuses* had *in vitro* cytotoxic properties towards different cancer cell lines [28]. Also, a number of benzocoumarins are described with cytotoxic activity against different cells [29]. Moreover, some phenanthrene-derived molecules and benzocoumarins also have anti-parasitic activity. The best example is halofantrine, a phenanthrene methanol derivative used in the treatment of malaria [30]. However, this molecule is no longer recommended in current therapies due to its cardiotoxicity [30,31]. More recently, other phenanthrene-based derivatives, particularly 3-hydroxy-N'-arylideneopropanehydrazonamides, have shown potent antimalarial *in vitro* activities with high selectivity indexes [32]. Different benzocoumarin scaffolds are highly toxic towards several parasites, including the protozoans *Plasmodium* spp and *Babesia* spp [29]. Since phenanthrene-derived compounds and benzocoumarins have previously demonstrated its efficacy as antiparasitic agents, our results strongly suggest that the molecules identified in the active fraction from *J. acutus* are responsible for its antitrypanosomal activity.

To the best of our knowledge, this is the first report of the potential antitrypanosomal activity of halophyte plants in general, and of the antitrypanosomal activity of *J. acutus* extract and fractions as well as the isolated compound, juncunol. Based on our results, it is likely that the molecules identified in the active fraction from *J. acutus* are responsible for its anti *T. cruzi* activity encouraging further research. In this sense studies aiming the isolation of the bioactive (s) compound (s) of this fraction are already in progress. Moreover, structure–activity relationship (SAR) studies may also disclose a renewed interest in the pharmacological applications of these molecules, by increasing its selectivity towards parasites.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgements

This work was supported by the XtremeBio (PTDC/MAR-EST/4346/2012) and MaNaCruzi projects (bilateral project, FCT/CAPES 2358, 2014/2015) funded by FCT – Foundation for Science and Technology and Portuguese National Budget; it also received national funds through FCT Project CCMAR/Multi/04326/2013 and P3D-Programa de Descoberta e Desenvolvimento de Drogas (PROEP/CNPq/FIOCRUZ 401988/2012-0). The authors thank the Program for Technological Development of Tools for Health-PDTIS-Fiocruz for use of its facilities. AJR and SMFM are CNPq Research Fellows. Luísa Custódio was supported by FCT Investigator Programme (IF/00049/2012) and Policarpo Sales by Programa Brasil Sem Miséria/Coordenação de Aperfeiçoamento de Pessoal de Ensino Superior (CAPES)/FIOCRUZ.

References

- [1] Rassi A Jr, Rassi A, Marin-Neto JA. Chagas disease. *Lancet* 2010; **375**: 1388–1402.
- [2] World Health Organization (WHO). *Working to overcome the global impact of neglected tropical diseases: first WHO report on neglected tropical diseases*. Geneva: World Health Organization; 2010. [Online]. Available from: http://apps.who.int/iris/bitstream/10665/44440/1/9789241564090_eng.pdf [Accessed on 29 February, 2016]
- [3] Hotez PJ. Neglected infections of poverty in the United States of America. *PLoS Negl Trop Dis* 2008; **2**: e256.
- [4] Hotez PJ. Neglected parasitic infections and poverty in the United States. *PLoS Negl Trop Dis* 2014; **8**: e3012.
- [5] Schmunis GA, Yadon ZE. Epidemiology of Chagas disease in non-endemic countries: the role of international migration. *Mem Inst Oswaldo Cruz* 2007; **102**(Suppl 1): 75–85.
- [6] World Health Organization (WHO). *Control and prevention of Chagas disease in Europe. Report of a WHO informal consultation (Jointly Organized by WHO headquarters and the WHO Regional Office for Europe) Geneva, Switzerland, 17–18 December 2009*. Geneva: WHO; 2010.. Report No: WHO/HTM/NTD/IDM/2010.1[Online]. Available from: http://www.fac.org.ar/1/comites/chagas/Chagas_WHO_Technical20Report_16_06_10.pdf.
- [7] Croft SL. Pharmacological approaches to antitrypanosomal chemotherapy. *Mem Inst Oswaldo Cruz* 1999; **94**: 215–220.
- [8] Efferth T, Zacchino S, Georgiev MI, Liu L, Wagner H, Panossian A. Nobel Prize for artemisinin brings phytotherapy into the spotlight. *Phytomedicine* 2015; **22**(13): A1–A3.
- [9] Ksouri R, Ksouri WM, Jallali I, Debez A, Magné C, Hiroko I, et al. Medicinal halophytes: potent source of health promoting biomolecules with medical, nutraceutical and food applications. *Crit Rev Biotechnol* 2012; **32**(4): 289–326.
- [10] Romanha AJ, Castro SL, Soeiro Mde N, Lannes-Vieira J, Ribeiro I, Talvani A, et al. *In vitro* and *in vivo* experimental models for drug screening and development for Chagas disease. *Mem Inst Oswaldo Cruz* 2010; **105**: 233–238.
- [11] DellaGreca M, Fiorentino A, Monaco P, Previtera L, Zarrelli A. A new dimeric 9,10-dihydrophenanthrenoid from the rhizome of *Juncus acutus*. *Tetrahedron Lett* 2002; **43**(14): 2573–2575.
- [12] DellaGreca M, Fiorentino A, Isidori M, Lavorgna M, Monaco P, Previtera L, et al. Phenanthrenoids from the wetland *Juncus acutus*. *Phytochemistry* 2002; **60**(6): 633–638.
- [13] DellaGreca M, Fiorentino A, Isidori M, Previtera L, Temussi F, Zarrelli A. Benzocoumarins from the rhizomes of *Juncus acutus*. *Tetrahedron* 2003; **59**(26): 4821–4825.
- [14] DellaGreca M, Fiorentino A, Monaco P, Previtera L, Temussi F, Zarrelli A. New dimeric phenanthrenoids from the rhizomes of *Juncus acutus*. Structure determination and antialgal activity. *Tetrahedron* 2003; **59**(13): 2317–2324.

- [15] DellaGreca M, Isidori M, Lavorgna M, Monaco P, Previtiera L, Zarrelli A. Bioactivity of phenanthrenes from *Juncus acutus* on *Selenastrum capricornutum*. *J Chem Ecol* 2004; **30**(4): 867-879.
- [16] Ogata Y. *Medicinal herb index in Indonesia*. 2nd ed. Jakarta: P. T. Eisai Indonesia; 1995.
- [17] Banskota AH, Attamimi F, Linn TZ, Usia T, Tezuka Y, Kalauni SK, et al. Novel norcassane-type diterpene from the seed kernels of *Caesalpinia crista*. *Tetrahedron Lett* 2003; **44**: 6879-6882.
- [18] Linn TZ, Awale S, Tezuka Y, Banskota AH, Kalauni SK, Attamimi F, et al. Cassane and norcassane-type diterpenes from *Caesalpinia crista* of Indonesia and their antimalarial activity against the growth of *Plasmodium falciparum*. *J Nat Prod* 2005; **68**: 706-710.
- [19] Wang FY, Li XQ, Sun Q, Yao S, Ke CQ, Tang CP, et al. Sesquiterpene lactones from *Inula cappa*. *Phytochem Lett* 2012; **5**: 639-642.
- [20] Seca AM, Grigore A, Pinto DC, Silva AM. The genus *Inula* and their metabolites: from ethnopharmacological to medicinal uses. *J Ethnopharmacol* 2014; **154**(2): 286-310.
- [21] El-Shamy A, Abdel-Razek AF, Nassar MI. Phytochemical review of *Juncus* L. genus (Fam. Juncaceae). *Arab J Chem* 2015; **8**(5): 614-623.
- [22] Rodrigues MJ, Gangadhar KN, Vizetto-Duarte C, Wubshet SG, Nyberg NT, Barreira L, et al. Maritime halophyte species from Southern Portugal as sources of bioactive molecules. *Mar Drugs* 2014; **12**(4): 2228-2244.
- [23] Sarkar H, Zerezhgi M, Bhattacharyya J. Dehydrojuncosol, a constituent of the roots of *Juncus roemerianus*. *Phytochemistry* 1988; **27**: 3006-3008.
- [24] Della Greca M, Fiorentino A, Mangoni L, Molinaro A, Monaco P, Previtiera L. 9,10-Dihydrophenanthrene metabolites from *Juncus effusus* L. *Tetrahedron Lett* 1992; **33**: 5257-5260.
- [25] Della Greca M, Fiorentino A, Mangoni A, Molinaro A, Monaco P, Previtiera L. A bioactive dihydrodibenzoxepin from *Juncus effusus*. *Phytochemistry* 1993; **34**: 1182-1184.
- [26] Kovács A, Vasas A, Hohmann J. Natural phenanthrenes and their biological activity. *Phytochemistry* 2008; **69**(5): 1084-1110.
- [27] Behery FA, Naeem ZE, Maatooq GT, Amer MM, Wen ZH, Sheu JH, et al. Phenanthrenoids from *Juncus acutus* L., new natural lipopolysaccharide-inducible nitric oxide synthase inhibitors. *Chem Pharm Bull* 2007; **55**: 1264-1266.
- [28] Ma W, Zhang Y, Ding YY, Liu F, Li N. Cytotoxic and anti-inflammatory activities of phenanthrenes from the medullae of *Juncus effusus* L. *Arch Pharma Res* 2016; **39**(2): 154-160.
- [29] Lv HN, Tu PF, Jiang Y. Benzocoumarins: isolation, synthesis, and biological activities. *Mini Rev Med Chem* 2014; **14**(7): 603-622.
- [30] World Health Organization (WHO). *WHO Guidelines for the treatment of malaria*. Geneva: World Health Organization; 2006.. Available from: https://books.google.pt/books?id=Am83eI7_poUC&printsec=frontcover&hl=pt-PT&source=gbs_ge_summary_r&cad=0#v=onepage&q&f=false [Accessed 15 February, 2016]
- [31] World Health Organization (WHO). *WHO Guidelines for the treatment of malaria*. 3rd ed. Geneva: World Health Organization; 2015. Available from: http://apps.who.int/iris/bitstream/10665/162441/1/9789241549127_eng.pdf [Accessed 16 February, 2016]
- [32] Leven M, Held J, Duffy S, Tschan S, Sax S, Kamber J, et al. Blood schizontocidal and gametocytocidal activity of 3-hydroxy-N'-arylidenepropanehydrazonamides: a new class of antiplasmodial compounds. *J Med Chem* 2014; **57**(19): 7971-7976.