


NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all
Aspects of Natural Products Research



Volume 11. Issue 6. Pages 707-882. 2016
ISSN 1934-578X (printed); ISSN 1555-9475 (online)
www.naturalproduct.us

EDITOR-IN-CHIEF**DR. PAWAN K AGRAWAL**

Natural Product Inc.
7963, Anderson Park Lane,
Westerville, Ohio 43081, USA
agrawal@naturalproduct.us

EDITORS**PROFESSOR ALEJANDRO F. BARRERO**

Department of Organic Chemistry, University of Granada,
Campus de Fuente Nueva, s/n, 18071, Granada, Spain
afbarre@ugr.es

PROFESSOR MAURIZIO BRUNO

Department STEBICEF,
University of Palermo, Viale delle Scienze,
Parco d'Orleans II - 90128 Palermo, Italy
maurizio.bruno@unipa.it

PROFESSOR DE-AN GUO

National Engineering Laboratory for TCM Standardization Technology,
Shanghai Institute of Materia Medica, Chinese Academy of Sciences,
Shanghai 201203, P. R. China
gda5958@163.com

PROFESSOR VLADIMIR I. KALININ

G.B. Elyakov Pacific Institute of Bioorganic Chemistry,
Far Eastern Branch, Russian Academy of Sciences,
Pr. 100-letya Vladivostoka 159, 690022,
Vladivostok, Russian Federation
kalininv@piboc.dvo.ru

PROFESSOR YOSHIHIRO MIMAKI

School of Pharmacy,
Tokyo University of Pharmacy and Life Sciences,
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan
mimakiy@ps.toyaku.ac.jp

PROFESSOR STEPHEN G. PYNE

Department of Chemistry, University of Wollongong,
Wollongong, New South Wales, 2522, Australia
spyne@uow.edu.au

PROFESSOR MANFRED G. REINECKE

Department of Chemistry, Texas Christian University,
Forts Worth, TX 76129, USA
m.reinecke@tcu.edu

PROFESSOR WILLIAM N. SETZER

Department of Chemistry, The University of Alabama in Huntsville,
Huntsville, AL 35809, USA
wsetzer@chemistry.uah.edu

PROFESSOR YASUHIRO TEZUKA

Faculty of Pharmaceutical Sciences, Hokuriku University,
Ho-3 Kanagawa-machi, Kanazawa 920-1181, Japan
y-tezuka@hokuriku-u.ac.jp

PROFESSOR DAVID E. THURSTON

Institute of Pharmaceutical Science
Faculty of Life Sciences & Medicine
King's College London, Britannia House
7 Trinity Street, London SE1 1DB, UK
david.thurston@kcl.ac.uk

HONORARY EDITOR**PROFESSOR GERALD BLUNDEN**

The School of Pharmacy & Biomedical Sciences,
University of Portsmouth,
Portsmouth, PO1 2DT U.K.
axuf64@dsl.pipex.com

ADVISORY BOARD

Prof. Viqar Uddin Ahmad
Karachi, Pakistan

Prof. Giovanni Appendino
Novara, Italy

Prof. Yoshinori Asakawa
Tokushima, Japan

Prof. Roberto G. S. Berlink
São Carlos, Brazil

Prof. Anna R. Bilia
Florence, Italy

Prof. Josep Coll
Barcelona, Spain

Prof. Geoffrey Cordell
Chicago, IL, USA

Prof. Fatih Demirci
Eskişehir, Turkey

Prof. Francesco Epifano
Chieti Scalo, Italy

Prof. Ana Cristina Figueiredo
Lisbon, Portugal

Prof. Cristina Gracia-Viguera
Murcia, Spain

Dr. Christopher Gray
Saint John, NB, Canada

Prof. Dominique Guillaume
Reims, France

Prof. Duvvuru Gunasekar
Tirupati, India

Prof. Hisahiro Hagiwara
Niigata, Japan

Prof. Judith Hohmann
Szeged, Hungary

Prof. Tsukasa Iwashina
Tsukuba, Japan

Prof. Leopold Jirovetz
Vienna, Austria

Prof. Phan Van Kiem
Hanoi, Vietnam

Prof. Niel A. Koorbanally
Durban, South Africa

Prof. Chiaki Kuroda
Tokyo, Japan

Prof. Hartmut Laatsch
Gottingen, Germany

Prof. Marie Lacaillle-Dubois
Dijon, France

Prof. Shoen-Sheng Lee
Taipei, Taiwan

Prof. Imre Mathe
Szeged, Hungary

Prof. M. Soledade C. Pedras
Saskatoon, Canada

Prof. Luc Pieters
Antwerp, Belgium

Prof. Peter Proksch
Düsseldorf, Germany

Prof. Phila Raharivelomanana
Tahiti, French Polynesia

Prof. Luca Rastrelli
Fisciano, Italy

Prof. Stefano Serra
Milano, Italy

Dr. Bikram Singh
Palampur, India

Prof. John L. Sorensen
Manitoba, Canada

Prof. Johannes van Staden
Scottsville, South Africa

Prof. Valentin Stonik
Vladivostok, Russia

Prof. Ping-Jyun Sung
Pingtung, Taiwan

Prof. Winston F. Tinto
Barbados, West Indies

Prof. Sylvia Urban
Melbourne, Australia

Prof. Karen Valant-Vetschera
Vienna, Austria

INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

To Subscribe: Natural Product Communications is a journal published monthly. 2016 subscription price: US\$2,595 (Print, ISSN# 1934-578X); US\$2,595 (Web edition, ISSN# 1555-9475); US\$2,995 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

Red Wine Inhibits Aggregation and Increases ATP-diphosphohydrolase (CD39) Activity of Rat Platelets *in Vitro*

Elisabetta Caiazzo¹, Idolo Tedesco², Carmela Spagnuolo², Gian Luigi Russo², Armando Ialenti¹ and Carla Cicala^{1*}

¹Department of Pharmacy, University of Naples "Federico II", Naples, Italy

²Institute of Food Sciences, National Research Council, Avellino, Italy

cicala@unina.it

Received: July 31st, 2015; Accepted: April 13th, 2016

Moderate consumption of red wine has been shown to exert a peculiar cardioprotective effect compared with other alcoholic beverages; inhibition of platelet aggregation seems to be one of the mechanisms underlying this beneficial effect. CD39/ATP-diphosphohydrolase is an integral membrane glycoprotein metabolizing ATP and ADP to AMP; in concert with CD73/ecto-5'-nucleotidase, it contributes to extracellular adenosine accumulation. CD39 is considered a key modulator of thrombus formation; it inhibits platelet aggregation by promoting ADP hydrolysis. There is evidence that red wine consumption increases CD39 activity in platelets from streptozotocin-induced diabetic rats. Here we show that two kinds of Aglianico red wines inhibit aggregation and increase ATP – and ADPase activity in rat platelets.

Keywords: ADP, CD39, Platelets, Rat, Red wine, Thrombosis.

Wine is an alcoholic beverage derived by yeast fermentation of grapes harvested from *Vitis vinifera*. Wine, in particular red wine, has been shown to exert a peculiar cardioprotective effect compared with other alcoholic beverages; inhibition of platelet aggregation seems to be one of the mechanisms underlying this beneficial effect. The antiplatelet effect of red wine has been supposed to be due to a reduced prostanoid synthesis, an increased nitric oxide production and/or an enhanced platelet c-AMP level [1,2]; however, the mechanism is still unclear. CD39 (ATP-diphosphohydrolase) is an integral membrane glycoprotein metabolizing ATP and ADP to AMP; in concert with CD73 (ecto-5'-nucleotidase), it contributes to extracellular adenosine accumulation [3]. CD39 is considered a key modulator of thrombus formation, since it may inhibit platelet aggregation promoting ADP hydrolysis and, in combination with CD73, promotes adenosine accumulation that is a vasodilator and a platelet antiaggregating agent [4-7]. Loss of CD39 activity from the endothelium sustains platelet aggregation and thrombogenesis [8-13]. There is evidence that moderate red wine consumption increases CD39 activity in platelets from streptozotocin-induced diabetic rats [14]; furthermore, flavonoids, which are major wine constituents and well known vasoprotective agents, also increase ATP - and ADPase activity in rat serum [15]. Here, we compared two different red wines, "R" (GAE, gallic acid equivalent, 1.09 mg/mL) and "F" (GAE, 2.19 mg/mL), produced from Aglianico grapes in 2008 and 2011 respectively. Concentrations of different classes of compounds present in the two red wines were investigated and their antioxidant activity was evaluated. The wines presented values that were different, according to the different vintage. As expected, anthocyanin content was higher in the youngest one ("F"; 2011) possibly for its limited concentration of anthocyanin polymers. Both FRAP and DPPH assays showed a direct correlation between the high concentration of phenolic compounds in F 2011 and its antioxidant capacity (Table 1). We found that both wines "R" and "F" inhibited ADP-induced platelet aggregation evaluated in PRP; it is worth noting that while "R" was active only at low ADP concentrations, the effect of "F" was evident even at the highest ADP concentration (Figure 1).

Table 1: Total flavonoids, phenols, anthocyanins and antioxidant properties of Aglianico red wines.

Wine	Total phenol content (mg/mL GAE)	Total flavonoids content (mM QE)	Total anthocyanin content (μM M3GE)	FRAP (mM Fe ²⁺ equivalent)	DPPH (% quenching)
F 2011	2.19 ± 0.1	26.92 ± 0.77	109.71 ± 10.18	7.60 ± 1.04	80.05 ± 1.29
R 2008	1.09 ± 0.002	15.17 ± 0.19	37.69 ± 2.88	4.94 ± 0.67	53.60 ± 2.9




Figure 1: The effect of wines "R" (panel A) and "F" (panel B) on concentration – response curve to ADP. PRP was incubated with either wine (R or F, 10 μL) or with the vehicle (ethanol 12 %) and aggregation in response to ADP (1-30 μM) was evaluated. Each point represents the mean ± standard error of n = 5. * $p < 0.05$ and ** $p < 0.01$ vs. vehicle (two ways ANOVA followed by Bonferroni's test).

Both wines also inhibited washed platelet aggregation induced by THR (0.2 U/ mL); the biological activity was still present following wine dilution; also in this case "F", containing higher levels of flavonoids, anthocyanins, and phenols, and also increased antioxidant activity compared with "R", was more active than "R" (Figure 2).

Both wines increased ATP and ADPase activity evaluated in platelet lysates (Figure 3); this increase had a functional significance since CD39 inhibitor, ARL67156, reversed the inhibitory effect of wines on platelet aggregation without modifying vehicle – treated platelet aggregation (Figure 4).




Figure 2: The effect of different dilutions of wine “R” and of wine “F” on THR – induced aggregation of washed platelets. * $p < 0.05$ and $p < 0.001$ (one way ANOVA followed by Dunnett’s test, $n = 6$) vs. vehicle (ethanol, 12 %).




Figure 3: Effect of wine “R” (panel A) and wine “F” (panel B) on ATP – (panel A) and ADP (panel B) – hydrolysis mediated by CD39 by washed platelets. Platelet ATP – and ADPase activity was evaluated by measuring Pi released as described in the experimental section. * $p < 0.05$ and ** $p < 0.01$ vs. vehicle (one way ANOVA followed by Dunnett’s test; $n = 5$).




Figure 4: The effect of wine “R” (panel A) and wine “F” (panel B), or of the vehicle (ethanol 12%), on platelet aggregation induced by ADP (3 μM) in presence of CD39 inhibitor (ARL67156, 100 μM). * $p < 0.05$ vs. vehicle; ## $p < 0.01$ vs. R (A) and vs. F (B); $n = 5$.

In the end, following incubation with “F” wine, there was increased platelet expression of CD39. Conversely, wine “R” did not modify platelet CD39 expression (Figure 5).

Thus, we show that two different kinds of Aglianico red wines inhibit platelet aggregation, and concomitantly increase platelet ATP and ADPase activity; of particular interest is the finding that these effects may be obtained with the wine *in toto*. Furthermore, the youngest wine, “F”, was more active than “R”, in accordance with its content of total flavonoids, phenols and anthocyanin and its increased antioxidant activity compared with “R”. Thus, we suggest that a moderate wine consumption through increasing ATP-diphosphohydrolase (CD39) activity of platelets and avoiding ADP accumulation may offer protection from thrombosis.





Figure 5: Representative results of Western blot analysis of CD39 expression on platelets following incubation with wine “R” or “F” or with the vehicle (ethanol 12 %). Graph bar represents optical density (O.D.) relative to β actin. * $p < 0.05$ vs. vehicle (repeated measures ANOVA followed by Dunnett’s test, $n = 5$).

Experimental

Preparation of red wines: Two different “Aglianico” red wines were employed in the present study: one from the 2010 vintage (indicated with the letter F 2011) and the other from the 2007 vintage (named R 2008). Both wines were obtained following controlled microvinification procedures. Shortly, grapes from *Vitis vinifera* cv. were harvested from vineyards located in the area surrounding the town of Avellino (Campania, Italy). After grape destemming and crushing, must was treated with potassium metabisulfite (100 mg/L of must) and enzymes were added (2 g/hL Lafazym CL; Polsinelli, Frosinone, Italy). Subsequently, must was fermented in stainless steel tanks (100 L) at 22–23°C in the presence of 20 g/hL of *Saccharomyces cerevisiae* strain D254 (Lallemand Inc., Castel d’Azzano, Italy). Maceration of the pomace lasted 10 days; subsequently, the must was pressed to obtain the final wine. After 24 h, wines were racked from gross lees, 100 mg/L of potassium metabisulfite was added, and the wine bottled. No malolactic fermentation occurred. The wines were analyzed at bottling time. After opening, samples were divided into aliquots, stored in the dark at room temperature and used once for all experiments.

Total flavonoid, phenol and anthocyanin contents: The total flavonoid content was determined according to Bunea [16]. “F” and “R” wines (10 μL) or standard (quercetin up to 50 μM) were diluted to a final volume of 1 mL with distilled water. After the addition of 5% NaNO_2 (60 μL), the mixture was incubated at room temperature for 5 min. Then, an equal volume of 10% AlCl_3 was added and incubated for 6 min before the addition of 1 N NaOH (400 μL). Finally, absorbance was measured against blank at 510 nm. Total flavonoid content was expressed as millimolar of quercetin equivalent (QE). The total phenol contents of the wines were determined using the Folin-Ciocalteu’s method of Singleton and Rossi [17]. Briefly, red wines (1 μL) or standard (gallic acid up to 100 μM) were diluted with distilled water to 1 mL. After the addition of 50 μL Folin-Ciocalteu’s reagent, the mixture was incubated for 5 min at room temperature. At the end of incubation, 100 μL of 20% Na_2CO_3 was added and the absorbance measured at

760 nm. Results was expressed as mg/mL of gallic acid equivalents (GAE).The anthocyanin content of wines was estimated using a pH shift method [18]. The results were expressed as micromolar of malvidin-3-glucoside (M3GE), the major anthocyanin present in red wines, using the molar extinction coefficient of $28.000 \text{ M}^{-1} \text{ cm}^{-1}$.

Antioxidant activity of red wines: Antioxidant power was determined by 2 different methods. The Ferric Reducing Ability Power (FRAP) assay was as reported by Benzie [19] and results expressed in millimolar of a solution of pure FeSO_4 (up to $40 \mu\text{M}$). Briefly, red wines ($2 \mu\text{L}$) were added to a mixture containing 1.7 mM FeCl_3 and 0.85 mM of 2,4,6-Tris (2-pyridyl)-s-triazine (TPTZ) in 300 mM acetate buffer, pH 3.6 ($100 \mu\text{L}$ final volume) and incubated at 37°C for 6 min. Subsequently, absorbance was measured against blank at 590 nm (Microplate reader, Synergy HT BioTek, Milan, Italy).The second method to determine the antioxidant capacity, the 2,2-diphenyl-1-picrylhydrazyl (free radical DPPH) free radical scavenging assay, was performed according to the Wan-Ibrahim procedure, with some modification [20]. One mL of a methanolic solution of DPPH ($100 \mu\text{M}$) was incubated with “F” and “R” wines ($1 \mu\text{L}$), vortexed and incubated at room temperature for 30 min before absorbance measurement at 517 nm . The antiradical activity was defined as the amount of antioxidant necessary to decrease the initial DPPH absorbance. The radical scavenging activity was expressed as percentage of quenching.

Platelet preparation and aggregation: Blood was withdrawn by cardiac puncture from male Wistar rats (Harlan Nossan, 200-250 g) slightly anesthetized with enflurane, and anticoagulated with 3.8% (w/v) trisodium citrate (1:9, v/v). Platelet-rich plasma (PRP) and platelet-poor plasma (PPP) were prepared as previously described [21]. Platelet count in PRP was performed by a cell counter (Beckman Coulter s.r.l., Milano, Italy) and adjusted to 3×10^5 platelets / μL with autologous PPP. Platelet aggregation was monitored by a light transmission aggregometer (Chrono-Log, Cooperation, Mod.490, USA) by measuring changes in turbidity of 0.25 mL of re-calcified (CaCl_2 1 mM) PRP warmed at 37°C and under continuous stirring. A concentration response curve (1 - $30 \mu\text{M}$) to ADP was evaluated. To evaluate the effect of red wines “R” (1.09 mg/mL GAE) and “F” (2.19 mg/mL GAE) on ADP – induced platelet aggregation, $10 \mu\text{L}$ of either red wine (“R” or “F”) or of ethanol (12% , v/v) was added to PRP, and after 2 min incubation, a response curve to ADP (1 – $30 \mu\text{M}$) was performed. In some cases, the effect of different wine dilutions (ratio wine / ethanol, 0.5 ; 0.33 ; 0.10) was evaluated on aggregation induced by ADP ($3 \mu\text{M}$). Experiments were also performed in the presence of CD39 inhibitor, ARL67156 trisodium salt; in brief, following 1 min PRP incubation with either wine or vehicle, ARL67156 ($100 \mu\text{M}$ final concentration) or vehicle (distilled water) was added to the platelet suspension, and, after 1 min, aggregation to ADP ($3 \mu\text{M}$) was evaluated.

Washed platelets preparation: Platelets were washed by adding to PRP an equal volume of citric/trisodium citrate buffer (*plus* prostaglandin E_1 , $1 \mu\text{M}$) and centrifuged at $3000 \text{ rpm} \times 15 \text{ min}$. The platelet pellet was re-suspended in calcium - free Tyrode's buffer to the concentration $3 \times 10^5 / \mu\text{L}$. Platelet suspension was incubated with $10 \mu\text{L}$ of either wine or ethanol, as described above, and aggregation to thrombin (THR, 0.2 U/mL) was evaluated.

Western blot analysis: Washed platelets were incubated with either wine or ethanol, as described above, and then centrifuged at 2500

rpm at 25°C for 15 min. Platelets were homogenized in the following buffer: 50 mM Tris-HCl , pH 7.4; 150 mM NaCl ; 1 mM EDTA ; 1% sodium deoxycholate (0.25%); 1 mM phenylmethylsulphonyl fluoride; 10 mg/mL aprotinin; 20 mM leupeptin; 1 mM sodium orthovanadate. Total protein concentration in lysates was determined by Bradford assay, using BSA (bovine serum albumin) as standard. Proteins were separated on 8% SDS polyacrylamide gels and blotted onto nitrocellulose membrane. The membranes were saturated by incubation with 5% non-fat dry milk in PBS supplemented with 0.1% Tween-20 (PBS-T) for 30 min at room temperature and then incubated with anti-CD39 goat antibody ($1:200$; Santa Cruz Biotechnology, Italy) overnight at 4°C . Successively, membranes were washed and then incubated for 2 h at room temperature with the secondary antibody conjugated with horseradish peroxidase, and anti-goat IgG-HRP ($1:2000$; Dako, Denmark). Immune reactive proteins were visualized by enhanced chemiluminescence using Image Quant 400 GE Healthcare software (GE Healthcare, Italy). Successively, to confirm the equal protein loading, membranes were stripped and incubated with anti β -actin monoclonal antibody ($1:2000$; Sigma-Aldrich, Italy) and subsequently with anti-mouse IgG-HRP ($1:2000$; Dako, Denmark), both for 120 min at room temperature.

Measurement of platelet ATP - and ADPase activity: In another set of experiments, platelet lysates ($50 \mu\text{g}$) obtained from washed platelets incubated with red wines or vehicle, as described above, were pre-incubated in $200 \mu\text{L}$ of reaction buffer containing 10 mM NaCl ; 5 mM KCl ; 60 mM glucose; 5 mM CaCl_2 and 50 mM Tris-HCl buffer, pH 7.5, at 37°C for 10 min. The enzyme reaction was started by the addition of either ATP or ADP to a final concentration of 1 mM ; after 40 min at 37°C , the reaction was stopped by the addition of $200 \mu\text{L}$ of trichloroacetic acid (TCA). Following sample centrifugation at 3000 rpm for 10 min, at 37°C , the release of Pi was measured using malachite green as a colorimetric reagent and KH_2PO_4 as standard [22]. To have the net value of Pi produced following enzymatic reaction, aspecific Pi released in the absence of either ATP or ADP in each sample was evaluated and the value obtained was subtracted from the value obtained following incubation with the substrate. Protein concentration was measured by Bradford assay, using bovine serum albumin as standard, and results were expressed as Pi released $\text{nmol}/\mu\text{g}$ protein.

Statistical analysis: All results are expressed as mean \pm S.E. ($n=4$ - 6). Concentration – response curves were represented by non- linear regression and analysed by two way analysis of variance (ANOVA) followed by Bonferroni's test. In other cases, one way ANOVA followed by Bonferroni's or Dunnett's test, as appropriate, was used. All statistical analysis was performed using GraphPad, Prism V5.0 (Graphpad software, California, USA). A value of $p<0.05$ was considered significant.

Acknowledgments - We gratefully thank Dr Daniela Strollo from Mastroberardino S.r.l winery for providing the wines employed in the present study. We also thank Dr Stefania Bilotto for her help in performing the experiments. This work was also partially supported by a grant from program FESR Campania Region 2007/2013, objectives 2.1, 2.2, project CAMPUS-QUARC and by a grant from the Italian Ministry of Economy and Finance to the National Research Council for the project “Innovazione e Sviluppo del Mezzogiorno- Conoscenze Integrate per Sostenibilit edInnovazione del Made in Italy Agroalimentare -Legge 191/2009”.

References

- [1] Ruf JC. (2004) Alcohol, wine and platelet function. *Biological Research*, 37, 209-215.

- [2] Russo P, Tedesco I, Russo M, Russo GL, Venezia A, Cicala C. (2001) Effects of de-alcoholated red wine and its phenolic fractions on platelet aggregation. *Nutrition, Metabolism and Cardiovascular Diseases*, **11**, 25-29.
- [3] Zimmermann H, Zebisch M, Sträter N. (2012) Cellular function and molecular structure of ecto-nucleotidases. *Purinergic Signalling*, **8**, 437-502.
- [4] Atkinson B, Dwyer K, Enjyoji K, Robson SC. (2006) Ecto-nucleotidases of the CD39/NTPDase family modulate platelet activation and thrombus formation: Potential as therapeutic targets. *Blood Cells, Molecules and Diseases*, **36**, 217-222.
- [5] Deaglio S, Robson SC. (2011) Ectonucleotidases as regulators of purinergic signaling in thrombosis, inflammation, and immunity. *Advances in Pharmacology*, **61**, 301-332.
- [6] Fuentes E, Badimon L, Caballero J, Padró T, Vilahur G, Alarcón M, Pérez P, Palomo I. (2014) Protective mechanisms of adenosine 5'-monophosphate in platelet activation and thrombus formation. *Thrombosis and Haemostasis*, **111**, 491-507.
- [7] Qawi I, Robson SC. (2000) New developments in anti-platelet therapies: potential use of CD39/vascular ATP diphosphohydrolase in thrombotic disorders. *Current Drug Targets*, **1**, 285-296.
- [8] Robson SC, Kaczmarek E, Siegel JB, Candinas D, Koziak K, Millan M, Hancock WW, Bach FH. (1997) Loss of ATP diphosphohydrolase activity with endothelial cell activation. *The Journal of Experimental Medicine*, **185**, 153-163.
- [9] Robson SC. (2001) Thromboregulation by endothelial cells: significance for occlusive vascular diseases. *Arteriosclerosis, Thrombosis and Vascular Biology*, **21**, 1251-1252.
- [10] Schetinger MR, Morsch VM, Bonan CD, Wyse AT. (2007) NTPDase and 5'-nucleotidase activities in physiological and disease conditions: new perspectives for human health. *Biofactors*, **31**, 77-98.
- [11] Koziak K, Bojakowska M, Robson SC, Bojakowski K, Soin J, Cszimadia E, Religa P, Gaciong Z, Kaczmarek E. (2008) Overexpression of CD39/nucleoside triphosphate diphosphohydrolase-1 decreases smooth muscle cell proliferation and prevents neointima formation after angioplasty. *Journal of Thrombosis and Haemostasis*, **6**, 1191-1197.
- [12] Guckelberger O, Sun XF, Sévigny J, Imai M, Kaczmarek E, Enjyoji K, Kruskal JB, Robson SC. (2004) Beneficial effects of CD39/ecto-nucleoside triphosphate diphosphohydrolase-1 in murine intestinal ischemia-reperfusion injury. *Thrombosis and Haemostasis*, **91**, 576-586.
- [13] Jalkanen J, Yegutkin GG, Hollmén M, Aalto K, Kiviniemi T, Salomaa V, Jalkanen S, Hakovirta H. (2015) Aberrant circulating levels of purinergic signaling markers are associated with several key aspects of peripheral atherosclerosis and thrombosis. *Circulation Research*, **116**, 1206-1215.
- [14] Schmatz R, Mann TR, Spanevello R, Machado MM, Zanini D, Pimentel VC, Stefanello N, Martins CC, Cardoso AM, Bagatini M, Gutierrez J, Leal CA, Pereira LB, Mazzanti C, Schetinger MR, Morsch VM. (2013) Moderate red wine and grape juice consumption modulates the hydrolysis of the adenine nucleotides and decreases platelet aggregation in streptozotocin-induced diabetic rats. *Cell Biochemistry and Biophysics*, **65**, 129-143.
- [15] Spier AP, Bavaresco CS, Wyse ATS, Carvalho D, Freitas Sarkis JJ. (2007) Effects of resveratrol and purple grape juice on nucleotide hydrolysis by adult rat serum. *Food Chemistry*, **103**, 565-571.
- [16] Bunea A, Rugina OD, Pinteá AM, Sconța Z, Bunea CI, Socaciu C. (2011) Comparative polyphenolic content and antioxidant activities of some wild and cultivated blueberries from Romania. *Notulae Botanicae Horti Agrobotanici*, **39**, 70-76.
- [17] Singleton VL, Rossi JA. (1965) Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *American Journal of Enology and Viticulture*, **16**, 144-158.
- [18] Burns JI, Gardner PT, O'Neil J, Crawford S, Morecroft I, McPhail DB, Lister C, Matthews D, MacLean MR, Lean ME, Duthie GG, Crozier A. (2000) Relationship among antioxidant activity, vasodilation capacity, and phenolic content of red wines. *Journal of Agriculture and Food Chemistry*, **48**, 220-230.
- [19] Benzie IF, Strain JJ. (1996) The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. *Analytical Biochemistry*, **239**, 70-76.
- [20] Wan-Ibrahim WI, Sidik K, Kuppusamy UR. (2010) A high antioxidant level in edible plants is associated with genotoxic properties. *Food Chemistry*, **15**, 1139-1144.
- [21] Cicala C, Santacroce C, Itoh H, Douglas GJ, Page CP. (1997) A study on rat platelet responsiveness following intravenous endotoxin administration. *Life Science*, **60**, PL31-PL38.
- [22] Chan KM, Delfert D, Jung KD. (1986) A direct colorimetric assay for Ca^{2+} -stimulated ATPase activity. *Analytical Biochemistry*, **157**, 375-380.

A Validated HPLC Method for Simultaneous Determination of Caffeoyl Phenylethanoid Glucosides and Flavone 8-C-glycosides in <i>Haberlea rhodopensis</i> Dimitrina Zheleva-Dimitrova, Paraskev Nedialkov and Ulrich Giresser	791
A Novel Heterodimer from <i>Crotalaria ferruginea</i> Yi-Hong Zou, Xin Liu, Ye-Na Liu, Gui-Hua Tang and Sheng Yin	793
New 2-(2-Phenylethyl)chromone Derivatives and Inhibitors of Phosphodiesterase (PDE) 3A from Agarwood Takuji Sugiyama, Yuji Narukawa, Shunsuke Shibata, Ryo Masui and Fumiyuki Kiuchi	795
Influence of Different Carbohydrates on Flavonoid Accumulation in Hairy Root Cultures of <i>Scutellaria baicalensis</i> Chang Ha Park, Young Seon Kim, Xiaohua Li, Haeng Hoon Kim, Mariadhas Valan Arasu, Naif Abdullah Al-Dhabi, Sook Young Lee and Sang Un Park	799
Pyran-2-one Derivatives from the Roots of <i>Croton crassifolius</i> Weihuan Huang, Jijian Wang, Yeyin Liang, Yaolan Li and Guocai Wang	803
New Isocoumarins and Related Metabolites from <i>Talaromyces flavus</i> Jun-Wei He, Huai-Shuang Xu, Li Yang, Wei-Wei He, Chuan-Xi Wang, Feng Lin, Yang-Yun Lian, Bo-Hang Sun and Guo-Yue Zhong	805
A New Neolignan, and the Cytotoxic and Anti-HIV-1 Activities of Constituents from the Roots of <i>Dasymaschalon sootepense</i> Sakchai Hongthong, Chutima Kuhakarn, Thaworn Jaipetch, Pawinee Piyachaturawat, Surawat Jariyawat, Kanoknetr Suksen, Jitra Limthongkul, Narong Nuntasana and Vichai Reutrakul	809
New Phenyl Propanoids from <i>Cryptocarya bracteolata</i> Nurdin Saidi, Hiroshi Morita, Marc Litaudon, Mohd. Azlan Nafiah, Khalijah Awang and Mustanir	815
Anti-inflammatory Constituents from the Aerial Parts of <i>Iris minutiaurea</i> Kyeong Wan Woo, Ki Ho Lee, Ji Hun Jang, Min Suk Kim, Hyun Woo Cho, Jung Hee Cho and Byeongkwan An	817
New Aminonaphthoquinone from the Sea Urchins <i>Strongylocentrotus pallidus</i> and <i>Mesocentrotus nudus</i> Elena A. Vasileva, Natalia P. Mishchenko, Pavel A. Zadorozhny and Sergey A. Fedoreyev	821
Proteomic Analysis of the Antibacterial Mechanism of Action of Juglone against <i>Staphylococcus aureus</i> Jiayi Wang, Zixue Wang, Rina Wu, Donghua Jiang, Bing Bai, Dehong Tan, Tingcai Yan, Xiyun Sun, Qi Zhang, and Zhaoxia Wu	825
A Pair of New Polyketide Enantiomers from Three Endolichenic Fungal Strains <i>Nigrospora sphaerica</i>, <i>Alternaria alternata</i>, and <i>Phialophora</i> sp. Jun-Wei He, Chuan-Xi Wang, Li Yang, Guo-Dong Chen, Dan Hu, Liang-Dong Guo, Xin-Sheng Yao and Hao Gao	829
Insight into the Presence of Stilbenes in Medicinal Plants Traditionally Used in Croatian Folk Medicine Ivana Generalić Mekinić, Danijela Skroza, Ivica Ljubenkov and Višnja Katalinić	833
Two New Isomers of Palmityl-4-hydroxycinnamate from Flowers of <i>Taraxacum</i> Species Matej Dudáš, Mária Vilková, Tibor Béres, Miroslav Repčák and Pavol Mártonfi	837
Chemical Composition, Herbicidal and Antifungal Activity of <i>Satureja cuneifolia</i> Essential Oils from Spain David García-Rellán, Mercedes Verdeguer, Adele Salamone, María Amparo Blázquez and Herminio Boira	841
Composition of the Leaf Essential Oil of <i>Phoebe formosana</i> from Taiwan and its <i>in vitro</i> Cytotoxic, Antibacterial, and Antifungal Activities Yu-Chang Su and Chen-Lung Ho	845
Effects of Plant Density on the Number of Glandular Trichomes and on Yield and Quality of Essential Oils from <i>Oregano</i> Teresa Tuttolomondo, Salvatore La Bella, Claudio Leto, Giuseppe Bonsangue, Raffaele Leone, Maria Cristina Gennaro, Giuseppe Virga, Rosalinda Inguanta and Mario Licata	849
Antioxidant and Anti-inflammatory Activities of Essential Oils of <i>Actinodaphne macrophylla</i> and <i>A. pruinosa</i> (Lauraceae) Wan Mohd Nuzul Hakimi Wan Salleh and Farediah Ahmad	853
Chemical Composition, Antioxidant and Cytotoxic Activities of Essential Oil of the Inflorescence of <i>Anacamptis coriophora</i> subsp. <i>fragrans</i> (Orchidaceae) from Tunisia Ridha El Mokni, Saoussen Hammami, Stefano Dall'Acqua, Gregorio Peron, Khaled Faidi, Jeremy Phillip Braude, Houcine Sebei and Mohamed Hédi El Aouni	857
Essential Oil from <i>Origanum vulgare</i> Completely Inhibits the Growth of Multidrug-Resistant Cystic Fibrosis Pathogens Giovanna Pesavento, Valentina Maggini, Isabel Maida, Antonella Lo Nostro, Carmela Calónico, Chiara Sassoli, Elena Perrin, Marco Fondi, Alessio Mengoni, Carolina Chiellini, Alfredo Vannacci, Eugenia Gallo, Luigi Gori, Patrizia Bogani, Anna Rita Bilia, Silvia Campana, Novella Ravenni, Daniela Dolce, Fabio Firenzuoli and Renato Fani	861
<i>In Vitro</i> Trials of <i>Dittrichia graveolens</i> Essential Oil Combined with Antibiotics Dragoljub L. Miladinović, Budimir S. Ilić, Branislava D. Kocić, Marija S. Marković and Ljiljana C. Miladinović	865

Accounts/Reviews

Phytochemistry, Bioactivity and Potential Impact on Health of <i>Juglans</i>: the Original Plant of Walnut Dongdong Bi, Yicheng Zhao, Rui Jiang, Yan Wang, Yuxin Tian, Xiaoyi Chen, Shaojuan Bai and Gaimei She	869
---	-----

Natural Product Communications

2016

Volume 11, Number 6

Contents

Original Paper

- Pubinernoid A and Apo-9'-fucoxanthinone, Secondary Metabolites from a Gorgonian Coral *Pinnigorgia* sp.**
Hsin-Huan Chang, Yu-Chia Chang, Wu-Fu Chen, Tsong-Long Hwang, Lee-Shing Fang, Zhi-Hong Wen, Yu-Hsin Chen, Yang-Chang Wu and Ping-Jyun Sung 707
- A New Non-glucosidic Iridoid from the Roots of *Strychnos nux-blanda***
Jirapast Sichaem, Suttira Khumkratok, Pongpun Siripong and Santi Tip-pyang 709
- New Cytotoxic Guaianolides from *Centaurea aegyptiaca***
Hanan G. Sary, Abdel Nasser B. Singab and Khaled Y. Orabi 711
- Artemisinic Acid Serves as a Novel *ORCA3* Inducer to Enhance Biosynthesis of Terpenoid Indole Alkaloids in *Catharanthus roseus* Cambial Meristematic Cells**
Mingxuan Wang, Jiachen Zi, Jianhua Zhu, Shan Chen, Pu Wang, Liyan Song and Rongmin Yu 715
- New Antiplasmodial Diterpenes from *Gutierrezia sarothrae***
Qingxi Su, Seema Dalal, Michael Goetz, Maria B. Cassera and David G. I. Kingston 719
- A New Cassane-type Diterpene from the Seed of *Caesalpinia sappan***
Hai Xuan Nguyen, Nhan Trung Nguyen, Phu Hoang Dang, Phuoc Ho Thi, Mai Thanh Thi Nguyen, Mao Van Can, Dya Fita Dibwe, Jun-ya Ueda, Kinzo Matsumoto and Suresh Awale 723
- Two New Oleanane-type Triterpenoids from Methanolized Saponins of *Momordica cochinchinensis***
Rong Fan, Rong-Rong Cheng, Hong-Tao Zhu, Dong Wang, Chong-Ren Yang, Min Xu and Ying-Jun Zhang 725
- Isolation of a New Homomonoterpene from *Madhuca pasquieri* and Effect of Isolated Compounds on NO Production**
Le Son Hoang, Manh Hung Tran, Van Thu Nguyen, Quynh Mai Thi Ngo, Joo-Sang Lee, Seong-Soo Rho, Jeong Ah Kim, Jeong-Hyung Lee, Mi Hee Woo and Byung Sun Min 729
- New Bioactive Semisynthetic Derivatives of 31-Norlanostenol and Obtusifoliol from *Euphorbia officinarum***
Maria Bailen, Mourad Daoubi Khamlichi, Ahmed Benharref, Rafael A. Martinez-Diaz and Azucena Gonzalez-Coloma 733
- Variation of Ginsenosides in Ginseng of Different Ages**
Jian-Ming He, Yi-Zhen Zhang, Jian-Ping Luo, Wen-Ju Zhang and Qing Mu 739
- Colochiroside E, an Unusual Non-holostane Triterpene Sulfated Trioside from the Sea Cucumber *Colochirus robustus* and Evidence of the Impossibility of a 7(8)-Double Bond Migration in Lanostane Derivatives having an 18(16)-Lactone**
Alexandra S. Silchenko, Anatoly I. Kalinovsky, Sergey A. Avilov, Pelageya V. Andryjaschenko, Pavel S. Dmitrenko, Ekaterina A. Yurchenko, Igor Yu. Dolmatov, Salim Sh. Dautov, Valentin A. Stonik and Vladimir I. Kalinin 741
- Two New Steroidal Aglycones from Roots of *Cynanchun paniculatum***
Jia-wen Zhao, Feng-yang Chen, Li-juan Gao, Shi-fang Xu, Yi-ping Ye and Xiao-yu Li 747
- Steroid Sulfates from Ophiuroids (Brittle Stars): Action on Some Factors of Innate and Adaptive Immunity**
Anna K. Gazha, Lyudmila A. Ivanushko, Eleonora V. Levina, Sergey N. Fedorov, Tatyana S. Zaporozets, Valentin A. Stonik and Nataliya N. Besednova 749
- Cytotoxicity of Naturally Occurring Isoquinoline Alkaloids of Different Structural Types**
Jakub Chlebek, Ivo Doskočil, Daniela Hulcová, Kateřina Breiterová, Marcela Šafratová, Radim Havelek, Klára Habartová, Anna Hošťálková, Tereza Volštatová and Lucie Cahlíková 753
- Hypoglycemic and Antioxidant Activities of the Water Extract of *Aquilaria crassna* Leaves in Streptozotocin-Nicotinamide-Induced Type-2 Diabetic Mice**
Suchada Manok, Bunleu Sunthong, Hitoshi Sato, Erika Sugiyama and Vilasinee Hirunpanich Sato 757
- In Vitro* Antifungal Activity of Kambo Medicine Water Extracts against *Trichophyton rubrum***
Xia Da, Hitoshi Takahashi, Kyaw Zaw Hein and Eishin Morita 763
- Guarea kunthiana Bark Extract Enhances the Antimicrobial Activities of Human and Bovine Neutrophils**
Natalja Jerjomiceva, Hisham Seri, Ragheda Yaseen, Nicole de Buhr, William N. Setzer, Hassan Y. Naim and Maren von Köckritz-Blickwede 767
- Red Wine Inhibits Aggregation and Increases ATP-diphosphohydrolase (CD39) Activity of Rat Platelets *in Vitro***
Elisabetta Caiazzo, Idolo Tedesco, Carmela Spagnuolo, Gian Luigi Russo, Armando Ialenti and Carla Cicala 771
- How the Addition of Spices and Herbs to Virgin Olive Oil to Produce Flavored Oils Affects Consumer Acceptance**
Manel Issaoui, Guido Flamini, Sondess Soudi, Alessandra Bendini, Sara Barbieri, Ines Gharbi, Tullia Gallina Toschi, Pier Luigi Cioni and Mohamed Hammami 775
- Erythrina edulis* (Pajuro) Seed Protein: A New Source of Antioxidant Peptides**
Arturo Intiquilla, Karim Jiménez-Aliaga, Amparo I. Zavaleta, Inés Arnao, Carmen Peña, Elizabeth L. Chávez-Hidalgo and Blanca Hernández-Ledesma 781
- Cloning and Functional Analysis of Three Chalcone Synthases from the Flowers of Safflowers *Carthamus tinctorius***
Junichi Shinozaki, Hiromichi Kenmoku, Kenichi Nihei, Kazuo Masuda, Masaaki Noji, Katsuhiko Konno, Yoshinori Asakawa and Kohei Kazuma 787

Continued inside backcover