

Filarial infection caused by *Onchocerca boehmi* (Supperer, 1953) in a horse from Italy

Riccardo Paolo Lia¹ · Yasen Mutafovich^{1,2} · Vincenzo Veneziano³ · Alessio Giannelli¹ · Francesca Abramo⁴ · Mario Santoro⁵ · Maria Stefania Latrofa¹ · Cinzia Cantacessi⁶ · Coralie Martin⁷ · Domenico Otranto¹ · Andrea Bertuglia⁸ · Barbara Riccio⁹

Received: 18 July 2016 / Accepted: 28 September 2016
© Springer-Verlag Berlin Heidelberg 2016

Abstract Equids can be infected by a range of skin-dwelling filarial nematodes, including four species of the genus *Onchocerca*. Current literature on equine onchocercosis is fragmentary and often limited to isolated case reports. The present study aimed to describe a clinical case of equine onchocercosis caused by *Onchocerca boehmi* (Supperer, 1953) (syn. *Elaeophora boehmi*) in an 8-year-old gelding Belgian show jumper from northern Italy. The horse was presented with a firm and painless mass on the proximal third of the right metacarpal region. Ultrasound examination showed a peritendinous enlargement around the palmaro-lateral area of the tendons, characterized by an elongated hypoechoic and well-defined structure, embedding a coiled hyperechoic line.

The metacarpal nodule was resected and histologically examined. Fragments of a parasitic nematode were detected, isolated and examined. The morphological analysis allowed identifying the nematode as *O. boehmi*. In addition, total genomic DNA was extracted from individual fragments using a commercial kit for the nematode identification and a comparative sequence analysis of the nematode cytochrome oxidase subunit 1 (*cox1*) sequence with data available in the GenBankTM database revealed the closest identity (i.e. 91 %) with that of *Onchocerca lupi*. Thus far, *O. boehmi* has only been reported in Austria and Iran, and information about its life-cycle and vectors is lacking. The systematic position of this species within the genus *Onchocerca*, not in *Elaeophora* where it was originally described, is in concordance with the morphological and molecular analysis. In this article, we describe the first autochthonous case of equine onchocercosis in Italy caused by *O. boehmi* and discuss novel parasitological, clinical, and pathological data on these pathogens of horses.

✉ Riccardo Paolo Lia
riccardopaolo.lia@uniba.it

¹ Dipartimento di Medicina Veterinaria, Università degli Studi di Bari Aldo Moro, Valenzano, Bari, Italy

² Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, Sofia, Bulgaria

³ Dipartimento di Medicina Veterinaria e Produzioni Animali, Università di Napoli Federico II, Napoli, Italy

⁴ Dipartimento di Scienze Veterinarie, Università degli Studi di Pisa, Pisa, Italy

⁵ Istituto Zooprofilattico Sperimentale del Mezzogiorno, Portici, Napoli, Italy

⁶ Department of Veterinary Medicine, University of Cambridge, Cambridge, UK

⁷ UMR7245, MCAM, Museum National d'Histoire Naturelle, Paris, France

⁸ Dipartimento di Scienze Veterinarie, Università degli Studi di Torino, Torino, Italy

⁹ Veterinary practitioner, Torino, Italy

Keywords Equine onchocercosis · *Onchocerca boehmi* · Horse · Limb nodules · Ultrasound · Histology

Introduction

The genus *Onchocerca* (Spirurida, Onchocercidae) includes more than 30 species of nodule-inducing nematodes inhabiting different anatomical regions of the subcutaneous tissues, ligaments and aponeuroses of domestic mammals (Anderson 2000; Uni et al. 2015). The microfilariae released by the female nematodes migrate through the dermis of specific body areas, and they are ingested by blood feeding intermediate hosts (e.g. black flies and biting midges). In the vector, larvae moult twice, reaching the infective third larval stage (L3) in about 3–4 weeks. The L3s are then transmitted to

a susceptible vertebrate host via the blood meal (Onmaz et al. 2013). The infection is patent in about 12–16 months (Taylor et al. 2007). *Onchocerca reticulata* Diesing, 1841, and *Onchocerca cervicalis* Railliet and Henry, 1910, are the best-known filarial nematodes of equids due to their wide geographical distribution and potential clinical relevance (Muller 1979). In particular, *O. cervicalis* infection was firstly reported from Australia as “Queensland itch” (Riek 1953) and is characterised by the occurrence of an allergic dermatitis, likely induced by the skin-dwelling microfilariae (Lees et al. 1983). Microfilariae may also invade the eyes, causing ocular signs (Cello 1971; Munger 1983), while *O. cervicalis* adults may cause inflammatory reactions in the nuchal ligament, which range from acute oedematous necrosis to chronic granulomatous changes. Conversely, the infection by *O. reticulata* is usually characterised by the presence of subcutaneous nodules over or within the flexor tendons and suspensory ligaments, where it is potentially associated with swelling and lameness (Anderson 2000; Scott and Miller 2003). Equids may also be infected by *Onchocerca raillieti* Bain, Muller, Khamis, Guilhon and Schillhorn van Veen, 1976, a species detected in subdermal masses in the withers or penis and in the perimuscular conjunctive tissue of domestic donkeys in Africa (Bain et al. 1976). Another species of the genus, *Onchocerca boehmi* (Supperer, 1953) (syn *Elaeophora boehmi*), was described based on specimens collected in the arteries and veins of the limbs of horses from Austria. In most cases, horses infected by *O. boehmi* are asymptomatic (Supperer 1953).

The scientific literature on equine onchocercosis is fragmentary and often dated. For example, *O. cervicalis* has been long considered a synonym of *O. reticulata*, until Bain (1975) described the occurrence of morphological differences

between these two species. Similarly, epidemiological data on onchocercid species infecting horses are scarce. Infection by *O. cervicalis* has been diagnosed in the USA (Stannard and Cello 1975), Canada (Marcoux et al. 1977), Australia (Riek 1954), and Brazil (Marques and Scrofernecker 2004). In Europe, few studies have been performed (Anderson 2000), and onchocercids have seldom been identified at species level. In this article, we describe the first autochthonous case of equine onchocercosis in Italy caused by *O. boehmi* and discuss novel parasitological, clinical, and pathological data on these pathogens of horses.

Materials and methods

Case presentation

An 8-year-old 570 kg gelding Belgian horse, used in show-jump competitions, housed in northern Italy (Genoa, Liguria region, Italy), was presented in July 2013 at the Veterinary Teaching Hospital of the University of Turin (Piedmont, Italy) with an evident lump at the right metacarpal region. This lesion had appeared 6 months prior to presentation as a diffuse swelling, during the spring season, that had progressively increased in size. The owner sought the advice of clinicians in order to investigate the presence of a tendinitis in the mid-metacarpal region. During the clinical examination, the horse presented a firm and painless mass located palmaro-laterally on the proximal third of the right metacarpal region and a mild swelling in correspondence of the medial aspect of the left metacarpal region (Fig. 1). Numerous firm and small subcutaneous nodules were observed on the back of the animal, along the epiaxial muscles. The horse was mildly lame only at the beginning of the clinical presentation. Palpation did not allow defining the

Fig. 1 Right forelimb of the horse, showing a subcutaneous firm nodule in the palmaro-lateral aspect of the right metacarpal region. Lateral (a) and palmar (b) view of the limb



relationship of the mass with the superficial digital flexor tendon (SDFT). Previous treatments included DMSO (dimethyl sulfoxide) Gel 99.9 % as a topical application, twice daily over 3 weeks, to reduce the swelling. An oral administration of ivermectin paste was previously recommended by the practitioner, at double label dose (400 µg/kg body weight), on the basis of previous experience with similar subcutaneous nodules of suspected parasitic aetiology.

Ultrasonographic examination

An ultrasonographic examination was conducted using a mobile Logiq E Vet Ultrasound machine (General Electric Company Fairfield, CT, USA) with a linear multifrequency transducer (8–12 MHz). The examination was carried out on site, with the horse in standing position. No sedatives were administered. Prior to the ultrasound examination, both palmar metacarpal regions were prepared using standard procedures. Images were obtained using a standoff pad coupled to the transducer. The examination showed the presence of a peritendinous enlargement around the palmaro-lateral aspect of the SDFT, on the right forelimb, exerting a mass-effect on the whole soft tissues. The abnormal peritendinous mass was characterized by an elongated hypoechoic and well-defined structure, including a coiled hyperechoic line. On the left forelimb, the ultrasound examination revealed the same ultrasonographic pattern on the medial aspect of the mid metacarpal region, but with a more echogenic structure and lacking the hyperechoic linear structure. Ultrasonographic findings of both structures were consistent with a peritendinous localization of a verminous nodule (Fig. 2).

Surgical removal of the nodule

Surgical removal of the peritendinous mass was performed, with the horse standing and sedated using a constant infusion

rate. In particular, the infusion rate was prepared by adding 2 mg of medetomidine to a 0.5 l bag of saline (4 µg/mL), and this volume was administered at a rate of 1 drop/s (10 drops/mL infusion set drip rate), which provides approximately 80 min of infusion. A local analgesia was administered using a high metacarpal nerve block, with a 2 % solution of mepivacaine. The nodule was resected from the SDFT peritenon and the deep metacarpal fascia. Haemorrhage was controlled using an Esmark bandage, applied proximally to the carpal region. The skin was closed using routine procedures, and a half-limb bandage was applied post-operatively. Post-surgery standard anti-inflammatory and antibiotic therapies were administered over 3 days following the procedure, and the horse was not trained for 2 weeks post-surgery, and it had a complete functional outcome.

Histopathological analysis

Histopathological examination of the excised metacarpal nodule was performed; the tissue was fixed in a 10 % formalin solution (pH 7.4) and processed using standard procedures (Mutafchiev et al. 2013).

Parasitological and molecular analyses

A sub-section of the nodule was fixed and preserved in 70 % ethanol and dissected under a stereomicroscope. For light-microscopy, nematode fragments were cleared and examined as temporary mounts in lactophenol, while those used for scanning electron microscopy observations were prepared and studied, as described elsewhere (Mutafchiev et al. 2013). A female of *O. boehmi* (one microscopic slide) from the collection of Supperer deposited in the University of Veterinary Medicine Vienna (UVMV) was used as comparative material. In addition, total genomic DNA was extracted from parasite fragments recovered from an individual




Fig. 2 Transversal (**a, b**) and longitudinal (**c, d**) ultrasound scans of both mid metacarpal regions, showing a verminous nodule on the palmaro-lateral aspect of the right forelimb. The parasite appears as a coiled

hyperechoic line within a hypoechoic nodule, surrounding the superficial digital flexor tendon (**b**: red arrows). In the longitudinal scan (**d**) it is evident the localization at the level of the deep metacarpal fascia

specimen using a commercial kit (DNeasy Blood & Tissue Kit, Qiagen, GmbH, Hilden, Germany) in accordance with the manufacturer's instructions; a partial region of cytochrome c oxidase subunit 1 mitochondrial gene (*cox1*; ~689 bp) was amplified as previously described (Otranto et al. 2011). The amplicon obtained was purified using Ultrafree-DA columns (Amicon, Millipore; Bedford, USA) and sequenced directly using the Taq Dye Deoxy Terminator Cycle Sequencing Kit (v.2, Applied Biosystems) in an automated sequencer (ABI-PRISM 377). Sequences were determined from both strands (using the same primers individually as for the PCR), and the electropherograms were verified by eye. The nucleotide sequence of the *cox1* fragment was conceptually translated into amino acid sequences using the invertebrate mitochondrial code by MEGA 6.0 software (Tamura et al. 2013). Finally, the nucleotide sequence was compared with those available in the GenBankTM database by BLAST analysis.

Results

Histopathological analysis

Both haematoxylin and eosin and trichromic stains revealed a number of multifocal coalescing parasitic and necrotic granulomas. Each granuloma was characterised by a central cavity containing one or more parasitic sections (possibly due to coiled bodies); the cavity was lined by necrotic material and eosinophilic products of

degranulation, surrounded by macrophages and by an external layer of dense collagen. Parasitic granulomas were separated by a dense interstitial eosinophilic and macrophage infiltrate on a background of fibroplasia. Rare collagenolytic granulomas were scattered around the nodule. A visible body wall with an outer cuticle with subcuticular striations and an inner hypodermal layer could be visualised for some of the parasites. Small intestine and empty uteri were also observed. Based on their morphological features, the parasites were identified as nematodes (Fig. 3).

Morphological and molecular identification

Nematode fragments ($n = 83$) recovered from the nodule varied in length from 0.25 to 8.83 mm, amounting to 186 mm total length and a diameter ranging from 127 to 320 μm . The fragments contained only empty ovaries and were considered as belonging to an unknown number of unfertilized female nematodes (Fig. 4a, b). The anterior and posterior extremities were not detected. The cuticle was 16–25 μm thick with three distinct layers: an external layer 3–4 μm thick with transverse striations 7–12 μm apart interrupted along the medial lateral linings (Figs. 4c, d and 5a) and ornate with fine irregularly anastomosing crests (Figs. 4d, e and 5b); a median layer 10–18 μm thick, with annular striae with length corresponding to the distance between the external transverse striations (Fig. 4e) and an internal hyaline layer 3–5 μm thick. The somatic musculature was coelomyarian.

Fig. 3 Histopathology of the nodule. **a** Granulomatous reaction around a parasite: the cavity is lined by necrotic material with products of eosinophil degranulation (asterisk), macrophages and giant cells (>), collagen bundles, eosinophils and lymphocytes (trichrom stain); **b** Morphological details of a coiled parasite within a granuloma: small intestine, uteri (>) and lateral chord (*) (HE stain); **c** Subcuticular striations (HE stain); **d** Collagenolytic granuloma at the periphery of the nodule (HE stain)





Fig. 4 *Onchocerca boehmi*, light microscopy, horse from Italy. **a** Body fragment with intestine (arrow) and two uteri (arrowheads); **b** transverse section through body, note two uteri (arrowheads); **c** Surface of cuticle, note the interrupted externals transverse striations along median lateral line (C2); **d** Surface of cuticle exhibited when studied without coverslip, note internal striae (arrowheads), transverse striations (arrows) and ornamentation of fine irregularly anastomosing crests; **e** Detail of cuticle of two body fragments, note fine external crests on the surface (arrowheads) and internal annular striae of the median layer (arrows)



The morphological identification was confirmed by comparing samples with the voucher material collected by Supperer, which consisted of a single developing young and unfertilised female measuring 54.5 mm in length, without a posterior extremity. The specimen had a maximum body width of 170 µm at about mid-body and a width, measured at the level of vulva and oesophago-intestinal junction, of 104 µm; the oesophagus was 1259 µm long and vulva was situated at 575 µm from anterior body end. The cuticle at mid-body was 15–22 µm thick (thicker on lateral sides) with three distinct layers: an external layer 2 µm thick with fine transverse striations 3–5 µm apart, median layers 10–15 µm thick with annular striae with length coinciding with distance between external transverse striations and internal layer without specific detailed structure with regular thickness of 4–5 µm (Fig. 6).

A fragment of ~689 of the *cox1* gene was amplified. BLAST analysis of this sequence (Accession number

KX898458) revealed the highest nucleotide similarity (i.e. 91%) with that of *Onchocerca lupi* Rodonaja, 1967, available from GenBankTM (Accession Number EF521410).

Discussion

The present study describes a case of *O. boehmi* infection from a horse in Italy, where equine onchocercosis has never been reported, and it is therefore unknown to veterinary practitioners. In equine practice, the appearance of skin nodules is often asymptomatic, and it goes therefore unnoticed by owners (B. Riccio, personal communication). However, in the present report, the clinical presentation was accompanied by an impaired function of the suspensory ligament and occurrence of mild lameness. Interestingly, prior to this case, no clinical symptoms associated to infestation by *O. boehmi* had been described. Given the anatomical localisation of the nodules, we might hypothesize that the nematode




Fig. 5 *Onchocerca boehmi* scanning electron microscopy, horse from Italy. **a** Transverse striations (arrows) of cuticle surface; **b** Cuticle ornamentation, note transverse striations (arrows) and fine external crests (arrowheads)

undertook an erratic migration from the in circulatory system (i.e. the arteries and veins of limbs) to the subcutaneous tissues of the metacarpal region. *O. boehmi* had only been diagnosed in two isolated reports, and information about its biology is lacking. According to the original report by Supperer (1953), adults were detected in the medial or external layer of tissues within the artery wall in Austrian horses, while a second survey from Iran indicated




Fig. 6 *Onchocerca boehmi*, cuticle of young female, horse from Austria. Note the fine external crests (arrowheads) and the internal striae (arrows)

that 14 out 161 (8.69 %) horses examined had microfilariae in the blood (Mirzayans and Maghsoudloo 1977).

The occurrence of the parasite in the nodule allowed to assess the histopathological lesions caused by *O. boehmi*. Eosinophils were the main inflammatory cells found in the nodule, as reported in skin lesions caused by other *Onchocerca* species (Scott and Miller 2003). Besides a direct defence action against parasites, eosinophils play a pivotal role during hypersensitivity disorder and can also be detected in eosinophilic granulomas of horses, which are clinically characterized by the presence of cutaneous nodules and by histopathological occurrence of collagen flame figures (Scott and Miller 2003). Flame figures, albeit rare, were observed in the case herein described. Onchocercosis in horses can be characterised by both parasitic encystment (adult parasites) and hypersensitivity, the latter usually caused by microfilariae; nevertheless, dead or dying microfilariae were not observed in the tissue examined and the lesions were not pruritic.

The morphology of the cuticle of the nematode fragments collected resembled to that of the voucher material of *O. boehmi* from Austria; therefore, we consider both samples conspecific. In particular, while *O. boehmi* is characterized by cuticle without external ridges and three distinct layers with specific morphology, other *Onchocerca* parasitizing equids, (i.e. *O. cervicalis* and *O. reticulata*) have a cuticle with well-distinct external annular ridges (Bain 1981). Conversely, the cuticle of *O. railletii*, which is smooth and does not bear any external ridges, is thicker than that of *O. boehmi* (up to 50–55 µm vs 22–25 µm) and has longer striae (up to 16–20 µm vs 6–12 µm) (Bain et al. 1976; present study). The systematic position of this species within the genus *Onchocerca*, as suggested by Bain et al. (1976), and not in the genus *Elaeophora* is in concordance with the present morphological and molecular results.

Equine onchocercosis has been reported worldwide, but most epidemiological information date back from the 1970s. For instance, *Onchocerca* sp. has been diagnosed in horses from the USA, where Stannard and Cello (1975) reported a mean 48 % prevalence, whereas Lloyd and Soulsby (1978) recovered microfilariae in 61 % of examined animals from the eastern part of the country. Schmidt et al. (1982) examined the nuchal ligament of 83 horses from Midwestern US, and 37 % of them were positive for adult parasites. Klei et al. (1984) detected microfilariae in 76 % (out of 84) of ponies from the Gulf Coast area and in 82.4 % of horses (out of 51) from the Louisiana State. Of 664 horses from Southeaster and Midwestern USA, 341 (51.4 %) were positive for cutaneous microfilariae of *O. cervicalis* (Cummings and James 1985). Monahan et al. (1995) diagnosed *O. cervicalis* infection in 30.5 % (out of 82) of ponies in USA. Finally, Lyons et al. (2000) reported *O. cervicalis* in 24 % of horses (out of 157) examined for several species of internal parasites at necropsy in

Kentucky. Infection by *O. cervicalis* was reported also in Canada (Marcoux et al. 1977; Lees et al. 1983). Indeed, during a survey on 383 slaughtered horses from the western Canadian provinces, *O. cervicalis* microfilariae were detected in 11.8 % of umbilical samples (Polley 1984). Riek (1954) examined the nuchal ligaments of 282 Australian horses from Queensland and found that 79.8 % were infected with *Onchocerca* (erroneously reported as *O. reticulata*), whereas Ottley et al. (1983) sampled a small group of horses and ponies from Queensland and the Northern Territory, and diagnosed *O. cervicalis*, *O. gutturosa* and *O. reticulata* in those animals. In South America, Mancebo et al. (1997) found *O. cervicalis* microfilariae in 24 % of the 257 adult work horses examined in Argentina. A similar result was reported in Brazil by Marques and Scrofernecker (2004), who described *O. cervicalis* microfilariae in the midventral skin samples of 17.9 % (out of 1200) horses examined, while adult nematodes were recovered from the nuchal ligaments of 200 (16.6 %) animals. In Europe, a few studies have been performed thus far. In England, Mellor (1973) detected adult *Onchocerca* sp. in nuchal ligaments of 15.8 % (out of 209) British horses. Moignoux (1954) reported that 6 % of horses living in Camargue (France) were infected by subcutaneous *Onchocerca* microfilariae. However, Collobert et al. (1995) found that only 1 % (out of 368) of horses was positive for *Onchocerca* at post-mortem examinations in Normandy. In other European countries, out of 160 horse skin biopsies examined on horses in Spain and Poland showed a very low prevalence (3.7 %) of *Onchocerca* microfilariae (Franck et al. 2006). Finally, skin biopsies from 42 horses were all negative for microfilariae in Finland (Solismaa et al. 2008). These data indicate that equine onchocercosis is common in horse populations; however, as a consequence of the non-specific clinical presentation and challenging diagnosis, the infection prevalence is most likely underestimated. Additional large-scale studies are required to ascertain the presence and diffusion of *O. boehmi* and other onchocercid species in Italian and European horse populations.

The present case report suggests that parasitic granuloma should be included in the differential diagnosis of peritendinous swelling in horses and accurate ultrasound examination will permit to differentiate this condition from acute tendonitis or haematoma. The prevalence of parasitic granuloma associated with *O. boehmi* in the equine population is currently unreported, as well as its life cycle. Further studies are warrant to elucidate the biology of this poorly known onchocercid and its relationship with the equine species.

Acknowledgments The scanning electron microscope study was supported by the Bulgarian National Endowment Fund “13 Centuries Bulgaria”. Authors would like to thank Prof. Anja Joachim (UVMV) for lending the voucher material.

Conflict of interest statement The authors declare that they have no conflict of interest.

References

- Anderson RC (2000) The Superfamily Filarioidea. In: Nematode Parasites of Vertebrates; Their Development and Transmission, 2nd edn. CABI Publishing, New York, pp 517–523
- Bain O (1975) Redescription de cinq espèces d'onchocercques. Ann Parasitol Hum Comp 50:763–788
- Bain O (1981) Le genre *Onchocerca*: hypothèses sur son évolution et clé dichotomique des espèces. Ann Parasitol Hum Comp 56:503–526
- Bain O, Muller RL, Khamis Y, Guillon J, Schillhorn van Veen T (1976) *Onchocerca raillieti* sp. n. (Filarioidea) chez l'Ane domestique en Afrique. J Helminthol 50:287–293
- Cello RM (1971) Ocular onchocerciasis in the horse. Equine Vet J 3:148–154
- Collobert C, Bernard N, Lamidey C (1995) Prevalence of *Onchocerca* species and *Thelazia lacrimalis* in horses examined post mortem in Normandy. Vet Rec 136:463–465
- Cummings E, James ER (1985) Prevalence of equine onchocerciasis in southeastern and midwestern United States. J Am Vet Med Assoc 186:1202–1203
- Franck MT, Colombet J, Hugnet C, Ducos de Lahitte J, Desmaizières LM, Delverdier M, Franc M (2006) Research of skin microfilariae on 160 horses from Poland, France and Spain. Revue Méd Vét 157: 323–325
- Klei TR, Torbert B, Chapman MR, Foil L (1984) Prevalence of *Onchocerca cervicalis* in equids in the Gulf Coast region. Am J Vet Res 45:1646–1647
- Lees MJ, Kleider N, Tuddenham TJ (1983) Cutaneous onchocerciasis in the horse: five cases in Southwestern British Columbia. Can Vet J 24:3–5
- Lloyd S, Soulsby E (1978) Survey for infection with *Onchocerca cervicalis* in horses in eastern United States. Am J Vet Res 39: 1962–1963
- Lyons ET, Swerczek TW, Tolliver SC, Bair HD, Drudge JH, Ennis LE (2000) Prevalence of selected species of internal parasites in equids at necropsy in central Kentucky (1995–1999). Vet Parasitol 92:51–62
- Mancebo OA, Verdi JH, Bulman GM (1997) Comparative efficacy of moxidectin 2% equine oral gel and ivermectin 2% equine oral paste against *Onchocerca cervicalis* (Railliet and Henry, 1910) microfilariae in horses with naturally acquired infections in Formosa (Argentina). Vet Parasitol 73:243–248
- Marcoux M, Frechette JL, Morin M (1977) *Onchocerca cervicalis* infection in Quebec: clinical signs and diagnostic methods. Can Vet J 18: 108–110
- Marques SMT, Scrofernecker ML (2004) *Onchocerca cervicalis* in horses from Southern Brasil. Trop Anim Health Prod 36:633–636
- Mellor PS (1973) Studies on *Onchocerca cervicalis* Railliet and Henry 1910: I. *Onchocerca cervicalis* in British horses. J Helminthol 47: 97–110
- Mirzayans A, Maghsoudloo H (1977) Filarial infection of Equidae in the Tehran area of Iran. Trop Anim Health Prod 9:19–20
- Moignoux JB (1954) Enquête épidémiologique sur l'onchocercose cutanée des chevaux en Camargue. Rev Path Gen Comp 54:569–573
- Monahan CM, Chapman MR, French DD, Klei TR (1995) Efficacy of moxidectin oral gel against *Onchocerca cervicalis* microfilariae. J Parasitol 81:117–118
- Muller R (1979) Identification of *Onchocerca*. Symposia of the British Society for Parasitology, Vol. 17. In: Problems in the Identification of Parasites and their Vectors, ed. Taylor and Muller. Blackwell Scientific, London, pp 175–206
- Munger RJ (1983) Equine onchocercal keratoconjunctivitis. Equine Vet J 15:65–70

- Mutafchiev Y, Dantas-Torres F, Giannelli A, Abramo F, Papadopoulos E, Cardoso L, Cortes H, Otranto D (2013) Redescription of *Onchocerca lupi* (Spirurida: Onchocercidae) with histopathological observations. *Parasit Vectors* 6:309
- Onmaz AC, Beutel RG, Schneeberg K, Pavaloju AN, Komarek A, van den Hoven R (2013) Vector and vector-borne diseases of horses. *Vet Res Commun* 37:65–81
- Otranto D, Sakru N, Testini G, Gürlü VP, Yakar K, Lia RP, Dantas-Torres F, Bain O (2011) Case report: first evidence of human zoonotic infection by *Onchocerca lupi* (Spirurida, Onchocercidae). *Am J Trop Med Hyg* 84:55–58
- Ottley ML, Dallemagne C, Moorhouse DE (1983) Equine onchocerciasis in Queensland and the Northern Territory of Australia. *Aust Vet J* 60:200–203
- Polley L (1984) *Onchocerca* in horses from Western Canada and the Northwestern United States: An abattoir survey of the prevalence of infection. *Can Vet J* 25:128–129
- Riek RF (1953) Studies on allergic dermatitis (Queensland itch) of the horse. I. Description, distribution, symptoms and pathology. *Aust Vet J* 7:177–184
- Riek RF (1954) Studies on allergic dermatitis (Queensland itch) of the horse: the aetiology of the disease. *Aust J Agric Res* 5:109–129
- Schmidt GM, Krehbiel JD, Coley SC, Leid RW (1982) Equine onchocerciasis: lesion in the nuchal ligament of Midwestern US horses. *Vet Pathol* 19:16–22
- Scott DW, Miller WH (2003) *Equine dermatology*. Saunders, Elsevier Science, St. Louis, pp 242–245
- Solismaa M, Laaksonen S, Nylund M, Pitkänen E, Airakorpi R, Oksanen A (2008) Filarioid nematodes in cattle, sheep and horses in Finland. *Acta Vet Scand* 50:20
- Stannard AA, Cello RM (1975) *Onchocerca cervicalis* infection in horses from the western United States. *Am J Vet Res* 36:1029–1031
- Supperer R (1953) Filarosen der Pferde in Österreich. *Wiener Tierärztliche Monatsschrift* 40:193–220
- Uni S, Fukuda M, Agatsuma T, Bain O, Otsuka Y, Nakatani J, Matsubayashi M, Harada M, Omar H, Ramli R, Hashim R, Azirun MS, Hashim R (2015) *Onchocerca takaokai* n. sp. (Nematoda: Filarioidea) in Japanese wild boars (*Sus scrofa leucomystax*): description and molecular identification of intradermal females. *Parasitol Int* 64:493–502
- Tamura K, Stecher G, Peterson D, Filipowski A, Kumar S (2013) MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol Bio Evol* 30:2725–2729
- Taylor MA, Coop RL, Wall RL (2007) *Parasites of horses*. In *Veterinary Parasitology*. Blackwell, London, p 303–304