

Supplemental material of the article TSAF-D-15-00478:

TITLE

Morpho-anatomical and physiological traits of two Bougainvillea genotypes trained to two shapes under deficit irrigation

AUTHORS

Chiara Cirillo, Veronica De Micco, Youssef Roupheal, Angela Balzano, Rosanna Caputo, Stefania De Pascale*

Fig. S1.




Fig. S1.

Effect of the interaction between genotype [BxbR=*B. x buttiana* 'Rosenka'; Ba=*B. 'Lindleyana'* (= *B. 'Aurantiaca'*)] and irrigation treatment (C=control; DI= deficit irrigation) (a, b) and between genotype and canopy shape (Pyr=pyramid; Glo=globe) (c, d) on plant stem (a, c) and flower (b, d) biomass. Mean values and standard errors are shown. Different letters correspond to significantly different values (P ≤ 0.05).

Fig. S2.




Fig. S2.

Effect of the interaction among genotype [BxbR=*B. x buttiana* 'Rosenka'; Ba= *B. 'Lindleyana'* (= *B. 'Aurantiaca'*)], irrigation treatment (C=control; DI= deficit irrigation) and canopy shape (Pyr=pyramid; Glo=globe) on the length of stomata guard cells of two potted *Bougainvillea* genotypes. Mean values and standard errors are shown. Different letters correspond to significantly different values ($P \leq 0.05$).