

# Apolipoprotein A-I (ApoA-I) Mimetic Peptide P2a by Restoring Cholesterol Esterification Unmasks ApoA-I Anti-Inflammatory Endogenous Activity In Vivo<sup>[S]</sup>

Mariarosaria Bucci, Luisa Cigliano, Valentina Vellecco, Luca Domenico D'Andrea, Barbara Ziaco, Antonietta Rossi, Lidia Sautebin, Alessandro Carlucci, Paolo Abrescia, Carlo Pedone, Angela Ianaro, and Giuseppe Cirino

Departments of Experimental Pharmacology (M.B., V.V., A.R., L.S., A.I., G.C.) and Biological Sciences (L.C., A.C., P.A.), University of Naples "Federico II," Naples, Italy; and Biostructures and Bioimaging Institute, Consiglio Nazionale delle Ricerche, Naples, Italy (L.D.D., B.Z., C.P.)

Received October 24, 2011; accepted December 12, 2011

## ABSTRACT

The acute-phase protein haptoglobin (Hpt) binds apolipoprotein A-I (ApoA-I) and impairs its action on lecithin-cholesterol acyltransferase, an enzyme that plays a key role in reverse cholesterol transport. We have previously shown that an ApoA-I mimetic peptide, P2a, displaces Hpt from ApoA-I, restoring the enzyme activity in vitro. The aim of this study was to evaluate whether P2a displaces Hpt from ApoA-I in vivo and whether this event leads to anti-inflammatory activity. Mice received subplantar injections of carrageenan. Paw volume was measured before the injection and 2, 4, 6, 24, 48, 72, and 96 h thereafter. At the same time points, concentrations of HDL cholesterol (C) and cholesterol esters (CEs) were measured by high-performance liquid chromatography, and Hpt and ApoA-I plasma levels were evaluated by enzyme-linked immunosorbent assay. Western blotting analysis for nitric-oxide synthase

and cyclooxygenase (COX) isoforms was also performed on paw homogenates. CEs significantly decreased in carrageenan-treated mice during edema development and negatively correlated with the Hpt/ApoA-I ratio. P2a administration significantly restored the CE/C ratio. In addition, P2a displayed an anti-inflammatory effect on the late phase of edema with a significant reduction in COX2 expression coupled to an inhibition of prostaglandin E<sub>2</sub> synthesis, implying that, in the presence of P2a, CE/C ratio rescue and edema inhibition were strictly related. In conclusion, the P2a effect is due to its binding to Hpt with consequent displacement of ApoA-I that exerts anti-inflammatory activity. Therefore, it is feasible to design drugs that, by enhancing the physiological endogenous protective role of ApoA-I, may be useful in inflammation-based diseases.

## Introduction

Homeostasis of cholesterol (C) is essential for cell function and survival because cholesterol is toxic when it accumulates in the plasma membrane or within the cells. In atherogenesis, a critical role is played by the process known as reverse cholesterol transport (RCT), through which the accumulated cholesterol is transported from the vessel wall to the liver for excretion. This process includes acceptors such as high-den-

sity lipoproteins (HDLs), apolipoprotein A-I (ApoA-I), apolipoprotein E, lecithin-cholesterol acyltransferase (LCAT) (EC 2.3.1.43), and several other enzymes (Rader, 2006; Rader et al., 2009). Evidence has suggested a protective role for HDLs in atherosclerosis based on their ability to promote RCT (Rader, 2006). At present, it is still unclear whether part of the "protective" effect of HDLs is due to functions beyond RCT and whether these functions could be enhanced or reduced after the interaction with circulating proteins. Indeed, it is known that HDLs undergo dramatic modification in structure and composition as a result of the concerted actions of the acute-phase response protein and inflammation (Khovidhunkit et al., 2004; Esteve et al., 2005). In these conditions, HDL particles progressively lose normal biologi-

M.B. and L.C. contributed equally to this work.

Article, publication date, and citation information can be found at <http://jpet.aspetjournals.org>.

<http://dx.doi.org/10.1124/jpet.111.189308>.

[S] The online version of this article (available at <http://jpet.aspetjournals.org>) contains supplemental material.

**ABBREVIATIONS:** C, cholesterol; RCT, reverse cholesterol transport; HDL, high-density lipoprotein; ApoA-I, apolipoprotein A-I; LCAT, lecithin-cholesterol acyltransferase; Hpt, haptoglobin; CE, cholesteryl ester; BSA, bovine serum albumin; COX, cyclooxygenase; iNOS, inducible nitric-oxide synthase; PGE<sub>2</sub>, prostaglandin E<sub>2</sub>; P2as, scramble peptide; HPLC, high-performance liquid chromatography; ELISA, enzyme-linked immunosorbent assay; HDL-C, unesterified and total cholesterol; eNOS, endothelial nitric-oxide synthase.

cal activities and acquire altered properties (Kontush and Chapman, 2006). For example, the replacement of ApoA-I with serum amyloid A, occurring in small, dense HDLs upon induction of the acute phase of inflammation (Parks and Rudel, 1985; Coetzee et al., 1986), has been reported to have proatherogenic effects (Cabana et al., 1996; Lewis et al., 2004; O'Brien et al., 2005).

The critical role of inflammation at all stages of atherosclerosis is now recognized, including triggers, mediators, and end-effectors (Kontush and Chapman, 2006; Libby, 2007). In several studies, we have shown that the acute-phase protein haptoglobin (Hpt) binds the HDL protein components ApoA-I and apolipoprotein E, impairing their key functions in RCT (Porta et al., 1999; Spagnuolo et al., 2005; Salvatore et al., 2007, 2009; Cigliano et al., 2009). Haptoglobin is a polymorphic glycoprotein that exhibits phenotype prevalence in cardiovascular diseases (Delanghe et al., 1997). Its circulating levels are enhanced during the acute phase of inflammation to capture and transport free hemoglobin to the liver (Lanlois and Delanghe, 1996). ApoA-I, the major protein component of HDL, plays a key role in reverse cholesterol transport. This protein stimulates the efflux of cellular cholesterol and activates the enzyme LCAT, which in turn converts the free cholesterol into cholesteryl esters (CEs), addressing them to HDLs for transport into the circulation (Rader and Daugherty, 2008). The binding of Hpt to ApoA-I is associated with inhibition of LCAT activity and reduction of ApoA-I-mediated delivery of cholesterol to hepatocytes leading to 1) poor cholesterol removal from peripheral cells and 2) a low level of HDL cholesterol in the circulation (Spagnuolo et al., 2005; Salvatore et al., 2007; Cigliano et al., 2009). Moreover, several studies have confirmed that high levels of Hpt are associated with increased risk of developing cardiovascular events or myocardial infarction (Braeckman et al., 1999; De Bacquer et al., 2001; Matuszek et al., 2003). We have previously shown that an ApoA-I mimetic peptide with an amino acid sequence overlapping the stimulatory site for LCAT (P2a: acetyl-LSPLGGEEMRDRARAHVDALRTHLA-amide) efficiently displaced Hpt from ApoA-I. In addition, in an *in vitro* setting, P2a was able to rescue the stimulatory function of ApoA-I in the presence of high Hpt levels, whereas, when incubated without Hpt, it did not affect LCAT cholesterol esterification (Spagnuolo et al., 2005). Because the anti-inflammatory activity of HDLs and ApoA-I has been well documented (Kontush and Chapman, 2006; Rader, 2006; Gomaschi et al., 2008; Sherman et al., 2010), we hypothesized that the ApoA-I mimetic peptide P2a could also be effective *in vivo* in displacing Hpt from ApoA-I, leaving this apolipoprotein available for anti-inflammatory activity. In this study, we show that the peptide P2a rescues LCAT-dependent cholesterol esterification *in vivo* causing, in addition, an anti-inflammatory effect.

## Materials and Methods

**Drugs and Reagents.** Sheep anti-ApoA-I and sheep anti-Hpt polyclonal antibody were purchased from Serotec (Oxford, UK). [ $1\alpha,2\alpha$ - $^3\text{H}$ ]Cholesterol (45 Ci/mmol) was obtained from PerkinElmer Life and Analytical Sciences (Waltham, MA). SIL G plates for thin-layer chromatography (thickness 0.25 mm) were obtained from Macherey-Nagel (Düren, Germany). Chemicals of the highest purity, bovine serum albumin (BSA), cholesterol, cholesteryl linoleate, donkey anti-sheep IgG horseradish-linked, *o*-phenylenediamine, dextran sulfate (Dextralip 50), carrageenan, molecular mass markers,

and SUPELCOSIL LC-18 (5- $\mu\text{m}$  particle size,  $250 \times 4.6$  mm i.d.) were obtained from Sigma-Aldrich (St. Louis, MO). Polystyrene 96-well plates were purchased from Nunc (Roskilde, Denmark). Bradford reagent was obtained from Bio-Rad Laboratories (Milan, Italy). The antibodies against COX2 and iNOS were from Transduction Laboratories (Lexington, KY). [ $^3\text{H}$ ]PGE $_2$  was from Perkin-Elmer (Milan, Italy). All other reagents and compounds used were obtained from Sigma-Aldrich (Milan, Italy).

**Peptide Synthesis.** The peptide P2a (ApoA-I sequence from L141 to A164; acetyl-LSPLGGEEMRDRARAHVDALRTHLA-amide) and the scramble peptide control (P2as: acetyl-RLSARLTLHEGPVAL-DEMADRHA-amide) were solid phase-synthesized using PALPEG-PS resin (0.16 mmol/g) (Applied Biosystems, Foster City, CA) by standard *N*-(9-fluorenyl) methoxycarbonyl chemistry. Amino acid coupling was performed using a 10 M excess of *N*-(9-fluorenyl) methoxycarbonyl-amino acid, 9.9 Eq of 2-(1*H*-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate-1-hydroxybenzotriazole, and 20 Eq of *N,N*-diisopropylethylamine. A solution of 30% piperidine in *N,N*-dimethylformamide was used in the deprotection step (two 5-min washes). The coupling reaction was performed for 60 min followed by a 10-min acetylation step with acetic anhydride (2 M)-*N,N*-diisopropylethylamine (0.55 M)-1-hydroxybenzotriazole (0.06 M) in *N*-methylpyrrolidone. Cleavage from the resin was achieved by treatment with trifluoroacetic acid, triisopropyl silane, ethanedithiol, and water (94:1:2.5:2.5, v/v/v/v) at room temperature for 3 h. Peptide analysis and purification were performed by reverse-phase-HPLC on a C12 Proteo column (Phenomenex, Torrance, CA) using a gradient of CH $_3$ CN (0.1% trifluoroacetic acid) in water (0.1% trifluoroacetic acid) going from 20 to 50% in 30 min. Peptide identities were assessed by electrospray ionization mass spectrometry on a Thermo Finnigan MSQ liquid chromatograph-mass spectrometer (P2a: molecular mass experimental 2755.5 Da, molecular mass calculated 2756.1; P2as: molecular mass experimental 2755.8 Da, molecular mass calculated 2756.1 Da).

**Mouse Paw Edema.** Male Swiss mice (CD-1; Harlan, Correzzana Italy) weighing 28 to 30 g were used for *in vivo* experiments. The experimental procedures were performed in accordance with the *Guide for the Care and Use of Laboratory Animals* (National Institutes of Health, publication 86-23, revised 1985) as well as the specific guidelines of the Italian (N.116/1992) and European Council law (N.86/609/CEE). Animals were divided into groups ( $n = 6/\text{group}$ ) and lightly anesthetized with isoflurane. Each group of animals received subplantar injections of 50  $\mu\text{l}$  of carrageenan 1% (w/v) or 50  $\mu\text{l}$  of vehicle (saline) in the left hind paw. Paw volume was measured by using a hydroplethysmometer specially modified for small volumes (Ugo Basile, Comerio, Italy) immediately before the subplantar injection and 2, 4, 6, 24, 48, 72, and 96 h thereafter. The same operator always performed the double-blind assessment of paw volume. The increase in paw volume was calculated as the difference between the paw volume measured at each time point and the basal paw edema. Each group of animals received intraperitoneal (100  $\mu\text{l}$ ) administration of P2a peptide (0.3, 0.6, or 1 mg/kg), P2as peptide (1 mg/kg), or vehicle (saline). All peptides were administered immediately before the injection of carrageenan and 24 h thereafter.

**ApoA-I and Hpt Immunoassay.** The concentration of ApoA-I and Hpt was determined by ELISA using mouse antigens as standards for calibration, isolated as reported previously (Spagnuolo et al., 2003) and exhibiting more than 98% purity by electrophoresis and densitometry analysis. Aliquots of plasma (50  $\mu\text{l}$  from 1:1000, 1:10,000, 1:20,000, 1:45,000, 1:60,000, and 1:100,000 dilutions) were diluted in coating buffer (7.3 mM Na $_2$ CO $_3$ , 17 mM NaHCO $_3$ , and 1.5 mM NaN $_3$ , pH 9.6), loaded into the wells of a microtiter plate, and processed essentially according to a published procedure (Cigliano et al., 2005). In particular, sheep IgG (anti-ApoA-I or anti-Hpt, respectively) was used as the primary antibody and donkey anti-sheep IgG horseradish-linked IgG as the secondary antibody for color development. Measurements were performed using a calibration curve, obtained by determining the immunoreactivity of 1, 0.5, 0.25, 0.125,

0.065, 0.033, and 0.016 ng of standard Hpt protein and 0.4, 0.24, 0.16, 0.08, 0.04, and 0.016 ng of standard ApoA-I protein.

**LCAT Activity Assay.** Plasma samples from treated and control animals were treated with 50 mM  $\text{MnCl}_2$  and 0.1% dextran sulfate (50 kDa) according to a published procedure to remove very-low-density lipoprotein and low-density lipoprotein (Burstein et al., 1970). The LCAT activity in vitro was measured using proteoliposomes (ApoA-I/lecithin/cholesterol molar ratio of 1.5:200:18) as substrate as reported previously (Spagnuolo et al., 2005; Cigliano et al., 2009). The enzyme activity was expressed as enzyme units (nanomoles of cholesterol esterified per hour per milliliter of plasma).

**Titration of Cholesterol and Cholesteryl Esters in HDLs.** The ratio of CE with unesterified C in HDLs was assumed to reflect the LCAT activity in the plasma (Cigliano et al., 2005). Plasma samples of treated or control animals were treated with 50 mM  $\text{MnCl}_2$  and 0.1% dextran sulfate (50 kDa) according to a published procedure to remove very-low-density lipoprotein and low-density lipoprotein (Burstein et al., 1970). After this treatment, two 25- $\mu\text{l}$  aliquots were used for measuring the amounts of unesterified and total C. One aliquot was incubated (1 h; 75°C) with 0.25 ml of ethanol, whereas the other one was incubated with 0.25 ml of ethanol containing 5 M KOH. After incubation, both mixtures were supplemented with 0.15 ml of 1% NaCl and, after addition of 2 ml of ice-cold hexane, were vigorously shaken for 2 min. The hexane extract was added, and the lower phase was likewise extracted two more times. The three extracts were pooled and dried under a nitrogen stream. The residue was dissolved in 0.2 ml of acetonitrile-isopropanol (57:43, v/v), and 20  $\mu\text{l}$  were processed by reverse-phase HPLC. Chromatography was performed on a C18 column at 40°C with 1 ml/min flow rate, according to a published procedure (Cigliano et al., 2005). The amounts of unesterified and total cholesterol (HDL-C) were measured in samples processed without and with KOH, respectively, and used to calculate the amount of CEs as "total minus unesterified cholesterol." Calibration curves ( $r^2 \geq 0.9997$ ), obtained by injecting different amounts ( $n = 12$ ) of standard C, were used for quantitative analysis.

**Western Blot Analysis.** Paws from different groups of mice were harvested 24 and 48 h after carrageenan or vehicle injection and homogenized in modified radioimmunoprecipitation assay buffer (50 mM Tris-HCl, pH 7.4, 1% Triton X-100, 0.25% sodium deoxycholate, 150 mM NaCl, 1 mM EDTA, 1 mM phenylmethanesulfonyl fluoride, 10  $\mu\text{g}/\text{ml}$  aprotinin, 20 mM leupeptin, and 50 mM NaF) using a Polytron homogenizer (two cycles of 10 s at maximum speed) on ice. After centrifugation at 12,000 rpm for 15 min, the protein concentration was determined by Bradford assay using BSA as standard (Bio-Rad Laboratories), and 40  $\mu\text{g}$  of the denatured proteins were separated on 10% SDS-polyacrylamide gel electrophoresis and transferred to a polyvinylidene difluoride membrane. Membranes were blocked in phosphate-buffered saline-Tween 20 (0.1%, v/v) containing 5% nonfat dry milk and 0.1% BSA for 1 h at room temperature and then were incubated with anti-COX2 (1:1000) anti-iNOS (1:1000) overnight at 4°C. The filters were washed with phosphate-buffered saline-Tween 20 (0.1%, v/v) extensively for 30 min before incubation for 2 h at 4°C with the secondary antibody (1:5000) conjugated with horseradish peroxidase anti-mouse IgG. The membranes were then washed, and immunoreactive bands were visualized using ECL (GE Healthcare, Chalfont St. Giles, Buckinghamshire, UK).

**PGE<sub>2</sub> Exudate Levels.** Mice from different groups were euthanized 24 and 48 h after carrageenan or vehicle injection. Paws were cut and centrifuged at 4000 rpm for 30 min. Exudates (supernatants) were collected with 100  $\mu\text{l}$  of saline and used for PGE<sub>2</sub> quantification. To determine PGE<sub>2</sub> levels, proteins were removed from the exudates with 30%  $\text{ZnSO}_4$  for 15 min (Thomsen et al., 1990). PGE<sub>2</sub> was determined in deproteinized exudates by radioimmunoassay according to the manufacturer's instructions.

**Statistical Analysis.** Results are expressed as mean  $\pm$  S.E.M. or as mean  $\pm$  S.D. Statistical analysis was performed by analysis of


variance followed by the Bonferroni test for multiple comparisons or *t* test analysis where appropriate, using GraphPad Prism software (GraphPad Software Inc., San Diego, CA). Differences were considered statistically significant when  $p < 0.05$ . Each sample was processed at least in triplicate.

## Results


**Cholesterol Esterification Is Reduced in Mouse Edema.** The HDL fraction was isolated from plasma of carrageenan-treated mice after 2, 4, 6, 24, 48, 72, 96, and 144 h. The molar concentration of C and CEs was measured, and the molar ratio CE/C was calculated as an index of LCAT activity *ex vivo* (Subbaiah et al., 1997; Furbee et al., 2001).

In the early phase of carrageenan-induced inflammation (0–6 h), no differences were found in the CE/C ratio compared with that for the vehicle (data not shown). In the late phase of inflammation (24–48 h), the values of the CE/C ratio were found to be significantly decreased in carrageenan-treated mice ( $21.5 \pm 1$  at time 0 versus  $12 \pm 0.6$  at 24 h,  $p < 0.01$ ;  $21.5 \pm 1$  at time 0 versus  $10.5 \pm 0.9$  at 48 h,  $p < 0.01$ ) (Fig. 1), whereas no change was detected in controls. LCAT activity *in vitro* was not significantly modified by carrageenan injection at the time points tested (data not shown). The levels of apolipoprotein A-I, the protein that stimulates the enzyme LCAT, were found to be unchanged in inflamed mice (Fig. 2). These results suggest that, in carrageenan-treated mice, a plasma factor influences the cholesterol esterification, and this effect is not assessable *in vitro*.

**Correlation between the Ratio of Hpt/ApoA-I and CE/C.** Haptoglobin binds ApoA-I, the main stimulator of LCAT, thus inhibiting the enzyme activity and the efficiency of the reverse cholesterol transport (Spagnuolo et al., 2005; Salvatore et al., 2007; Cigliano et al., 2009). The plasma level of Hpt was measured in carrageenan-treated mice (Fig. 2). The change in Hpt level after carrageenan injection showed a biphasic trend: an early increase peaking 2 h after the carrageenan injection ( $29.1 \pm 1.5 \mu\text{M}$  in inflamed mice versus  $14.1 \pm 0.5 \mu\text{M}$  in controls;  $p < 0.01$ ) and a second late response peaking 48 h after ( $66.6 \pm 4 \mu\text{M}$  in inflamed mice versus  $13.8 \pm 2.8 \mu\text{M}$  in controls;  $p < 0.01$ ), returning to the physiological values at 144 h ( $14.2 \pm 3 \mu\text{M}$  in inflamed mice versus  $13.4 \pm 2.1 \mu\text{M}$  in controls). The Hpt/ApoA-I ratio



**Fig. 1.** C and CE ratio in HDLs isolated from mice plasma. Groups of three animals were sacrificed at 2, 4, 6, 24, 48, 72, 96, and 144 h after injection of carrageenan. The control group received vehicle only. Blood was collected, and the HDL fraction was isolated from plasma of carrageenan-treated mice and analyzed for C and CE content by HPLC. The molar concentrations of C and CE in HDLs were determined, and the molar ratio CE/C was calculated. The samples were analyzed in triplicate, and the data are expressed as means  $\pm$  S.D.




**Fig. 2.** Titration of Hpt in plasma from mice treated with carrageenan. Groups of three mice were sacrificed at 2, 4, 6, 24, 48, 72, 96, and 144 h after carrageenan injection. The control values are reported at time 0 in the graph. Blood was collected, and plasma from each animal was prepared. Samples were analyzed by ELISA for measuring the concentration of Hpt (●) and ApoA-I (□). The data are expressed as means  $\pm$  S.D.;  $n = 3$  separate experiments.


was found to be negatively correlated with the CE/C ratio ( $p < 0.01$ ;  $r = 0.94$ ) (Fig. 3). These data fit well with an inhibitory role of Hpt in the ApoA-I-dependent activity of cholesterol esterification on HDLs.

**P2a Peptide Inhibits Mouse Paw Edema in a Dose-Dependent Manner.** In the early phase of carrageenan-induced paw edema (0–6 h), no differences were found among groups treated with P2a at all doses tested compared with the vehicle group (Fig. 4). In the late phase of edema (24–96 h), injection of P2a caused a significant and dose-dependent inhibition of edema (Fig. 4). The specificity of the P2a effect was confirmed by the finding that the P2as did not affect edema development in both phases (Fig. 4).

**P2a Peptide Restores Cholesterol Esterification.** It is known that the binding of Hpt to ApoA-I is associated with reduced LCAT activity. We previously found that, in vitro, the peptide P2a, homologous to the ApoA-I sequence from Leu141 to Ala164, displaces Hpt from ApoA-I, restoring the activity of LCAT (Spagnuolo et al., 2005). To evaluate whether the P2a peptide displaces Hpt in vivo, different doses of P2a (0.3, 0.6, or 1 mg/kg i.p.) or of P2as were injected



**Fig. 3.** Correlation between Hpt/ApoA-I ratio and CE/C ratio. Plasma was obtained from carrageenan-treated mice and control mice at different time points. The plasma levels of Hpt and ApoA-I were measured by ELISA, and the molar concentrations of C and CE in the HDLs were analyzed by HPLC. Hpt/ApoA-I and CE/C ratios were calculated in triplicate. The Hpt/ApoA-I averages are plotted versus the homologous CE/C averages.




**Fig. 4.** Effect of P2a peptide on carrageenan-induced paw edema. P2a (0.3, 0.6, or 1 mg/kg), P2as (1 mg/kg), or vehicle was administered intraperitoneally immediately before the subplantar injection of carrageenan (50  $\mu$ l) and 24 h thereafter. Data are expressed as means  $\pm$  S.E.M.;  $n = 6$  for each group of treatment. \*\*,  $p < 0.01$ ; \*,  $p < 0.05$  versus vehicle.

in mice. When mice were treated with 0.6 or 1 mg/kg P2a, the CE/C ratio was found to be significantly restored (Fig. 5A). No effect of P2a on HDL-C was found (Fig. 5B). This result shows that P2a, administered intraperitoneally, efficiently engages Hpt, thus restoring ApoA-I function also in vivo.

**P2a Peptide Reduces COX2 Expression and PGE<sub>2</sub> Generation.** P2a exerted its anti-inflammatory action in the second phase of carrageenan-induced paw edema. To identify the molecular mechanisms responsible for this anti-inflammatory effect, the expressions of inducible isoforms of nitric oxide synthase and COX, i.e., iNOS and COX2, were evaluated at 24 and 48 h after carrageenan administration. As shown in Fig. 6, Western blot analysis did not reveal any significant alteration in COX2 expression at 24 h from edema induction in all three groups of mice treated with P2a. In contrast, 48 h after carrageenan injection, P2a (1 mg/kg) significantly reduced COX2 expression (Fig. 6). The COX2 involvement was also confirmed by the significant reduction of PGE<sub>2</sub> levels in paw exudates obtained from mice treated with P2a (Fig. 7). P2a administration did not affect iNOS and eNOS expression at any time point (supplemental figure).


## Discussion

It is known that the binding of Hpt to ApoA-I is associated with reduced LCAT activity in vitro (Spagnuolo et al., 2005). This binding, by decreasing the amount of free ApoA-I available for LCAT stimulation, impairs cholesterol esterification (Spagnuolo et al., 2005; Salvatore et al., 2007, 2009; Cigliano et al., 2009). The peptide P2a, which presents the ApoA-I amino acid sequence overlapping the domain required for LCAT stimulation, displaces Hpt from ApoA-I and rescues the enzyme activity in vitro (Spagnuolo et al., 2005). However, it has still not been determined whether this mechanism is relevant in vivo. To address this issue, we tested whether P2a could exert similar activity in vivo. By monitoring the plasma CE/C ratio and Hpt levels during edema development, we found an inverse correlation between CE/C




**Fig. 5.** A, effect of P2a on Hpt inhibition in cholesterol esterification. Blood was collected by intracardiac puncture from mice treated with P2a (0.3, 0.6, or 1 mg/kg i.p.) or P2as (1 mg/kg i.p.) 24 or 48 h after carrageenan injection. The HDL fraction was isolated from each plasma sample and analyzed for C and CE content by HPLC. Data are expressed as CE/C ratio. It is notable that the increase in the CE/C ratio was strictly related to the anti-inflammatory effect of P2a as described in Fig. 8C. The samples were analyzed in triplicate, and the data are expressed as means  $\pm$  S.D. For each group of treatment  $n = 6$ . \*\*,  $p < 0.001$ ; \*,  $p < 0.01$  versus treated mice 24 or 48 h after carrageenan injection. B, effect of P2a on HDL-C. The molar concentrations of C and CE in HDLs were determined as reported for A, and the total C was calculated as C + CE. The samples were analyzed in triplicate and the data are expressed as means  $\pm$  S.D.

and Hpt/ApoA-I ratios. The inflammatory response induced by carrageenan in mice is coupled with a significant increase in Hpt levels (Salvatore et al., 2007). Therefore, in this *in vivo* experimental model, Hpt acts as a competitive inhibitor of ApoA-I for LCAT activity, causing a reduction in cholesteryl ester production and confirming the pathophysiological relevance of this mechanism in inflammation. In the late phase of mouse paw edema, P2a displayed a clear anti-inflammatory effect. This effect did not involve changes in ApoA-I levels or LCAT activity, as confirmed by the finding that these two latter parameters were not modified in plasma harvested by mice treated with carrageenan. Thus, the anti-inflammatory activity of P2a should rely on different mechanism(s). We found that Hpt levels were elevated approximately 6-fold in the late phase of this inflammatory model. Thus, we hypothesize that Hpt, by binding ApoA-I, impairs LCAT function. In other words, as summarized for clarity in Fig. 8, the effect of P2a relies on the boosting of an endogenous mechanism, e.g., Hpt capture, rather than on a direct anti-inflammatory action. This hypothesis is sustained by the finding that systemic administration of P2a dose dependently increased the CE/C ratio, rescuing it to almost physio-




**Fig. 6.** Effect of P2a administration on COX2 expression in inflamed paws. Twenty-four and 48 h from carrageenan injection, mice paws harvested from different treatment groups were homogenized and Western blot analysis for COX2 was performed. A, representative Western blot for COX2. S, sham; V, vehicle; P2as, P2a scramble peptide; P2a 0.6, 0.6 mg/kg P2a; P2a 1, 1 mg/kg P2a. B, Western blot densitometric analysis for COX2 ( $n = 3$  experiments). Data are expressed as means  $\pm$  S.E.M. \*,  $p < 0.05$  versus sham; \*\*,  $p < 0.01$  versus sham; <sup>o</sup>,  $p < 0.05$  versus vehicle; <sup>oo</sup>,  $p < 0.01$  versus vehicle.



**Fig. 7.** Effect of P2a on PGE<sub>2</sub> levels in inflamed paws. Twenty-four and 48 h after carrageenan injection, mouse paws harvested from different groups of treatment were centrifuged, and exudates were collected and used for PGE<sub>2</sub> determination. Data are expressed as means  $\pm$  S.E.M. For each treatment group  $n = 4$ . \*\*,  $p < 0.01$ ; \*\*\*\*,  $p < 0.001$  versus sham; <sup>o</sup>,  $p < 0.05$ ; <sup>oo</sup>,  $p < 0.01$  versus vehicle.

logical levels at the higher dose tested (1 mg/kg). The specificity of the P2a effect was confirmed by the experiments performed using a scrambled peptide, for which the mismatching of amino acid positions led to a lack of efficacy. This effect of P2a on LCAT activity was paralleled by an anti-inflammatory effect on the late phase of carrageenan-induced paw edema (24–96 h). It is well known that in this mouse model of edema COX2-derived eicosanoids and/or iNOS- or eNOS-derived nitric oxide is involved at different time points (Posadas et al., 2004). P2a administration did not modulate eNOS expression. These findings correlate well with the lack of activity of P2a in the early phase of the edema. In the late phase, in which P2a significantly modifies the cholesterol biochemical pathway, there was also a reduction in COX2 expression as well as of PGE<sub>2</sub> levels. This finding implies



**Fig. 8.** Schematic depiction of P2a mechanism of action. A, in physiological conditions, the HDL ApoA-I domain interacts with LCAT, causing an increase in the CE/C ratio. B, Hpt is an acute-phase protein, and its levels are increased during the inflammatory response. This leads to increased binding of Hpt to ApoA-I and, by interference with LCAT activity, to a reduction in CE/C ratio (reverse cholesterol transport). C, P2a has the same amino acid sequence of the binding site for ApoA-I. After the treatment in vivo, P2a displaces Hpt from ApoA-I, restoring the CE/C ratio.

that the two phenomena observed in presence of P2a, i.e., CE/C ratio rescue and edema inhibition, are related.

Our data clearly imply that P2a, by virtue of its ability to bind Hpt, makes available more ApoA-I that acts as an endogenous anti-inflammatory signal. Indeed, the P2a anti-inflammatory effect is evident at the same time point at which recovery of the CE/C ratio occurs, i.e., 48 h after carrageenan injection. This hypothesis is further supported by the fact that P2a significantly inhibits exclusively the second phase of the edema at which the Hpt level reaches its maximum.

In conclusion, we have demonstrated that P2a, an ApoA-I-derived peptide with an amino acid sequence overlapping the stimulatory site for LCAT has an anti-inflammatory effect in vivo. Discoveries in the past decade have shed light on the complex metabolic and antiatherosclerotic pathways of HDLs. These insights have fueled the development of HDL-targeted drugs. In particular, many efforts were devoted to the design of ApoA-I mimetic peptides mimicking the functionality of ApoA-I (Kruger et al., 2005; Navab et al., 2005; Buga et al., 2006; Peterson et al., 2007; Degoma and Rader, 2011). Our study suggests that it is feasible to design drugs that can enhance the physiological endogenous protective role of ApoA-I, which may have an application in inflammation-based cardiovascular diseases such as atherosclerosis.

#### Authorship Contributions

*Participated in research design:* Bucci, Cigliano, Abrescia, and Cirino.

*Conducted experiments:* Bucci, Cigliano, Vellecco, Rossi, Carlucci, and Ziaco.

*Contributed new reagents or analytic tools:* D'Andrea and Ziaco.

*Performed data analysis:* Cigliano, Vellecco, and Ianaro.

*Wrote or contributed to the writing of the manuscript:* Bucci, Cigliano, Sautebin, Abrescia, Pedone, Ianaro, and Cirino.

#### References

- Braeckman L, De Bacquer D, Delanghe J, Claeys L, and De Backer G (1999) Associations between haptoglobin polymorphism, lipids, lipoproteins and inflammatory variables. *Atherosclerosis* **143**:383–388.
- Buga GM, Frank JS, Mottino GA, Hendizadeh M, Hakhamian A, Tillisch JH, Reddy ST, Navab M, Anantharamaiah GM, Ignarro LJ, et al. (2006) D-4F decreases brain arteriole inflammation and improves cognitive performance in LDL receptor-null mice on a Western diet. *J Lipid Res* **47**:2148–2160.
- Burstein M, Scholnick HR, and Morfin R (1970) Rapid method for the isolation of lipoproteins from human serum by precipitation with polyanions. *J Lipid Res* **11**:583–595.
- Cabana VG, Lukens JR, Rice KS, Hawkins TJ, and Getz GS (1996) HDL content and composition in acute phase response in three species: triglyceride enrichment of HDL a factor in its decrease. *J Lipid Res* **37**:2662–2674.
- Cigliano L, Pugliese CR, Spagnuolo MS, Palumbo R, and Abrescia P (2009) Haptoglobin binds the antiatherogenic protein apolipoprotein E—impairment of apolipoprotein E stimulation of both lecithin:cholesterol acyltransferase activity and cholesterol uptake by hepatocytes. *FEBS J* **276**:6158–6171.
- Cigliano L, Spagnuolo MS, Cuomo G, Valentini G, Niglio A, and Abrescia P (2005) Apolipoprotein A-I-dependent cholesterol esterification in patients with rheumatoid arthritis. *Life Sci* **77**:108–120.
- Coetzee GA, Strachan AF, van der Westhuyzen DR, Hoppe HC, Jeenah MS, and de Beer FC (1986) Serum amyloid A-containing human high density lipoprotein 3. Density, size, and apolipoprotein composition. *J Biol Chem* **261**:9644–9651.
- De Bacquer D, De Backer G, Langlois M, Delanghe J, Kesteloot H, and Kornitzer M

- (2001) Haptoglobin polymorphism as a risk factor for coronary heart disease mortality. *Atherosclerosis* **157**:161–166.
- Degoma EM and Rader DJ (2011) Novel HDL-directed pharmacotherapeutic strategies. *Nat Rev Cardiol* **8**:266–277.
- Delanghe J, Cambier B, Langlois M, De Buyzere M, Neels H, De Bacquer D, and Van Cauwelaert P (1997) Haptoglobin polymorphism, a genetic risk factor in coronary artery bypass surgery. *Atherosclerosis* **132**:215–219.
- Esteve E, Ricart W, and Fernández-Real JM (2005) Dyslipidemia and inflammation: an evolutionary conserved mechanism. *Clin Nutr* **24**:16–31.
- Furbee JW Jr, Francone O, and Parks JS (2001) Alteration of plasma HDL cholesterol ester composition with transgenic expression of a point mutation (E149A) of human LCAT. *J Lipid Res* **42**:1626–1635.
- Gomaschi M, Calabresi L, Rossoni G, Iametti S, Franceschini G, Stonik JA, and Remaley AT (2008) Anti-inflammatory and cardioprotective activities of synthetic high-density lipoprotein containing apolipoprotein A-I mimetic peptides. *J Pharmacol Exp Ther* **324**:776–783.
- Khovidhunkit W, Kim MS, Memon RA, Shigenaga JK, Moser AH, Feingold KR, and Grunfeld C (2004) Effects of infection and inflammation on lipid and lipoprotein metabolism: mechanisms and consequences to the host. *J Lipid Res* **45**:1169–1196.
- Kontush A and Chapman MJ (2006) Functionally defective high-density lipoprotein: a new therapeutic target at the crossroads of dyslipidemia, inflammation, and atherosclerosis. *Pharmacol Rev* **58**:342–374.
- Kruger AL, Peterson S, Turkseven S, Kaminski PM, Zhang FF, Quan S, Wolin MS, and Abraham NG (2005) D-4F induces heme oxygenase-1 and extracellular superoxide dismutase, decreases endothelial cell sloughing, and improves vascular reactivity in rat model of diabetes. *Circulation* **111**:3126–3134.
- Langlois MR and Delanghe JR (1996) Biological and clinical significance of haptoglobin polymorphism in humans. *Clin Chem* **42**:1589–1600.
- Lewis KE, Kirk EA, McDonald TO, Wang S, Wight TN, O'Brien KD, and Chait A (2004) Increase in serum amyloid A evoked by dietary cholesterol is associated with increased atherosclerosis in mice. *Circulation* **110**:540–545.
- Libby P (2007) Inflammatory mechanisms: the molecular basis of inflammation and disease. *Nutr Rev* **65**:S140–S146.
- Matuszek MA, Aristoteli LP, Bannon PG, Hendel PN, Hughes CF, Jessup W, Dean RT, and Kritharides L (2003) Haptoglobin elutes from human atherosclerotic coronary arteries—a potential marker of arterial pathology. *Atherosclerosis* **168**:389–396.
- Navab M, Anantharamaiah GM, Reddy ST, Hama S, Hough G, Frank JS, Grijalva VR, Ganesh VK, Mishra VK, Palgunachari MN, et al. (2005) Oral small peptides render HDL antiinflammatory in mice and monkeys and reduce atherosclerosis in ApoE null mice. *Circ Res* **97**:524–532.
- O'Brien KD, McDonald TO, Kunjathoor V, Eng K, Knopp EA, Lewis K, Lopez R, Kirk EA, Chait A, Wight TN, et al. (2005) Serum amyloid A and lipoprotein retention in murine models of atherosclerosis. *Arterioscler Thromb Vasc Biol* **25**:785–790.
- Parks JS and Rudel LL (1985) Alteration of high density lipoprotein subfraction distribution with induction of serum amyloid A protein (SAA) in the nonhuman primate. *J Lipid Res* **26**:82–91.
- Peterson SJ, Husney D, Kruger AL, Olszanecki R, Ricci F, Rodella LF, Stacchiotti A, Rezzani R, McClung JA, Aronow WS, et al. (2007) Long-term treatment with the apolipoprotein A1 mimetic peptide increases antioxidants and vascular repair in type I diabetic rats. *J Pharmacol Exp Ther* **322**:514–520.
- Porta A, Cassano E, Balestrieri M, Bianco M, Picone R, De Stefano C, and Abrescia P (1999) Haptoglobin transport into human ovarian follicles and its binding to apolipoprotein A-1. *Zygote* **7**:67–77.
- Posadas I, Bucci M, Rovizzo F, Rossi A, Parente L, Sautebin L, and Cirino G (2004) Carrageenan-induced mouse paw oedema is biphasic, age-weight dependent and displays differential nitric oxide cyclooxygenase-2 expression. *Br J Pharmacol* **142**:331–338.
- Rader DJ (2006) Molecular regulation of HDL metabolism and function: implications for novel therapies. *J Clin Invest* **116**:3090–3100.
- Rader DJ, Alexander ET, Weibel GL, Billheimer J, and Rothblat GH (2009) The role of reverse cholesterol transport in animals and humans and relationship to atherosclerosis. *J Lipid Res* **50**:S189–S194.
- Rader DJ and Daugherty A (2008) Translating molecular discoveries into new therapies for atherosclerosis. *Nature* **451**:904–913.
- Salvatore A, Cigliano L, Bucci EM, Corpillo D, Velasco S, Carlucci A, Pedone C, and Abrescia P (2007) Haptoglobin binding to apolipoprotein A-I prevents damage from hydroxyl radicals on its stimulatory activity of the enzyme lecithin-cholesterol acyl-transferase. *Biochemistry* **46**:11158–11168.
- Salvatore A, Cigliano L, Carlucci A, Bucci EM, and Abrescia P (2009) Haptoglobin binds apolipoprotein E and influences cholesterol esterification in the cerebrospinal fluid. *J Neurochem* **110**:255–263.
- Sherman CB, Peterson SJ, and Frishman WH (2010) Apolipoprotein A-I mimetic peptides: a potential new therapy for the prevention of atherosclerosis. *Cardiol Rev* **18**:141–147.
- Spagnuolo MS, Cigliano L, and Abrescia P (2003) The binding of haptoglobin to apolipoprotein AI: influence of hemoglobin and concanavalin A. *Biol Chem* **384**:1593–1596.
- Spagnuolo MS, Cigliano L, D'Andrea LD, Pedone C, and Abrescia P (2005) Assignment of the binding site for haptoglobin on apolipoprotein A-I. *J Biol Chem* **280**:1193–1198.
- Subbaiah PV, Liu M, and Witt TR (1997) Impaired cholesterol esterification in the plasma in patients with breast cancer. *Lipids* **32**:157–162.
- Thomsen LL, Ching LM, and Baguley BC (1990) Evidence for the production of nitric oxide by activated macrophages treated with the antitumor agents flavone-8-acetic acid and xanthenone-4-acetic acid. *Cancer Res* **50**:6966–6970.

---

**Address correspondence to:** Dr. Giuseppe Cirino, Department of Experimental Pharmacology, Faculty of Pharmacy, University of Naples “Federico II,” via Domenico Montesano 49, 80131 Naples, Italy. E-mail: cirino@unina.it

---