

Tinnitus psychopharmacology: A comprehensive review of its pathomechanisms and management

Michele Fornaro
Matteo Martino

Department of Neuroscience, Section
of Psychiatry, University of Genova,
Genova, Italy

Background: Subjective tinnitus is a frequent, impairing condition, which may also cause neurotransmitter imbalance at the cochlea. Psychopharmacologic agents, although not being the first-line treatment for tinnitus, may modulate cochlear neurotransmission, thereby influencing the subjective tinnitus experience.

Method: A comprehensive review of MEDLINE literature (from January 1990–January 2010) was performed searching for: “tinnitus”, major classes of psychopharmacological agents, and psychiatric disorders. The most relevant clinical evidence is reported briefly along with a concise description of the main neurotransmitters purported to be involved in tinnitus, in order to provide the reader with a rational evaluation of tinnitus therapy with psychopharmacological agents.

Results: Although strong methodological issues limit the reliability of the current results, a broad number of psychopharmacological agents have already been considered for tinnitus, both as candidate triggers or potential therapies.

Conclusions: Selected psychopharmacological drugs may play a role in the clinical management of this disorder. While the rational use of these agents for the treatment of tinnitus should not be overlooked, research should be undertaken on their neuromodulating actions at the cochlea.

Keywords: tinnitus, psychopharmacology, cochlea

Introduction

Tinnitus, from the Latin word “*tinntus*” meaning “ringing”¹ is a perception of sound in the absence of external sounds (ICD-10 code, H93.1).² This phenomenon can be divided into two broad groups: *objective* and *subjective* tinnitus. Objective tinnitus is provoked by sound generated in the body reaching the ear through conduction in body tissues, while subjective tinnitus is meaningless sounds that are not associated with any physical sound and it can be heard only by the suffering person. Although subjective tinnitus represents a far more prevalent condition compared to objective tinnitus,³ and thereby constitutes a more accessible phenomenon for investigators, current information available on its etiology is unsatisfactory. In fact, while a broad number of heterogeneous pathomechanisms and causes have been postulated (Table 1), no univocal consensus has been reached to date and its management is still a well debated issue. Furthermore, since there are many kinds of subjective tinnitus, the search for a (unique) cure is futile. As consequence, both researchers and clinicians have progressively extended the portfolio of potential therapies (including nonfirst-choice treatments such as psychopharmacological agents) especially for those patients whose tinnitus may be related to comorbid psychological stressors.⁴

Correspondence: Michele Fornaro
Ospedale San Martino, Largo Rosanna
Benzi n.16, ZIP 16100, Genova, Italy
Tel +39 347 414 0003
Fax +39 0103537669
Email dott.fornaro@gmail.com

Table 1 Main hypothesized pathomechanisms and causes of tinnitus**Main peripheral auditory system theories of subjective tinnitus**

Spontaneous otoacoustic emissions
Edge theory
Discordant theory

Small acoustic signals perceived as tinnitus
Increased spontaneous activity in the edge area
Discordant dysfunction of damaged outer hair cells and intact inner hair cells

Main central auditory system and somatosensory theories of subjective tinnitus

The dorsal cochlear nucleus
Auditory plasticity theory
Crosstalk theory

Hyperactivity/plastic readjustment of DCN
Enhanced neural activity due to cochlear damage
Ephaptic coupling between nerve fibers

Main causes of subjective tinnitus

Otologic problems and hearing loss

Loud noise, presbycusis, Ménière's disease, acoustic neuroma, external ear infection

Pharmacological causes

Analgesics, antibiotics, chemotherapy and antiviral drugs, loop diuretics, antidepressants, psychedelic drugs (5-MeO-DET, 5-Methoxy-diisopropyltryptamine, diisopropyltryptamine, harmaline, N,N-dimethyltryptamine, psilocybin, salvinorin A)

Neurologic disorders

Traumatic brain injury, meningitis, encephalitis, strokes, multiple sclerosis, chiari malformation, auditory nerve injury

Metabolic disorders

Thyroid disorder, hyperlipidemia, vitamin B12 deficiency, iron deficiency, anemia

Psychiatric disorders

Anxious and depressive states

Other

Tension myositis syndrome, fibromyalgia, head and neck muscle spasm, temporomandibular joint disorders, thoracic outlet syndrome, lyme disease, hypnagogia, sleep paralysis, glomus tympanicum, herpes infections

Main causes of objective tinnitus

Pulsatile tinnitus: Altered blood flow or increased blood turbulence near the ear

Atherosclerosis, venous hum, carotid artery aneurysm, carotid artery dissection

Muscle contractions that cause clicks or crackling around the middle ear

Since some forms of tinnitus may be exacerbated or moderated by psychopharmacological agents, the aim of the present paper is to provide the reader with an updated survey on their rational use in subjective tinnitus.

Data source and selection

A comprehensive review of English-written MEDLINE results was retrieved using the following queries: "Tinnitus AND neurobiology"; "Tinnitus AND therapy"; "Tinnitus AND antidepressant"; "Tinnitus AND SSRI"; "Tinnitus AND venlafaxine"; "Tinnitus AND duloxetine"; "Tinnitus AND bupropion"; "Tinnitus AND tricyclic"; "Tinnitus AND benzodiazepine"; "Tinnitus AND lithium"; "Tinnitus AND antiepileptic"; "Tinnitus AND anxiety"; "Tinnitus AND depression"; "Tinnitus AND bipolar disorder"; "Tinnitus AND schizophrenia"; "Tinnitus AND ADHD"; and "Tinnitus AND dementia". Results were kept within a January 1990–January 2010 time limit. Finally, meta-analysis and randomized clinical trials (RCTs) data were prioritized when available.

Pathophysiology

The changes in the auditory nervous system, especially at the dorsal (DCN) and ventral cochlear nucleus (VCN) underpinning tinnitus are often represented by a reduction in the inhibitory rather than an excitatory input, resulting in a shift in the balance between inhibition and excitation.

Deprivation of input may cause neural plasticity to change the relationship between inhibition and excitation and protein synthesis⁵ and finally, impact on neurotransmission—especially at the DCN, the inferior colliculus (IC), together with the primary and secondary auditory cortices—eventually affecting the thalamic and dorsal cortex transmission, the amygdala, and other structures.⁶ The rerouting of information may cause structures of the central nervous system (CNS) that are not normally involved in processing auditory information to become activated by sound stimulation (ie, the abnormal involvement of the nonclassical–nonspecific/extralemniscal–pathways). Yet to date, no univocal or exhaustive appreciation of tinnitus determining neural abnormalities has been

reached; with its pathophysiological correlates still remaining a debated issue.

Among others, the following evidence and hypotheses have been postulated:⁷

Peripheral auditory system

Spontaneous otoacoustic emissions

Small acoustic signals presumed to be generated by the electromotile activity of the outer hair cells (OHCs) of the cochlea and propagated into the external auditory canal may be abnormally perceived as tinnitus.⁸

Edge theory

Increased spontaneous activity in the edge area, which represents a transition from normal OHCs on the apical side of a lesion to OHCs toward the basal side that are missing or altered, may contribute to tinnitus perception.⁹

Discordant theory

The discordant dysfunction of damaged OHCs and intact inner hair cells (IHCs) may result in the disinhibition and increased spontaneous activity of neurons in the DCNs, that receive excitation from IHCs, but with no inputs from the damaged OHCs, therefore playing a role in tinnitus phenomenon.¹⁰

Central auditory system

The dorsal cochlear nucleus

Hyperactivity by disinhibition or plastic readjustments of the DCN, triggered for instance by OHC damage or a reduction in auditory nerve input have been supposed to concur in tinnitus genesis.¹¹

Auditory plasticity theory

Damage to the cochlea enhances neural activity in the central auditory pathway, such as the IC and the temporal lobe of the auditory association cortex (similar to the phantom limb sensation in amputees).¹²

Crosstalk theory

The compression from blood vessels or a tumor on auditory nerve causes ephaptic coupling between nerve fibers, which may result in the phase-locking of spontaneous activity of groups of auditory neurons that may be perceived as sound.¹³

Somatosensory system

Abnormal DCN activity could also be influenced by the stimulation of the somatosensory system. Furthermore, pain

signals from the cochlea carried by the cochlear C fibers may be interpreted by the CNS as tinnitus.¹⁴

Neurotransmitters

It has been postulated that numerous cochlear neurotransmitters play a role in tinnitus sensation.

GABA

It is known that the number of gamma-aminobutyric acid (GABA) immunoreactive neurons in the auditory nuclei decreases with age;¹⁵ together with its concentration and release, the glutamatergic acid decarboxylase activity (the rate-limiting enzyme in the formation of GABA), GABA_B receptor binding and the presynaptic GABA releasing terminals availability.¹⁶ This trend has been postulated to be a higher risk of noise-induced hearing loss as an important tinnitus etiological factor, especially in older men¹⁷, and has prompted both researchers and clinicians to consider GABAergic modulators as a candidate therapy for tinnitus patients. Anxiolytic benzodiazepines (BDZ), as well as anti-epileptic drugs that enhance GABA_A receptors (including those used as mood-stabilizers); or GABA_B agonists (eg, baclofen, a drug used for multiple sclerosis and proposed for morphine withdrawal syndrome¹⁸) have all been suggested both for animal and human models of tinnitus.^{19,20} Finally, while anxiety *per se* does not produce tinnitus, it may strongly exacerbate its perception, and may be modulated by GABAergic neurotransmission enhancers with the final result of tinnitus relief.

Serotonin

Serotonergic (5-HTergic) modulation (eg, by salicylate administration) has been reported to provoke tinnitus,²¹ although human genetic studies on the regulatory region of the 5-HT transporter (5-HTTLPR) SLC6A4 gene failed to show any significant effect on the susceptibility to chronic tinnitus, thus indicating an indirect, rather than a direct, modulating role of 5-HT in tinnitus etiology and CNS-adaptive mechanisms.²² Modulation of 5-HT is a core mechanism of a very large number of routinely prescribed drugs including the selective serotonin reuptake inhibitor (SSRI) antidepressants, thus their putative role in increasing the sensation of tinnitus should therefore be taken into account.²³ Nonetheless, the biological theory of depression provides a rationale for the association of emotional distress and chronic tinnitus determined by the respective impact of dysfunctional neural reorganization on limbic and auditory structures,²⁴ with a specific role of brain-derived

neurotrophic factor (BDNF) in promoting plasticity-induced recovery, has been postulated in depressive states,²⁵ which could make some affective (or anxiety) disordered individuals more genetically prone²⁶ or stress-vulnerable²⁷ for developing tinnitus.

Dopamine

Tinnitus perception takes place in prefrontal, primary temporal and temporoparietal associative areas, as well in the limbic system. Dopamine (DA) neurotransmission acts through prefrontal, primary temporal, temporoparietal associative areas and the limbic system. Tinnitus perception and DA-ergic pathway share the same cerebral structures, which control attention, stress, emotions, learning, memory and motivated behavior. Distress from tinnitus emanates from these same cerebral functions, and could therefore be potentially modulated by either DA-blockers (ie, typical and atypical antipsychotics) or DA-agonists (eg, bupropion, a norepinephrine/DA reuptake inhibitor, antidepressant and by many anti-Parkinson drugs such as levodopa, bromocriptine, pramipexole and others).

Glutamate

N-methyl D-aspartate (NMDA) glutamate receptor is most likely involved in tinnitus, as it is involved in many forms of central neuropathic pain.²⁸ Specifically, it has been shown that acetylsalicylic acid activates cochlear NMDA receptors²⁹ and that the use of NMDA-antagonists at the round window abolishes tinnitus,²⁹ while NMDA receptor agonists may induce tinnitus-like behaviors.³⁰ Memantine, used to treat neuropathic pain and mild to moderate forms of Alzheimer's disease,³¹ blocks NMDA transmission in hair cells (the same as salicylate acts upon)³² also modulating the cholinergic transmission. Acamprosate, a drug acting as both a NMDA-blocker and GABA-enhancer is used for alcohol withdrawal maintenance,³³ recently proposed as an (augmentation) strategy to improve cognitive depressive symptoms,³⁴ has recently been trialed for the management tinnitus with some success.³⁵ Salicylate amplifies cochlear NMDA-mediated responses but has little or no effect on α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic (AMPA) glutamate receptor and glutamatergic kainite-mediated responses.³⁶ AMPA receptors (AMPA) are both glutamate receptors and cation channels that are integral to plasticity and synaptic transmission at many postsynaptic membranes. AMPA receptors mediate the excitotoxic effect of excessive noise trauma, leading to excessive amounts of glutamate in the synaptic cleft that results in neurite loss,³⁷ and are also able

to upregulate themselves through plastic changes.³⁸ Application of NMDA antagonists onto the round window membrane abolishes tinnitus, even in animals receiving treatment with the 5-HT_{2C} anxiogenic agent 1-(3-chlorophenyl) piperazine (mCPP),³⁹ suggesting NMDA blocking, even for depressed or chronically anxious tinnitus patients.⁴⁰

Ion channel and others

The inner cells of the cochlea have only the L-type of calcium (Ca²⁺) channels, that are involved in the regulation of *in situ* glutamate release and are blocked by salicylate.⁴¹ In the IC, the current through the L-type channels, although unable to directly trigger the neurotransmitter release, contributes to GABAergic transmission by activating the second messenger system and/or by increasing the intracellular Ca²⁺ connection;⁴² while salicylate also blocks the outward and delayed rectifier potassium (K⁺) channels in rat IC, hence decreasing the GABAergic transmission.⁴³

Glycine receptors (whose number significantly decreases with age in DCN) along with acetylcholine muscarinic and nicotinic receptors (whose number also decreases with age in the VCN and IC) antagonism has also been investigated for their implication in the genesis of tinnitus^{44,45} and may lead to new pharmacological strategies (such as with varenicline, a smoking-cessation drug and often prescribed to psychiatric patients⁴⁶) in the treatment of tinnitus.

Psychopharmacological treatment

Many drugs, including psychopharmacological agents, are routinely employed in the treatment of tinnitus (Table 2). Such drugs could be prescribed as both a potential therapy for tinnitus, or just to patients with psychopathological problems who may also present with tinnitus, or to patients who may develop tinnitus (possibly as iatrogenic phenomenon).

Antidepressants

Baldo and colleagues performed a meta-analysis in 2006 on tricyclic antidepressants (TCAs) and SSRIs, which questioned substantially their efficacy in the treatment of tinnitus, although most evidence may have been invalidated by methodological bias or the reliance on single case reports.⁴⁷ In 2007, an analysis of four RCTs by Robinson found that higher doses of SSRIs and TCAs appear to work for tinnitus patients who also exhibit depression and anxiety⁴⁸ or insomnia,⁴⁹ both of which are frequent comorbidities among chronic otolaryngic patients.⁵⁰ What appears to be irreversible disability of otologic origin may, in part, be a reversible disability of

Table 2 Main pharmacological agents for subjective tinnitus

Drugs	Rationale	Evidence	
Antidepressants			
TCAs	NA- and 5HT-ergic transmission modulation, anticholinergic action, BDNF synthesis enhancement	Nortriptyline	Superior to placebo (RTC) (higher doses of TCAs and SSRIs appear to work for tinnitus patients who also exhibit depression and anxiety or insomnia)
		Amitriptyline	Some success (case reports and a single blind study). Low doses sometimes related to tinnitus onset
		Trimipramine Clomipramine	Less effective than placebo (RTC) High doses associated with tinnitus onset
SSRIs	5HT-ergic transmission modulation, BDNF synthesis enhancement	Paroxetine	No significant improvement for nondepressed patients (RTC)
SNRIs	NA- and 5HT-ergic transmission modulation, BDNF synthesis enhancement	Sertraline	More effective than placebo for severe refractory tinnitus cases (RTC)
		Fluoxetine Duloxetine	Worsening of most of the tinnitus cases Withdrawal has been seldom associated with tinnitus
		Venlafaxine	Some success (case reports); withdrawal has been seldom associated with tinnitus
NDRIs	NA- and DA-ergic transmission modulation, BDNF synthesis enhancement	Bupropion	Limited-sample reports indicated its potential induction of tinnitus
Heterocyclics		Mianserin Trazodone	Lack of evidence Lack of evidence
Sedative-hypnotics			
Benzodiazepines	GABA-A-ergic transmission modulation (alprazolam, clonazepam, midazolam: short half/average-life and pro 5-HT-ergic features)	Alprazolam	Superior to placebo in the treatment of severe disabling tinnitus of predominantly cochlear origin (effective especially for those patients presenting high levels of co-morbid anxiety)
		Clonazepam	Superior to placebo (effective also for pulsatile tinnitus when associated with beta-blocker agents)
		Midazolam	Superior to placebo
GABA-B agonists	Muscle relaxing and antispastic action	Baclofen	Effective for pulsatile tinnitus
Mood stabilizers			
Antiepileptics	GABA-ergic transmission enhancement, GABA transaminase inhibition, alpha (2) delta calcium channel antagonism, others (histone deacetylases role, GSK-3 remodeling and synapsin I clustering)	Carbamazepine	Some success in case reports and trials (limited sampled reports indicated its potential induction of objective tinnitus)
		Gabapentin	Effective in reducing subjective or objective tinnitus, especially in individuals with associated acoustic trauma (RTC)
		Lamotrigine	Ineffective (RTC)
		Valproate	Effective in a single case study (while another single-case evidence reported its potential role in inducing tinnitus)
		Vigabatrin Pregabalin	Animal studies only No systematic evidence
Lithium	Inhibitory effect on the phosphoinositide cascade, proserotonergic at higher doses? Others	Not systematically assessed for tinnitus	
Glutamatergic compounds			
Acamprosate	NDMA receptor blockade and GABA-ergic transmission enhancing	Superior than placebo in a double-blind study (potentially effective even for depressed or chronically anxious tinnitus patients)	
Memantine	Hair cells NMDA transmission blockade and cholinergic modulation	Not yet investigated in humans for tinnitus treatment	

(Continued)

Table 2 (Continued)

Drugs	Rationale	Evidence
Antipsychotics		
Typical	Auditolimbic D ₂ DA-ergic modulation	Cases of schizophrenic patients treated with antipsychotics for tinnitus-like sensations
Atypical	Dose-dependant D ₂ receptor antagonism (postsynaptic D ₂ -blockade at high doses and presynaptic auto-inhibitory D ₂ -blockade at low doses with DA release instead of reduction), D ₂ partial agonism, others	Sulpiride Superior to placebo in a single blind (low doses) placebo-controlled study and in a RTC + hydroxyzine or melatonin
Lidocain	L-type CA ⁺⁺ channels blockade, indirectly contributing to glutamatergic, GABA-ergic and cholinergic transmission modulation	Partially effective
Melatonin	Pineal neurohormone with melatonergic agonism and anti-DA-ergic actions	Partially effective, especially in patients with sleep disturbance
Botulinum toxin	Inhibition of acetylcholine release at the neuromuscular junction	Partially effective in somatic tinnitus
Zinc, antioxidants, minerals, vitamins, ginko biloba and other herbal remedies	Different mechanisms of action	Alternate results or ineffective
Unassessed potential targets		
Agomelatine	MT1/MT2 melatonergic agonism and 5-HT _{2C} -ergic antagonism	No evidence to date
Varenicline	Nicotinic acetylcholine receptor partial agonism	No evidence to date

Abbreviations: DA, dopamine; TCA, tricyclic antidepressants; CA, calcium; GABA gamma-aminobutyric acid; SSRIs, selective serotonin reuptake inhibitors; SNRIs, serotonin norepinephrine reuptake inhibitors; NDRI, norepinephrine and dopamine reuptake inhibitors; NDMA, N-methyl D-aspartate; RTC, relaxation training control; BDNF, brain-derived neurotrophic factor; NA, noradrenaline; GSK-3, glycogen synthase kinase-3; MT, metallothionein.

psychiatric origin when treated^{51,52} with drugs proven to be effective antidepressants such as nortriptyline.⁵³

Some TCAs have proven not to be effective for tinnitus,^{54,55} or were even less effective than placebo⁵⁶ (or possibly being associated with tinnitus onset with high doses of clomipramine).⁵⁷

A possible dose effect may exist as lower doses have sometimes been related to tinnitus onset,^{58,59} even when the same agents were reported as being effective⁶⁰ (as in non-TCA heterocyclic antidepressants such as trazodone⁶¹ and mianserin).⁶²

Sertraline was found to be more effective than placebo for severe refractory tinnitus cases⁶³ although its discontinuation was associated with tinnitus onset.⁶⁴ Other SSRIs (eg, paroxetine) provided no significant improvement for nondepressed patients⁶⁵ or even exacerbated tinnitus (eg, fluoxetine).⁶⁶

While newer antidepressants have not been investigated as thoroughly as SSRIs and TCAs, little data on their role in the treatment of tinnitus has been reported and which has sometimes led to inhomogeneous evidence.

Serotonin norepinephrine reuptake inhibitors (SNRIs) withdrawal has seldom been associated with tinnitus (both for venlafaxine^{67,68} and duloxetine⁶⁹), while the norepinephrine dopamine reuptake inhibitor (NDRI) bupropion may mimic

a transient ischemic attack that may induce tinnitus,^{70,71} that is most likely due to DA-ergic enhancement.

Antipsychotics

Although pharmacological treatment is well-established for tinnitus in clinical practice systematic, rigorous literature evidence on antipsychotics is lacking, especially with regard to recently introduced atypical compounds.

Also, it is sometimes difficult to separate tinnitus from psychotic symptoms, thus increasing the cases of schizophrenic patients who may receive antipsychotics for tinnitus-like sensations or hallucinations.⁷²

To date the most accurately investigated antipsychotic medication for tinnitus is sulpiride.

A 2007 prospective, randomized, single blinded, placebo-controlled study by Lopez-Gonzalez on sulpiride plus hydroxyzine (an antihistamine derivate and subcortical sedative attenuating the limbic system) indicated the efficacy of the antipsychotic drug in reducing the sensation of tinnitus, possibly mediated by the auditolimbic D₂ DA-ergic modulation.⁷³ An RCT investigation from the same group also reported the efficacy of sulpiride in association with melatonin (a pineal hormone with anti-DA-ergic actions) in attenuating tinnitus sensation.⁷⁴ Remarkably, both studies used

low doses of sulpiride (up to 50 mg/8 hours) and this should be taken into account when using (atypical) antipsychotics, since the presynaptic autoinhibitory D₂-blockade should lead to DA release instead of reduction (as evidenced by high dose-mediated postsynaptic D₂ heteroreceptors blockade).

The melatonergic role has also been considered for associations with antipsychotics other than sulpiride,⁷⁵ and is worthy of further investigation, even for other classes of drugs (ie, the newly introduced antidepressant agomelatine, acting as metallothionin-1(MT1)/MT2 melatonergic agonist and 5-HT_{2C}-ergic antagonist).⁷⁶

Mood stabilizers

Many antiepileptics used as mood stabilizers have been considered for the treatment of tinnitus, although almost all available evidence does not come from systematic investigations and therefore do not lead to an univocal consensus.

Among others, carbamazepine has been proposed both as stand-alone treatment for subjective tinnitus⁷⁷ or with lidocaine,⁷⁸ salicylate,⁷⁹ or steroid⁸⁰ augmentation/switch option, while limited-sampled reports indicate its potential induction of objective tinnitus (by auditory nerve vascular compression).^{81,82}

Valproate, another popular antiepileptic/mood-stabilizer, has also been investigated as a candidate for tinnitus therapy.^{83,84} However a single case has been reported suggesting its potential role in inducing tinnitus (eventually by magnesium metabolism impairment),^{85,86} with tinnitus eventually being misinterpreted as psychotic symptom.⁸⁷ The only RCT evidence for lamotrigine concluded it to be ineffective for tinnitus.⁸⁸

To date, gabapentin is the most thoroughly investigated antiepileptic drug for tinnitus. A placebo-controlled trial by Bauer and colleagues reported its effectiveness in reducing subjective or objective tinnitus, especially in individuals with associated acoustic trauma.⁸⁹ However, more recent, placebo-controlled studies using high doses of gabapentin (1800 mg/day and 900–3600 mg/day respectively) have disproved its efficacy versus placebo.^{90–92} Remarkably, no systematic studies on the use of pregabalin (a gabapentin-related GABA-modulator plus alpha (2) delta calcium channel antagonism) have been reported to date. However, evidence from the literature on tinnitus relief by CA²⁺ antagonists⁹³ suggests that an investigation into applicability in the treatment of tinnitus is warranted.⁴¹

Lithium, a proven nonantiepileptic mood-stabilizer, has not been systematically assessed for its suitability for the treatment of tinnitus, although its inhibitory effect on the phosphoinositide cascade, maximized in case of

over-the-counter assumption, may lead to vasodilatory shock as well as to hearing deficiency.⁹⁴

Finally, while GABA transaminase inhibition is considered a basic antiepileptic feature, antimanic actions seem to be independent of it, with recent acquisition focusing on the role of histone deacetylases⁹⁵ and the glycogen synthase kinase-3 remodeling and synapsin I clustering.⁹⁶ This consideration contributes to explain why the mood stabilizers class includes so much different compounds, whereas the GABA modulation is just part of the therapeutic portfolio, thus making particularly difficult to investigate a single action hypothesized to modulate tinnitus perception.

Sedative-hypnotics

Benzodiazepines (BDZs) and other sedative-hypnotics agents are among the most prescribed drugs worldwide and the pivotal role of BDZ receptor distribution in severe, intractable tinnitus has been studied by Daftary and colleagues.⁹⁷ It is therefore unsurprising that BDZ GABA-A-ergic modulation has repeatedly been considered for treating severe disabling tinnitus of predominantly cochlear origin.⁹⁸

Although systematic investigations are lacking, the majority of evidence suggests their potential role in tinnitus management, especially for those patients presenting with high levels of comorbid anxiety. Placebo-controlled trials for alprazolam,⁹⁹ midazolam,¹⁰⁰ and clonazepam¹⁰¹ have produced the most promising results (remarkably, these drugs share the short half/average-life and the pro-5-HT-ergic features). Clonazepam has also been reported as being effective for the treatment of pulsatile tinnitus when associated with beta-blocker agents,¹⁰² although less effective than baclofen (a muscle relaxer and an antispasmodic GABA-B-agonist agent).¹⁰³ While no systematic trials have been conducted on the long half-life BDZs, it has been observed that the chance of protracted tinnitus due to abrupt discontinuation of diazepam (30 mg/day long-term treatment)¹⁰⁴ or oxazepam¹⁰⁵ is most likely due to a receptor upregulation phenomena.

Conclusions

Tinnitus pharmacological therapies include a wide range of proposed medicaments. Among others, zinc,¹⁰⁶ melatonin,⁷⁵ lidocaine,¹⁰⁷ botulinum toxin,¹⁰⁸ antioxidants minerals, vitamins, ginkgo biloba, and other herbal remedies,¹⁰⁹ have been considered in the treatment of tinnitus.

Indeed, psychological (eg, cognitive behavioral therapy) and retraining approaches have been suggested repeatedly as preferred integrative or stand-alone treatments.¹¹⁰ However,

psychopharmacological agents are often considered choices as well, although it is difficult to assess their efficacy.

In fact, major limits influence the validity of available literature evidence. Few studies follow RCT standards and most reports rely on small-sized samples or are just single-case reports, with the literature evidence considered by this review relying on MEDLINE.

Also clinical diagnosis of tinnitus are extremely heterogeneous and could be “influenced” by concomitant comorbidities/therapies, especially in case of anxious and depressed patients. These considerations inevitably limit the comprehension of the underpinning neurobiological substrates of tinnitus, further reducing the chance of developing selective pharmacological agents.

Nonetheless, while psychopharmacological agents include a heterogeneous, broad number of compounds, the rational acknowledgment of studies in the current literature and hypothesized biological causes should be considered mandatory in order to avoid iatrogenic phenomena or to effectively consider the psychopharmacological treatment strategy.

Acknowledgments

The authors acknowledge Mrs Rita Santi Amantini for her secretarial services.

Disclosures

The authors received no financial support for this work and report no conflict of interest relevant to this work.

References

- Sequist LV. Protein degradation/chaperone system and histone deacetylase pathway. *J Thorac Oncol*. 2009;4(11 Suppl 3):S1077–S1078.
- World Health Organization. *International Statistical Classification of Diseases and Related Health Problems; Tenth edition (ICD-10)*. Geneva, Switzerland: World Health Organization; 1992.
- Demeester K, van Wieringen A, Hendrickx JJ, et al. Prevalence of tinnitus and audiometric shape. *B-ENT*. 2007;3(Suppl 7):37–49.
- Adoga AA, Adoga AS, Obindo JT. Tinnitus and the prevalence of co-morbid psychological stress. *Niger J Med*. 2008;17(1):95–97.
- Sie KC, Rubel EW. Rapid changes in protein synthesis and cell size in the cochlear nucleus following eighth nerve activity blockade or cochlea ablation. *J Comp Neurol*. 1992;320(4):501–508.
- Tzounopoulos T. Mechanisms of synaptic plasticity in the dorsal cochlear nucleus:plasticity-induced changes that could underlie tinnitus. *Am J Audiol*. 2008;17(2):S170–S175.
- Han BI, Lee HW, Kim TY, Lim JS, Shin KS. Tinnitus:characteristics causes mechanisms and treatments. *J Clin Neurol*. 2009;5(1):11–19.
- Kemp DT. Stimulated acoustic emissions from within the human auditory system. *J Acoust Soc Am*. 1978;64:1386–1391.
- Kiang NY, Moxon EC, Levine RA. Auditory-nerve activity in cats with normal and abnormal cochleas. In: *Sensorineural Hearing Loss*. Chichester, West Sussex: Wiley, Ciba Found Symp; 1970: 241–273.
- Jastreboff PJ, Hazell JW. A neurophysiological approach to tinnitus:clinical implications. *Br J Audiol*. 1993;27:7–17.
- Kaltenbach JA, Zhang J, Finlayson P. Tinnitus as a plastic phenomenon and its possible neural underpinnings in the dorsal cochlear nucleus. *Hear Res*. 2005;206:200–226.
- Salvi RJ, Wang J, Ding D. Auditory plasticity and hyperactivity following cochlear damage. *Hear Res*. 2000;147:261–274.
- Møller AR. Pathophysiology of tinnitus. *Ann Otol Rhinol Laryngol*. 1984;93:39–44.
- Levine RA. Somatic (craniocervical) tinnitus and the dorsal cochlear nucleus hypothesis. *Am J Otolaryngol*. 1999;20:351–362.
- Ling LL, Hughes LF, Caspary DM. Age-related loss of the GABA synthetic enzyme glutamic acid decarboxylase in rat primary auditory cortex. *Neuroscience*. 2005;132(4):1103–1113.
- Campos ML, de Cabo C, Wisden W, Juiz JM, Merlo D. Expression of GABA(A) receptor subunits in rat brainstem auditory pathways:cochlear nuclei superior olivary complex and nucleus of the lateral lemniscus. *Neuroscience*. 2001;102(3):625–638.
- Rosenhall U, Karlsson AK. Tinnitus in old age. *Scand Audiol*. 1991;20(3):165–171.
- Diaz SL, Barros VG, Antonelli MC, Rubio MC, Balerio GN. Morphine withdrawal syndrome and its prevention with baclofen: Autoradiographic study of mu-opioid receptors in prepubertal male and female mice. *Synapse*. 2006;60(2):132–140.
- Brozoski TJ, Spires TJ, Bauer CA. Vigabatrin a GABA transaminase inhibitor reversibly eliminates tinnitus in an animal model. *J Assoc Res Otolaryngol*. 2007;8(1):105–118.
- Shulman A. Gabapentin and tinnitus relief. *Int Tinnitus J*. 2008;14(1):1–5.
- Yang G, Lobarinas E, Zhang L, et al. Salicylate-induced tinnitus:behavioral measures and neural activity in auditory cortex of awake rats. *Hear Res*. 2007;226(1–2):244–253.
- Tyler RS, Coelho C, Noble W. Tinnitus: standard of care, personality differences, genetic factors. *ORL J Otorhinolaryngol Relat Spec*. 2006;68(1):14–19; discussion 20–2.
- Folmer RL, Shi YB. SSRI use by tinnitus patients:interactions between depression and tinnitus severity. *Ear Nose Throat J*. 2004;83(2):107–112.
- Lockwood AH, Salvi RJ, Coad ML, Towsley ML, Wack DS, Murphy BW. The functional neuroanatomy of tinnitus:evidence for limbic system links and neural plasticity. *Neurology*. 1998;50(1):114–120.
- Brunoni AR, Lopes M, Fregni F. A systematic review and meta-analysis of clinical studies on major depression and BDNF levels:implications for the role of neuroplasticity in depression. *Int J Neuropsychopharmacol*. 2008;11(8):1169–1180.
- Tan J, Rüttiger L, Panford-Walsh R, et al. Tinnitus behavior and hearing function correlate with the reciprocal expression patterns of BDNF and Arg3.1/arc in auditory neurons following acoustic trauma. *Neuroscience*. 2007;145(2):715–726.
- Moller AR. Tinnitus and pain. *Prog Brain Res*. 2007;166:47–53.
- Puel JL. Cochlear NMDA receptor blockade prevents salicylate-induced tinnitus. *B-ENT*. 2007;3(Suppl 7):19–22.
- Guitton MJ, Caston J, Ruel J, Johnson RM, Pujol R, Puel JL. Salicylate induces tinnitus through activation of cochlear NMDA receptors. *J Neurosci*. 2003;23(9):3944–3952.
- Guitton MJ, Dudai Y. Blockade of cochlear NMDA receptors prevents long-term tinnitus during a brief consolidation window after acoustic trauma. *Neural Plast*. 2007;2007:80904.
- Pomara N, Ott BR, Peskind E, Resnick EM. Memantine treatment of cognitive symptoms in mild to moderate Alzheimer disease:secondary analyses from a placebo-controlled randomized trial. *Alzheimer Dis Assoc Disord*. 2007;21(1):60–64.
- Oliver D, Ludwig J, Reisinger E, Zoellner W, Ruppertsberg JP, Fakler B. Memantine inhibits efferent cholinergic transmission in the cochlea by blocking nicotinic acetylcholine receptors of outer hair cells. *Mol Pharmacol*. 2001;60(1):183–189.
- Mason BJ. Treatment of alcohol-dependent outpatients with acamprosate:a clinical review. *J Clin Psychiatry*. 2001;62(Suppl 20):42–48.

34. Mann K, Kiefer F, Spanagel R, Littleton J. Acamprosate: Recent findings and future research directions. *Alcohol Clin Exp Res*. 2008;32(7):1105–1110.
35. Azevedo AA, Figueiredo RR. Tinnitus treatment with acamprosate: double-blind study. *Braz J Otorhinolaryngol*. 2005;71(5):618–623.
36. Peng BG, Chen S, Lin X. Aspirin selectively augmented N-methyl-D-aspartate types of glutamate responses in cultured spiral ganglion neurons of mice. *Neurosci Lett*. 2003;343(1):21–24.
37. Puel JL, Ruel J, Gervais d'Aldin C, Pujol R. Excitotoxicity and repair of cochlear synapses after noise-trauma induced hearing loss. *Neuroreport*. 1998;9(9):2109–2114.
38. Muly SM, Gross JS, Potashner SJ. Noise trauma alters D-[3H]aspartate release and AMPA binding in chinchilla cochlear nucleus. *J Neurosci Res*. 2004;75(4):585–596.
39. Guitton MJ, Pujol R, Puel JL. m-Chlorophenylpiperazine exacerbates perception of salicylate-induced tinnitus in rats. *Eur J Neurosci*. 2005;22(10):2675–2678.
40. Weise C, Heinecke K, Rief W. Stability of physiological variables in chronic tinnitus sufferers. *Appl Psychophysiol Biofeedback*. 2008;33(3):149–159.
41. Sziklai I. The significance of the calcium signal in the outer hair cells and its possible role in tinnitus of cochlear origin. *Eur Arch Otorhinolaryngol*. 2004;261(10):517–525.
42. Narváez M, Figueras A, Capellà D, Laporte JR. Tinnitus with calcium-channel blockers. *Lancet*. 1994;343(8907):1229–1230.
43. Puel JL, Guitton MJ. Salicylate-induced tinnitus: molecular mechanisms and modulation by anxiety. *Prog Brain Res*. 2007;166:141–146.
44. Eggermont JJ. Tinnitus: neurobiological substrates. *Drug Discov Today*. 2005;10(19):1283–1290.
45. Elgoyhen AB, Katz E, Fuchs PA. The nicotinic receptor of cochlear hair cells: a possible pharmacotherapeutic target? *Biochem Pharmacol*. 2009;78(7):712–719.
46. Jimenez-Ruiz C, Berlin I, Hering T. Varenicline: a novel pharmacotherapy for smoking cessation. *Drugs*. 2009;69(10):1319–1338.
47. Baldo P, Doree C, Lazzarini R, Molin P, McFerran DJ. Antidepressants for patients with tinnitus. *Cochrane Database Syst Rev*. 2006;(4):CD003853.
48. Robinson S. Antidepressants for treatment of tinnitus. *Prog Brain Res*. 2007;166:263–271.
49. Dobie RA. Depression and tinnitus. *Otolaryngol Clin North Am*. 2003;36(2):383–388.
50. Chandra RK, Epstein VA, Fishman A.J. Prevalence of depression and antidepressant use in an otolaryngology patient population. *Otolaryngol Head Neck Surg*. 2009;141(1):136–138.
51. Sullivan M, Katon W, Russo J, Dobie R, Sakai C. A randomized trial of nortriptyline for severe chronic tinnitus. Effects on depression disability and tinnitus symptoms. *Arch Intern Med*. 1993;153(19):2251–2259.
52. Dobie RA, Sakai CS, Sullivan MD, Katon WJ, Russo J. Antidepressant treatment of tinnitus patients: report of a randomized clinical trial and clinical prediction of benefit. *Am J Otol*. 1993;14(1):18–23.
53. Dobie RA, Sakai CS, Sullivan MD, Katon WJ, Russo J. Antidepressant treatment of tinnitus patients. Interim report of a randomized clinical trial. *Acta Otolaryngol*. 1992;112(2):242–247.
54. Evans DL, Golden RN. Protriptyline and tinnitus. *J Clin Psychopharmacol*. 1981;1(6):404–406.
55. Golden RN, Evans DL, Nau CH. Doxepin and tinnitus. *South Med J*. 1983;76(9):1204–1205.
56. Mihail RC, Crowley JM, Walden BE, Fishburne J, Reinwall JE, Zajtchuk JT. The tricyclic trimipramine in the treatment of subjective tinnitus. *Ann Otol Rhinol Laryngol*. 1988;97(2 Pt 1):120–123.
57. Ackerman DL, Greenland S, Bystritsky A, Katz RJ. Relationship between early side effects and therapeutic effects of clomipramine therapy in obsessive-compulsive disorder. *J Clin Psychopharmacol*. 1996;16(4):324–328.
58. Mendis D, Johnston M. An unusual case of prolonged tinnitus following low-dose amitriptyline. *J Psychopharmacol*. 2008;22(5):574–575.
59. Podoshin L, Ben-David Y, Fradis M, Malatskey S, Hafner H. Idiopathic subjective tinnitus treated by amitriptyline hydrochloride/biofeedback. *Int Tinnitus J*. 1995;1(1):54–60.
60. Bayar N, Böke B, Turan E, Belgin E. Efficacy of amitriptyline in the treatment of subjective tinnitus. *J Otolaryngol*. 2001;30(5):300–303.
61. Dib GC, Kasse CA, Alves de Andrade T, Gurgel Testa JR, Cruz OL. Tinnitus treatment with trazodone. *Braz J Otorhinolaryngol*. 2007;73(3):390–397.
62. Marais J. Cochleotoxicity due to mianserin hydrochloride. *J Laryngol Otol*. 1991;105(6):475–476.
63. Zoger S, Svedlund J, Holgers KM. The effects of sertraline on severe tinnitus suffering—a randomized double-blind placebo-controlled study. *J Clin Psychopharmacol*. 2006;26(1):32–39.
64. Leiter FL, Nierenberg AA, Sanders KM, Stern TA. Discontinuation reactions following sertraline. *Biol Psychiatry*. 1995;38(10):694–695.
65. Robinson SK, Viirre ES, Bailey KA, Gerke MA, Harris JP, Stein MB. Randomized placebo-controlled trial of a selective serotonin reuptake inhibitor in the treatment of nondepressed tinnitus subjects. *Psychosom Med*. 2005;67(6):981–988.
66. Shemen L. Fluoxetine for treatment of tinnitus. *Otolaryngol Head Neck Surg*. 1998;118(3 Pt 1):421.
67. Pondrom MF, Brahm NC. Venlafaxine-associated tinnitus. *Am J Health Syst Pharm*. 2009;66(18):1606–1608.
68. Ahmad S. Venlafaxine and severe tinnitus. *Am Fam Physician*. 1995;51(8):1830.
69. De Marinis M, Santilli V. Tinnitus in postherpetic neuralgia. *J Headache Pain*. 2010;11(1):83–84.
70. Humma LM, Swims MP. Bupropion mimics a transient ischemic attack. *Ann Pharmacother*. 1999;33(3):305–307.
71. Settle EC. Tinnitus related to bupropion treatment. *J Clin Psychiatry*. 1991;52(8):352.
72. D'Amelio R, Delb W. Comorbidity of schizophrenic psychosis and tinnitus. A hitherto neglected theme in research and therapy. *HNO*. 2008;56(7):670–672.
73. Lopez-Gonzalez MA, Moliner-Peiro F, Alfaro-Garcia J, Esteban-Ortega F. Sulpiride plus hydroxyzine decrease tinnitus perception. *Auris Nasus Larynx*. 2007;34(1):23–27.
74. Lopez-Gonzalez MA, Santiago AM, Esteban-Ortega F. Sulpiride and melatonin decrease tinnitus perception modulating the audiolimbic dopaminergic pathway. *J Otolaryngol*. 2007;36(4):213–219.
75. Neri G, Baffa C, De Stefano A, et al. Management of tinnitus: oral treatment with melatonin and sulodexide. *J Biol Regul Homeost Agents*. 2009;23(2):103–110.
76. Owen RT. Agomelatine: A novel pharmacological approach to treating depression. *Drugs Today (Barc)*. 2009;45(8):599–608.
77. Hulshof JH, Vermeij P. The value of carbamazepine in the treatment of tinnitus. *ORL J Otorhinolaryngol Relat Spec*. 1985;47(5):262–266.
78. Sanchez TG, Balbani AP, Bittar RS, Bento RF, Câmara J. Lidocaine test in patients with tinnitus: rationale of accomplishment and relation to the treatment with carbamazepine. *Auris Nasus Larynx*. 1999;26(4):411–417.
79. Zheng Y, Hooton K, Smith PF, Darlington CL. Carbamazepine reduces the behavioural manifestations of tinnitus following salicylate treatment in rats. *Acta Otolaryngol*. 2008;128(1):48–52.
80. She W, Dai Y, Du X, Chen F, Ding X, Cui X. Treatment of subjective tinnitus: a comparative clinical study of intratympanic steroid injection vs. oral carbamazepine. *Med Sci Monit*. 2009;15(6):P135–P139.
81. Levine RA. Typewriter tinnitus: a carbamazepine-responsive syndrome related to auditory nerve vascular compression. *ORL J Otorhinolaryngol Relat Spec*. 2006;68(1):43–47.
82. Brantberg K. Paroxysmal staccato tinnitus: a carbamazepine responsive hyperactivity dysfunction symptom of the eighth cranial nerve. *J Neurol Neurosurg Psychiatry*. 2010;81(4):451–455.
83. Mansbach AL, Freyens P. Tinnitus: current data and treatment with sodium valproate. *Acta Otorhinolaryngol Belg*. 1983;37(5):697–705.

84. Menkes DB, Larson PM. Sodium valproate for tinnitus. *J Neurol Neurosurg Psychiatry*. 1998;65(5):803.
85. Hori A, Kataoka S, Sakai K, et al. Valproic acid-induced hearing loss and tinnitus. *Intern Med*. 2003;42(11):1153–1154.
86. Yildirim C, Yağiz R, Uzun C, Taş A, Bulut E, Karasalihoğlu A. The protective effect of oral magnesium supplement on noise-induced hearing loss. *Kulak Burun Bogaz Ihtis Derg*. 2006;16(1):29–36.
87. Reeves RR, Mustain DW, Pendarvis JE. Valproate-induced tinnitus misinterpreted as psychotic symptoms. *South Med J*. 2000;93(10):1030–1031.
88. Simpson JJ, Gilbert AM, Weiner GM, Davies WE. The assessment of lamotrigine an antiepileptic drug in the treatment of tinnitus. *Am J Otol*. 1999;20(5):627–631.
89. Bauer CA, Brozoski TJ. Effect of gabapentin on the sensation and impact of tinnitus. *Laryngoscope*. 2006;116(5):675–681.
90. Witsell DL, Hannley MT, Stinnet S, Tucci DL. Treatment of tinnitus with gabapentin: a pilot study. *Otol Neurotol*. 2007;28(1):11–15.
91. Piccirillo JF, Finnell J, Vlahiotis A, Chole RA, Spitznagel E Jr. Relief of idiopathic subjective tinnitus: is gabapentin effective? *Arch Otolaryngol Head Neck Surg*. 2007;133(4):390–397.
92. Bakhshae M, Ghasemi M, Azarpazhooh M, et al. Gabapentin effectiveness on the sensation of subjective idiopathic tinnitus: a pilot study. *Eur Arch Otorhinolaryngol*. 2008;265(5):525–530.
93. Davies E, Knox E, Donaldson I. The usefulness of nimodipine an L-calcium channel antagonist in the treatment of tinnitus. *Br J Audiol*. 1994;28(3):125–129.
94. Plenge U, Moller AM. Development of sustained vasodilatory shock and permanent loss of hearing after severe lithium carbonate poisoning. *Ugeskr Laeger*. 2008;170(5):354.
95. Hobara T, Uchida S, Otsuki K, et al. Altered gene expression of histone deacetylases in mood disorder patients. *J Psychiatr Res*. 2010;44(5):263–270.
96. Hall AC, Brennan A, Goold RG, et al. Valproate regulates GSK-3-mediated axonal remodeling and synapsin I clustering in developing neurons. *Mol Cell Neurosci*. 2002;20(2):257–270.
97. Daftary A, Shulman A, Strashun AM, Gottschalk C, Zoghbi SS, Seibyl JP. Benzodiazepine receptor distribution in severe intractable tinnitus. *Int Tinnitus J*. 2004;10(1):17–23.
98. Bahmad FM, Venosa AR, Oliveira CA. Benzodiazepines and GABAergic in treating severe disabling tinnitus of predominantly cochlear origin. *Int Tinnitus J*. 2006;12(2):140–144.
99. Jalali MM, Kousha A, Naghavi SE, Soleimani R, Banan R. The effects of alprazolam on tinnitus: a cross-over randomized clinical trial. *Med Sci Monit*. 2009;15(11):PI55–PI60.
100. Panford-Walsh R, Singer W, Rüttiger L, et al. Midazolam reverses salicylate-induced changes in brain-derived neurotrophic factor and arg3.1 expression: implications for tinnitus perception and auditory plasticity. *Mol Pharmacol*. 2008;74(3):595–604.
101. Ganança MM, Caovilla HH, Ganança FF, et al. Clonazepam in the pharmacological treatment of vertigo and tinnitus. *Int Tinnitus J*. 2002;8(1):50–53.
102. Albertino S, Assuncao AR, Souza JA. Pulsatile tinnitus: treatment with clonazepam and propranolol. *Braz J Otorhinolaryngol*. 2005;71(1):111–113.
103. Szczepaniak WS, Moller AR. Effects of (-)-baclofen clonazepam and diazepam on tone exposure-induced hyperexcitability of the inferior colliculus in the rat: possible therapeutic implications for pharmacological management of tinnitus and hyperacusis. *Hear Res*. 1996;97(1–2):46–53.
104. Busto U, Fornazzari L, Naranjo CA. Protracted tinnitus after discontinuation of long-term therapeutic use of benzodiazepines. *J Clin Psychopharmacol*. 1988;8(5):359–362.
105. Busto U, Sellers EM, Naranjo CA, Cappell H, Sanchez-Craig M, Sykora K. Withdrawal reaction after long-term therapeutic use of benzodiazepines. *N Engl J Med*. 1986;315(14):854–859.
106. Coelho CB, Tyler R, Hansen M. Zinc as a possible treatment for tinnitus. *Prog Brain Res*. 2007;166:279–285.
107. Ueda S. Treatment of tinnitus with intravenous lidocaine. *Nippon Jibiinkoka Gakkai Kaiho*. 1992;95(9):1389–1397.
108. Láinez MJ, Piera Botulinum toxin for the treatment of somatic tinnitus. *Prog Brain Res*. 2007;166:335–338.
109. Enrico P, Sirca D, Mereu M. Antioxidants minerals vitamins and herbal remedies in tinnitus therapy. *Prog Brain Res*. 2007;166:323–330.
110. Delb W, D'Amelio R, Boisten CJ, Plinkert PK. Evaluation of the tinnitus retraining therapy as combined with a cognitive behavioral group therapy. *HNO*. 2002;50(11):997–1004.

Neuropsychiatric Disease and Treatment

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS, and is the official

Submit your manuscript here: <http://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>

Dovepress

journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.