



ELSEVIER

Contents lists available at ScienceDirect

Marine Environmental Research

journal homepage: www.elsevier.com/locate/marenvres

Comparative sensitivity of *Crassostrea angulata* and *Crassostrea gigas* embryo-larval development to As under varying salinity and temperature

Anthony Moreira^a, Etelvina Figueira^a, Giovanni Libralato^b, Amadeu M.V.M. Soares^a,
Marco Guida^b, Rosa Freitas^{a,*}

^a Department of Biology & CESAM, University of Aveiro, Campus Universitário de Santiago, Aveiro, Portugal

^b Department of Biology, University of Naples Federico II, via Cinthia ed. 7, 80126, Naples, Italy

ARTICLE INFO

Keywords:

Arsenic
Climate change
Development
Embryotoxicity
Oyster
Thermohaline

ABSTRACT

Oysters are a diverse group of marine bivalves that inhabit coastal systems of the world's oceans, providing a variety of ecosystem services, and represent a major socioeconomic resource. However, oyster reefs have become inevitably impacted from habitat destruction, overfishing, pollution and disease outbreaks that have pushed these structures to the break of extinction. In addition, the increased frequency of climate change related events promise to further challenge oyster species survival worldwide.

Oysters' early embryonic development is likely the most vulnerable stage to climate change related stressors (e.g. salinity and temperature shifts) as well as to pollutants (e.g. arsenic), and therefore can represent the most important bottleneck that define populations' survival in a changing environment. In light of this, the present study aimed to assess two important oyster species, *Crassostrea angulata* and *Crassostrea gigas* embryo-larval development, under combinations of salinity (20, 26 and 33), temperature (20, 24 and 28 °C) and arsenic (As) exposure (0, 30, 60, 120, 240, 480, 960 and 1920 µg. As L⁻¹), to infer on different oyster species capacity to cope with these environmental stressors under the eminent threat of climate change and increase of pollution worldwide.

Results showed differences in each species range of salinity and temperature for successful embryonic development. For *C. angulata*, embryo-larval development was successful at a narrower range of both salinity and temperature, compared to *C. gigas*.

Overall, As induced higher toxicity to *C. angulata* embryos, with calculated EC50 values at least an order of magnitude lower than those calculated for *C. gigas*.

The toxicity of As (measured as median effective concentration, EC50) showed to be influenced by both salinity and temperature in both species. Nonetheless, salinity had a greater influence on embryos' sensitivity to As. This pattern was mostly noticed for *C. gigas*, with lower salinity inducing higher sensitivity to As. Results were discussed considering the existing literature and suggest that *C. angulata* populations are likely to become more vulnerable under near future predictions for temperature rise, salinity shifts and pollution.

1. Introduction

For centuries, oyster reefs formed the dominant structural habitat in temperate estuarine systems of the world. Such biogenic features are formed by one or few oyster species that provide the physical structure for entire ecosystems (Beck et al., 2009). Oyster reefs render a variety of ecosystem services, including the formation of habitat structure for commercially important fish species, water quality improvement, shoreline defence against coastal erosion and carbon dioxide storage (Coen et al., 2007; Ridge et al., 2017; Fodrie et al., 2017; Grabowski et al., 2012). Currently, these ecosystems are globally threatened, and

the majority are classified as functionally extinct (Beck et al., 2011).

The main factors affecting the loss of important oyster reefs around the globe include overexploitation and habitat degradation (Cranfield et al., 1999; Leniham and Peterson, 1998), pollution (Ruano, 1997), exotic species introduction (Miossec et al., 2009; Ruesink et al., 2005), and disease outbreaks (Virvilis and Angelidis, 2006; Comps et al., 1988). In addition, climate change related impacts (i.e. temperature rise, seawater acidification and salinity regime alterations) will further challenge oyster species survival worldwide (Brander, 2007; Dekshenieks et al., 2000; Hoehg-Guldberg and Bruno, 2010; Knight, 2017; Levinton et al., 2011). The comprehension of the factors

* Corresponding author. Departamento de Biologia, Universidade de Aveiro Campus, Universitário de Santiago, 3810-193, Aveiro, Portugal.
E-mail address: rosafreitas@ua.pt (R. Freitas).

<https://doi.org/10.1016/j.marenvres.2018.06.003>

Received 2 April 2018; Received in revised form 29 May 2018; Accepted 4 June 2018
0141-1136/ © 2018 Elsevier Ltd. All rights reserved.

influencing oysters' environmental sensitivity can increase the background knowledge and the possibility to better protect and manage this important biological resource.

According to FAO (2015), oysters form an important global aquaculture shellfish resource. Particularly, *Crassostrea gigas* (Thunberg, 1793) a native species to Japan, is nowadays virtually present in coastal systems of the entire planet (Miossec et al., 2009). In Europe, *C. gigas* is cultured in several countries, with France leading *C. gigas* production and consumption statistics (Heral, 1989; Buestel et al., 2009). However, wild populations have become naturalized and established in several European countries, and reports on *C. gigas* natural occurrences stretch from Norway to southern Portugal (Miossec et al., 2009 and references therein).

Another important species contributing for global oyster fishery landings, *Crassostrea angulata* (Lamarck, 1819), is the main species cultured in southern China (Hsiao et al., 2016; Qin et al., 2012), also occurring in southern Europe (Batista et al., 2015) and North Africa (Fabioux et al., 2002). At present, the geographical distribution of *C. angulata* in Europe is limited to the Iberian Peninsula (Buestel et al., 2009; Batista et al., 2015), where the most pristine population naturally occurs in the Sado estuary (Portugal), making it an important biological resource to be protected.

Under the increasing pressure of climate change related phenomena to estuarine systems worldwide, the combined effects of stressors such as temperature rise, salinity alterations and pollution (Harley et al., 2006; Lejeusne et al., 2010; Robins et al., 2016), are likely to increase the selective pressure on oyster species in the environment. Although adult *C. angulata* and *C. gigas* can be fairly resilient to fluctuations of abiotic factors and to pollutants (Cross et al., 2014; Moreira et al., 2016a; b; Zanette et al., 2011), early developmental stages are more susceptible to physico-chemical environmental changes compared to juveniles and adults (Beiras and His, 1994; His et al., 1999; MacInnes and Calabrese, 1979; Ringwood, 1990). Thus, they represent the bottleneck that define oyster species resilience to environmental change (Byrne et al., 2012), and therefore are important to be studied to infer on different species sensitivities in a changing environment.

The aim of the present study was to compare the sensitivities of *C. angulata* and *C. gigas* embryo-larval development, considering various exposure scenarios including changes in arsenic (As) concentration, salinity and temperature. For this, *C. angulata* from the Sado estuary and *C. gigas* from Guernsey Island embryo-larval development were studied under different combinations of salinity and temperature to assess possible impacts of climate driven alterations, and the combined effect of these conditions with As. Arsenic was chosen as reference stressor due to: a) its toxicity to oyster embryos is strongly influenced by salinity and temperature variations (Moreira et al., 2018); b) its ubiquity in estuarine ecosystems (Neff, 2002) and tendency to remobilize from sediment (De Gieter et al., 2005; Ereira et al., 2015; Masson et al., 2007); and c) its propensity to increase in concentration from climatic events (Henke, 2009; Galloway et al., 2018). Arsenic concentrations in estuarine systems are therefore highly dynamic (Henke, 2009). For instance Ereira et al. (2015) reported As concentrations of $5.2 \mu\text{g L}^{-1}$ in seawater, 20 mg kg^{-1} in sediment and $260 \mu\text{g kg}^{-1}$ in suspended particulate matter, evidencing the potential exposure of relatively high As concentrations to resident biota.

Embryotoxicity tests on *C. angulata* and *C. gigas*, were carried out considering a range of As concentrations (30, 60, 120, 240, 480, 960 and $1920 \mu\text{g L}^{-1}$), different salinity (20, 26 and 33) and temperature (20, 24 and 28°C) levels, as well as different time of exposure (24 and 48 h) to investigate: i) the effect of varying salinity and temperature on both species embryo-larval development; ii) As embryotoxicity to both species; and iii) the effects of varying salinity and temperature on embryos sensitivity to As.

2. Methods

2.1. Experimental setup

Experiments were conducted separately for each species, using previously cleaned and sterilised glassware for the entire experimental setup. Analytical grade artificial seawater (Tropic Marine Sea Salt) from the same batch was used for exposure media preparation and spawning, prepared according to the manufacturer's instructions using reverse osmosis (RO) water, 3 days before the experiments took place to achieve a salinity of 33 (i.e reference salinity) (Leverett and Thain, 2013). After complete salt dissolution and equilibration (24 h) seawater was filtered ($0.2 \mu\text{m}$) through cellulose acetate filters (Millipore) using a vacuum filtration unit. Seawater salinity was adjusted to obtain 3 separate batches at three different salinity levels ($20, 26$ and 33 ± 1) for exposure media preparation using RO water as dilution media (measured pH for all batches of seawater used ranged from 8.16 to 8.29 (Hanna-Instruments)). Salinity levels were chosen in accordance to Moreira et al. (2018), that showed As toxicity to *C. gigas* embryos could restrict embryo-larval development within 20–33 of salinity.

A stock solution of sodium arsenate (Na_3AsO_4) (CAS no. 10048-95-0, Sigma-Aldrich, Missouri, USA) was prepared in ultra-pure water and spiked in separate volumetric vessels to achieve nominal As concentrations of 30, 60, 120, 240, 480, 960 and $1920 \mu\text{g L}^{-1}$, for each target salinity. Exposure concentrations were chosen based on previous studies on *C. gigas* embryonic development and As exposure (Martin et al., 1981; Mamindy-Pajany et al., 2010). Exposure solutions were distributed in 24-well sterile capped polystyrene microplates (VWR), giving one microplate per salinity level, 3 wells (3 ml each) per exposure condition (As concentration) including negative controls. Each microplate corresponding to salinity and As conditions, were 3 fold replicated for incubations at different temperatures (20, 24 and 28°C) to test two levels of 4°C increase of temperature within projected global surface temperature rise by the end of the 21st century relative to years 1986–2005 (RCP8.5) (IPCC, 2014), while testing a temperature range that would allow to infer on embryo development under As exposure based on previous data (Moreira et al., 2018). All the above stated microplates were further replicated twice, to test all conditions after two different timeframes of embryonic development, to allow for the assessment of As induced delayed development to oyster embryos (Moreira et al., 2018), while testing valid exposure time criteria of 24 h (His et al., 1999) and 48 h (Knezovich et al., 1981) for *C. gigas*. Each microplate was incubated at the desired temperature overnight in separate climatic chambers ($\pm 1^\circ\text{C}$) before spawning induction took place to stabilize testing media at the target temperature. Copper (Cu (NO_3)₂) was used as reference toxicant (positive control) (Libralato et al., 2009) at 3, 6, 12, 18 and, $30 \mu\text{g L}^{-1}$ of Cu, for which incubations took place at standard salinity 33 and temperature 24°C .

Each stock of seawater used for the embryotoxicity assays (at every combination of salinity and As concentration) were analysed by Inductively Coupled Plasma - Mass Spectrometry (ICP-MS) to determine effective As concentrations for each condition. Multi-Element Standard IV - 71A (Inorganic Ventures) was used as standard for As quantification, and calibration curve verified with standard reference material (NIST SRM 1643f) (Supplementary Table S6).

2.2. Spawning and fertilization

C. gigas progenitors were obtained from Guernsey Sea Farms (UK), and *C. angulata* progenitors were provided by an aquaculture facility operating in the Sado estuary (Exporsado). *C. angulata* were collected during spawning season, from intertidal growing tables, and stimulated to spawn one day after collection. Spawning was induced by thermal stimulation, by consecutively changing oysters from seawater baths set at 18 and 28°C in 30 min consecutive intervals (Libralato et al., 2007). Gamete emission and quality (oocyte shape and sperm motility) were

visually inspected under a microscope. The number of male and female oysters selected for fertilization were 4 and 3 (*C. gigas*) and 6 and 5 (*C. angulata*), respectively.

Selected females were left to spawn in separate beakers, oocyte suspensions were filtered through a 100 μm nylon mesh and mixed in a final 500 ml suspension (salinity 33, 24 °C). Male gametes were collected in separate, filtered through a 45 μm nylon mesh into a mixed suspension, and left to activate for 20 min. Oocyte suspensions were fertilized by adding approximately 1 to 10⁶ oocyte-to-sperm ratios, and fertilization success verified by microscopy.

Zygotes were immediately transferred to microplates containing the exposure media to reach approximately 200 embryos per well. Microplates with the exposure media (0, 30, 60, 120, 240, 480, 960 and 1920 $\mu\text{g L}^{-1}$ of As) at different salinities (20, 26, 33) and fertilized oocytes were left to incubate in the dark, at different temperatures (20, 24 and 28 \pm 1 °C) for 24 and 48 h post fertilization. Embryo-larval development was stopped by adding buffered formalin (4%). Analysis followed visual inspection of embryo-larval development under an inverted microscope and camera (Leica: DMIL-1; MC170 HD) by counting 100 embryos per well, and characterizing the relative frequencies of different types of development (D-shape, pre-D, protruded mantle, kidney shape, indented shell and dead larvae) according to His et al. (1997) (Fig. 1).

The zygote suspensions from each species that remained unused from embryotoxicity assays were analysed for trace elements (As, Cd, Cr, Cu, Hg, Ni, Pb, Sn and Zn), following an adaptation of Weng and Wang (2017) protocol as follows. Zygote suspensions of each species were centrifuged at 3000 g for 20 min (4 °C) followed by pellet collection. The pellets consisting of each species zygotes were freeze dried, and posteriorly digested overnight with analytical grade nitric acid and hydrogen peroxide (30% w/w) (Chem-Lab NV), alongside with TORT-3 (Lobster Hepatopancreas Reference Material for Trace Metals, NRC Canada) for analytical control by elements recovery percentages calculation. The digested samples were analysed for As, Cd, Cr, Cu, Hg, Ni, Sn and Zn by ICP-MS, with IV- 71A (Inorganic Ventures) as standard, and calibration curves verified with standard reference material (NIST SRM 1643f), calculated measure of trueness over 90%. Recovery percentages based on TORT-3 reference values varied between 93.7 and 111.6%.

2.3. Data analysis

Results obtained were analysed as percentages of abnormal larvae (including pre-D and other malformations) according to His et al. (1999), and the validity of the experiments verified by results for negative and positive controls, considering the acceptability ranges proposed by ASTM (2004) (> 70% D-shape larvae) for negative controls, and Libralato et al. (2009) (9.47 $\mu\text{g. L}^{-1}$ \leq EC50 \leq 21.72 $\mu\text{g. L}^{-1}$) for positive controls.

To represent embryo-larval development under different salinity and temperature levels, frequencies of malformed larvae observed at different salinities and temperature (negative controls), after 24 and 48 h development were represented in Contour plots using SigmaPlot version 11.0.

To assess As embryotoxicity, data on malformation frequency from As exposures were corrected for the effects observed in the respective negative control (salinity and temperature) using the Abbott's formula (ASTM, 2004). Data were submitted to non-linear regression analysis using GraphPad Prism version 6.01. Effective As concentrations retrieved from ICP-MS for each condition were used for toxicity calculations, after lognormal transformation. The best fit dose-response curve values allowed to estimate the median effect concentrations (EC50s) and respective 95% confidence intervals for conditions where larvae development was successful. To test the effects of salinity and temperature on embryos sensitivities to As, the following null hypotheses were tested: i) H₀ⁱ: no differences existed among EC50s at different salinity levels considering fixed temperature; ii) H₀ⁱⁱ: no differences existed among EC50s at different temperature levels considering fixed salinity; iii) H₀ⁱⁱⁱ: no differences existed between EC50s at 24 and 48 h for each combination of salinity and temperature. To test for differences among EC50 values between conditions, analysis of variance was performed using one-way ANOVA, followed by Tukey's multiple comparisons tests, based on values of Log(EC50) and the associated standard error returned by the software for each estimated curve (GraphPad Prism version 6.01). EC50 values are provided for each species in separate graphs showing the relative 95% confidence intervals, and significant differences (*p*-values \leq 0.05) were presented with different letters: lower-case letters for comparisons of EC50s at different temperature and different salinities after 24 h exposure; upper-case letters for comparisons of EC50s at different




Fig. 1. Photographic record of embryo-larvae types observed in *Crassostrea angulata* after embryotoxicity exposures: a) D-shape; b) pre-D; c) protruded mantle; d) kidney shape; e) indented shell; and f) dead larvae, after His et al. (1997).




Fig. 2. Contour plots representing the observed percentage of malformations at different combinations of salinity and temperature, for *C. angulata* after 24 h (A) and 48 h (B) development; and *C. gigas* after 24 h (C) and 48 h (D) development. Each colour represents a different category of percent frequency malformation. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

temperature levels and different salinities, after 48 h exposure; asterisks (*) when statistical differences between 24 h and 48 h exposures at each fixed combination of salinity and temperature were observed.

3. RESULTS and DISCUSSION

3.1. Embryonic development, salinity and temperature

3.1.1. *Crassostrea angulata*

C. angulata embryo-larval development under different combinations of salinity and temperature are depicted in Fig. 2A and B (24 and 48 h post fertilization respectively). Contour plots show that *C. angulata* presented low frequencies of malformed larvae (< 20%) at intermediate and high salinity levels (26 and 32) and temperatures of 24 and 28 °C, after 24 h development (Fig. 2B). On the other hand, *C. angulata* presented high frequencies of malformed larvae at every salinity level combined to low temperature (20 °C) (100% identified as Pre-D) after 24 h development (Fig. 2A). At higher temperatures and low salinity (20), results showed higher percentages of malformed larvae of 90% (28 °C) and 75% (24 °C).

Studies concerning this species teratogenesis date back to the 19th century, when the first trials on artificial fecundation of *C. angulata* were described for full salinity and 22 °C (Bouchon-Brandely, 1882), however scarce information is available on the effects of varying

salinity and temperature on this species embryonic development in laboratory conditions. Nonetheless, early studies by Amemiya (1926) found that the lowest limit of salinity for *C. angulata* embryonic development was 21, and optimum salinity to be between 28 and 35 (studies performed at 20–23,5 °C, 24 h). Similarly, our data showed that low salinity (20) induced poor embryonic development (75–100% malformations) at all tested temperatures, and malformation frequency decreased with the increase of salinity.

After 48 h of embryo-larval development, *C. angulata* showed lower frequencies of malformed larvae at the intermediate salinity (26) at all temperature levels (Fig. 2B), while relatively higher malformation percentages were observed at the highest salinity (33), at 26 and 28 °C at 48 h, compared to results obtained at 24 h (Fig. 2A).

The relative increase of the range of temperature and salinity at which embryos successfully developed into D-shape larvae observed between 24 and 48 h, reflected differences in developing rates that depend on the surrounding media characteristics, namely temperature and salinity. This was due to the fact that increasing the time of exposure allowed for embryo development completion. at suboptimal temperature and salinity conditions, because both these environmental factors affect oyster embryo development rate. For instance, studies on *Pinctata imbricata* pearl oysters from the tropical Atlantic (Urban, 2000) showed that suboptimal temperature and salinity delayed embryonic development (O'Connor and Lawler, 2004). This

Table 1

Summary table of median effect concentrations (EC50s) of As and relative 95% confidence intervals for *C. gigas* and *C. angulata* at different combinations of salinity and temperature, after 24 and 48 h exposures.

T (°C)	Salinity	<i>C. angulata</i>		<i>C. gigas</i>	
		24 h	48 h	24 h	48 h
20	20	n.c	n.c	n.c	101.1 (96.9–105.5)
	26	n.c	31.0 (n.d)	119.0 (114.6–123.5)	161.9 (147.2–178.0)
	33	n.c	n.c	285.5 (267.3–305.0)	530.0 (465.2–603.8)
24	20	n.c	n.c	101.9 (101.7–102.0)	80.71 (80.70–80.73)
	26	29.8 (29.8–29.8)	41.8 (38.7–45.0)	179.7 (178.6–180.7)	214.3 (206.1–222.9)
	33	39.2 (38.8–39.6)	18.7 (17.1–20.5)	451.5 (312.6–652.2)	663.5 (516.5–852.3)
28	20	n.c	n.c	103.4 (90.64–117.9)	103.3 (103.0–103.5)
	26	39.6 (39.5–39.7)	50.5 (44.0–58.1)	190.1 (180.9–199.8)	198.9 (185.5–213.3)
	33	36.2 (28.5–46.1)	15.9 (14.6–17.3)	303.7 (229.6–401.5)	493.7 (429.9–567.0)

n.c - EC50 value not calculated due to low or zero percentage of developed larvae (D-shape).

n.d - 95% confidence interval not determined due to few points for curve fit parameter estimation.

delayed effect is also observed in later stages, namely D-shape larvae of *C. angulata* (Thiyagarajan and Ko, 2012), *C. gigas* (His et al., 1989) and *P. imbricata* (Dove and O'Connor et al., 2007). Moreover, data obtained in the present study show that intermediate salinity (26) is likely closest to the optimum salinity for this species under laboratory conditions, the salinity at which we observed the widest thermal tolerance range (Le Dantec, 1968) (most noticed after 48 h development). This is in accordance with studies by Thiyagarajan and Ko (2012) that described optimum salinity for *C. angulata* D-shape larvae development between 24 and 27. Also in line with environmental studies showing higher survival rates of *C. angulata* larvae at temperatures over 22 °C and salinities between 28 and 32 (Le Dantec, 1968).

3.1.2. *Crassostrea gigas*

C. gigas embryo-larval development under different salinity and temperature combinations is depicted in Fig. 2C and D (24 and 48 h respectively). Contour plots show that *C. gigas* presented low frequencies of malformed larvae (< 30%) at all combinations of salinity and temperature after 24 h development, except for embryos exposed to low salinity (20) and low temperature (20 °C), where high percentages of malformed larvae were observed (100% identified as pre-D, Supplementary Table S2) (Fig. 2C).

Previous studies showed low percentages of D-shape *C. gigas* at 20 °C and full salinity (Gamain et al., 2017) (40% D-shape larvae after 24 h post fertilization), or even complete arrest of development at salinities ranging from 15 to 32 at 20 °C (100% trochophore stage at 24 h) (Moreira et al., 2018).

C. gigas embryos exposed to the same conditions for 48 h, showed high frequencies of developing D-shape larvae at all combinations of salinity and temperature (< 35% malformed larvae), including low salinity (20) and low temperature (20 °C) (Fig. 2D).

These results indicate some degree of variability in the developing rate of *C. gigas* at suboptimal temperature (20 °C) and are likely related to differences in development rates at suboptimal temperature. This is corroborated by results obtained at 48 h post-fertilization, for which high frequencies of D-shape larvae were observed in the present study, and in accordance with results from Parker et al. (2010) that also observed high rates of embryonic development at 48 h and low temperature (20 °C). Differences in methodology during the embryo-larval assay, namely the origin of seawater source (His et al., 1997; Beiras and His, 2004), as well as phenotypic plasticity and genetic variability (Pace et al., 2006; Taris et al., 2006 and references therein), could explain the differences observed between our findings and some of the results reported in literature.

3.1.3. Comparison between species

Differences between species successful embryonic development under different combinations of salinity and temperature were evident,

with *C. angulata* embryos presenting a narrower range of salinity and temperature for complete development than *C. gigas*. Implications of such findings under the eminence of climate change could mean differentiated resilience capacities between these species to cope with near future shifts in salinity and temperature regimes, and possibly constrain *C. angulata* populations' geographical distribution comparing to that of *C. gigas*.

3.2. Arsenic embryotoxicity

Results obtained from the embryotoxicity assay for each species showed that negative controls for both species were within the accepted values described for *C. gigas* embryotoxicity standard protocols (ASTM, 2004; His et al., 1999), with frequencies of normal developed larvae (D-shape) after 24 h development exceeding 70%, at standard salinity (33) and temperature (24 °C) for both species (*C. gigas* 75% and *C. angulata* 90%). Moreover, results obtained regarding positive controls (Cu) were within previously reported values ($9.47 \mu\text{g L}^{-1} \leq \text{EC50} \leq 21.72 \mu\text{g L}^{-1}$) (Libralato et al., 2009), and were similar for both species (10.07 (8.98–11.30) $\mu\text{g L}^{-1}$ of Cu for *C. angulata* and 11.40 (9.59–13.54) $\mu\text{g L}^{-1}$ of Cu for *C. gigas*). These results confirm the validity of the experiment and show that the sensitivity of both species to Cu were identical.

3.2.1. *Crassostrea angulata*

Arsenic toxicity measured in terms of EC50 considering standard toxicity assay conditions with oyster embryos (24 °C, 33 salinity, 24 h) for *C. angulata* was 39.2 $\mu\text{g L}^{-1}$ As (18.7 $\mu\text{g L}^{-1}$ As at 48 h) (Table 1). To our knowledge, this is the first study on the effects of contaminants to *C. angulata* embryonic development. The toxicity of As (EC50) measured in the present study for *C. angulata* was in the lower range of previously reported toxicity threshold values for related oyster species such as *C. virginica* (with LC50 of 7.2 mg L^{-1}) (Calabrese et al., 1973) and *C. gigas* (EC50 ranging from ca. 1–920 $\mu\text{g L}^{-1}$ of As) (Moreira et al., 2018; Mamindy-Pajany et al., 2013).

3.2.2. *Crassostrea gigas*

EC50 considering standard conditions for *C. gigas* (24 °C, 33 salinity, 24 h) of 452 $\mu\text{g L}^{-1}$ As (663.5 $\mu\text{g L}^{-1}$ As at 48 h) (Table 1), was within previously reported EC50 values (920 $\mu\text{g L}^{-1}$ As from Mamindy-Pajany et al. (2013) and 326 $\mu\text{g L}^{-1}$ As from Martin et al. (1981)). Moreover, in a previous study (Moreira et al., 2018) As toxicity determined at standard salinity and temperature assay conditions revealed an EC50 lower than 1 $\mu\text{g L}^{-1}$ As.

Among the few available data on As toxicity to bivalve embryos, a great variability is observed, in contrast to other types of pollutants (His et al., 1999). These discrepancies could be related to different reference seawater characteristics (Beiras and Albentosa, 2004), as observed in a

previous study (Moreira et al., 2018), or different methodological approaches (Martin et al. (1981), that used different incubation temperature (20 °C), different time (48 h) and As form AsO₅, or no reference to real As concentrations (Mamindy-Pajany et al., 2013).

3.2.3. Comparisons between species

Our results suggest that *C. angulata* is likely more sensitive to As than *C. gigas*, with at least an order of magnitude lower EC50 values (Table 1), despite that positive control for Cu toxicity showed similar results for both species, indicating sensitivity to As should be different between species, and was not related to batch's overall sensitivity. Several factors could be hypothesised to have influenced these results, but the most important one could be a differentiated species-specific sensitivity to As.

Indeed, species specific tolerance to As seemed to be the main reason explaining the differentiated toxicity observed. These results are supported by other studies, namely those that showed taxon related differences between *C. angulata* (Spanish origin) and *C. gigas* (French origin), regarding each progenies' maturation traits as well as mortality (Soletchnik et al., 2002). Also, Huvet et al. (2002), suggested that *C. gigas* could present overall better gamete quality than *C. angulata*, while studying hybrid crosses between both species. To our knowledge this is the first study on *C. angulata* embryotoxicity, and our data add to the body of evidence that show differences in ecophysiological traits between these closely related species (Soletchnik et al., 2002; His et al., 1972; Gouletquer et al., 1999; Heral et al., 1986; Moreira et al., 2016).

Other factors such as broodstock origin or parental exposure history could also have influenced each species sensitivity to As but were unlikely given our results and the existing literature as follows.

Gamain et al. (2017) studied *C. gigas* from different origins (hatchery, cultivated and wild) and found no major differences in embryotoxicity of metolachlor to embryos of different parental origins. However, and in contrast with our study, progeny from hatchery oysters were more sensitive to Cu than wild and cultivated ones. Considering this, it would be unlikely that oysters (hatchery *C. gigas*, and cultivated *C. angulata*) from our study, would have presented the observed differentiated response to As, solely based on broodstock origin (hatchery and cultivated).

A recent study by Weng and Wang (2017) demonstrated maternal transfer of trace metals in *C. hongkongensis* adults to their progeny (oocytes and larvae) reflecting parental exposure from contaminated sites, with negative impacts on the most contaminated embryos' development. Therefore, parental transfer of trace elements to zygotes in *C. angulata* could have influenced final As toxicity, knowing that the Sado estuary is anthropogenically polluted (Costa et al., 2009). However, the results obtained in the present study on trace element content in fertilized oocytes showed only marginal differences in trace element concentrations between species, except for Cr, Ni (higher in *C. angulata*) and Zn (higher in *C. gigas*) (Table 2). Trace element content from both *C. angulata* and *C. gigas* were in the lower range of those described by Weng and Wang (2017) for *C. hongkongensis* embryos, considering the

Table 2

Trace element concentration ($\mu\text{g}\cdot\text{g}^{-1}$ dry weight) determined for *C. angulata* and *C. gigas* fertilized oocytes suspensions used for embryotoxicity experiments (mean \pm RSD).

	<i>C. angulata</i>	<i>C. gigas</i>
As	3.43 \pm 0.031	3.29 \pm 0.035
Cd	0.11 \pm 0.0046	0.10 \pm 0.0054
Cr	3.69 \pm 0.049	0.36 \pm 0.0047
Cu	7.75 \pm 0.078	7.06 \pm 0.078
Hg	0.04 \pm 0.0051	0.03 \pm 0.0031
Ni	1.91 \pm 0.026	0.76 \pm 0.012
Sn	0.46 \pm 0.0085	0.40 \pm 0.0051
Zn	56.69 \pm 0.58	78.94 \pm 0.86

C. angulata




Fig. 3. Median effect concentrations (EC50s) of As and relative 95% confidence intervals for *C. angulata* at different salinity and temperature conditions, after 24 and 48 h exposure. Different letters represent statistical differences between tested groups ($p \leq 0.05$): lower-case letters for comparisons between combinations of salinity and temperature conditions after 24 h exposure; upper-case letters for comparisons between combinations of salinity and temperature conditions after 48 h exposure; asterisks (*) for statistical differences between 24 and 48 h exposures at each combination of salinity and temperature.

lowest reported values (lowest contaminated sites) for As, Cd, Cr, Cu, Ni and Zn (4.26 ± 0.83 ; 0.21 ± 0.11 ; 0.71 ± 0.39 ; 45.3 ± 21.0 ; 1.42 ± 1.03 ; $300 \pm 87.8 \mu\text{g}\cdot\text{g}^{-1}$ dry weight, respectively). Hence, in the present study, parental element transfer to embryos was not likely to have influenced final As toxicity in either species, given that we observed overall low contamination, and few differences in element concentrations in embryos between species.

3.3. Effects of salinity and temperature on species sensitivity to As

3.3.1. *Crassostrea angulata*

EC50 values determined for *C. angulata* at all tested combinations of salinity and temperature levels, at both 24 and 48 h of exposure are depicted in Fig. 3. For *C. angulata*, EC50 was not possible to calculate at the lowest salinity (20), regardless of temperature and duration of exposure. Moreover, at higher salinities and 20 °C, EC50 was only possible to determine for the intermediate salinity (26) and only at 48 h development (Fig. 3). These findings reflected the observed thermohaline tolerance of this species, with overall lower toxicity at optimum salinity (26).

To assess the effects of salinity on embryo sensitivity to As, pairwise comparisons among EC50 values obtained at each salinity were performed considering fixed temperature. Pairwise comparisons showed that at 24 h, EC50 values between salinities were significantly different only at 24 °C, for which EC50 was higher at salinity 33 than at salinity 26. On the other hand, results for *C. angulata* embryos exposed for 48 h, showed significantly higher EC50 values at salinity 26 at both compared temperature levels (24 and 28 °C) (Fig. 3). These findings illustrate that different salinities as well as time of exposure altered the sensitivity of embryos to As.

Salinity can influence oyster embryo development rate, by increasing the rate with the increase of salinity (O'Connor and Lawler, 2004), and therefore the retardation effect of As previously described (Moreira et al., 2018) at intermediate salinity (26) could be more noticeable than at higher salinity (33), as observed after 24 h development. However, these differences were counteracted after 48 h exposure, for which significantly higher EC50 values were observed at salinity 26 comparing to salinity 33, for both comparable temperature levels (24 and 28 °C). These results, showed that extending the exposure period from 24 to 48 h had a beneficial effect at salinity 26 (significant increase of EC50 at 24 °C), corroborating the above stated hypothesis that developing rates influenced final As toxicity. On the other hand, sensitivity to As (lower EC50) increased significantly at higher salinity (33) comparing 24 and 48 h exposures, likely resulting from poor

physiological status at higher salinities (Thiyagarajan and Ko, 2012), also supported by results obtained in negative controls at the same conditions for which we observed relatively high frequencies of dead larvae (data not shown).

To evaluate the effect of temperature on embryos sensitivity to As, pairwise comparisons between EC50 values obtained at different temperatures, within each salinity level were performed. Results showed significant increases of EC50s with the increase of temperature (24 and 28 °C) at salinity 26 for both 24 and 48 h exposures. At salinity 33 no significant differences were observed between temperature levels, at either 24 and 48 h exposures (Fig. 3).

Studies on the effects of temperature on the toxicity of pollutants to bivalve embryos, have reported contrasting results. For instance, increased toxicity with the increase of temperature has been shown for Cu and Ag (Boukadida et al. (2016), and for Pb (Hrs-Brecko, 1977) in *Mytillus galloprovincialis*. However, studies on *C. virginica* showed that Cu was more toxic at lower temperatures, presumably because closer to suboptimal temperature conditions would induce higher toxicity (MacInnes and Calabrese, 1979). Assuming that higher temperature increases the rate of chemical reactions in the media, the uptake of contaminants through biological membranes can be promoted at higher temperatures (Hazel, 1995), hence toxicity could be expected to increase at higher temperatures.

In the present study results obtained for *C. angulata* at optimum salinity (26) and both exposure periods (24 and 48 h) showed the opposite trend, for which the effect of increasing temperature resulted in lower As toxicity. These results were likely related to a counteractive effect of increased rates of embryo development at higher temperatures, with the retarding effect of As (Moreira et al., 2018), thus resulting in higher EC50s at higher temperatures.

3.3.2. *Crassostrea gigas*

Median effect concentrations (EC50's) of As under different salinity and temperature conditions after 24 and 48 h incubation for *C. gigas* are presented in Fig. 4. Results obtained show a clear increase of EC50's with the increase of salinity, considering both 24 and 48 h larval development, and fixed temperature levels (Fig. 4).

These results are in line with previous studies using *C. gigas* exposed to As (Moreira et al., 2018), Ag and Cu (Coglianese, 1982), Cu and metalochlor (Gamain et al., 2016) at different salinities. The increased sensitivity to As (lower EC50s) with the decrease of salinity is likely related to different osmoregulation status of oyster embryos in response to lower salinities. Under hypoosmotic conditions, the ion flux between the embryos and surrounding media may increase, inducing higher

uptake of soluble contaminant through active transport processes (Connel, 1989; Grosell et al., 2007). Another factor contributing to higher sensitivity to As at lower salinities, could be related to differences in As speciation with salinity variation, however we discarded such possibility, because under a similar range of salinity (and temperature) conditions, As speciation analysis showed no important differences in prevailing As species and bioavailability with varying salinity and temperature (Moreira et al., 2018).

The effect of varying temperature on *C. gigas* embryos' sensitivity to As, inferred by pairwise comparisons of EC50 values among temperature levels, were variable within each salinity level and incubation period (Fig. 4). At the lowest salinity (20), no significant differences were observed among EC50's obtained for all temperature levels (at both 24 and 48 h). At salinity 26, EC50 was significantly lower at 20 °C, comparing to the remaining temperature levels, after 24 h incubation. At salinity 33 after 24 h embryo development, the highest EC50 value was observed at 24 °C, with significant differences towards that obtained at 20 °C (Fig. 4).

These findings suggest that higher sensitivity (lower EC50s) observed at lower temperatures (20 °C, 24 h) for all tested salinity levels likely resulted from an additive effect of delayed development induced by both As and low temperature, considering that the developing rate of oyster larvae decrease at lower temperatures (His et al., 1989; Dove and O'Conner, 2007), and that As induces a retarding effect on embryo development (Moreira et al., 2018).

Results further revealed that after 48 h no significant differences were observed among temperature levels at salinity 26 and 33 (Fig. 4). These findings demonstrate that the arresting effect of As on *C. gigas* larvae development previously described is not permanent, because extending the exposure period to 48 h revealed higher frequencies of embryo-larval development completion (lower frequencies of Pre-D larvae) (Supplementary Table S2), which in turn resulted in similar toxicity effects to larvae exposed at different temperature levels at 48 h, unlike results obtained at 24 h for which temperature showed a higher effect.

Comparisons between EC50 values obtained for each combination of salinity and temperature at 24 h incubation and the corresponding conditions after 48 h exposure (Fig. 4), revealed no differences in EC50 values at any combination of low (20) and intermediate salinities (26) at every temperature level between 24 and 48 h exposures (Fig. 4 and Supplementary Table S3). However, significant differences were observed at all temperature levels tested at salinity 33, where significantly higher EC50 values were observed after 48 h comparing to values obtained after 24 h. It is important to note that at salinity 20 at 20 °C, EC50 was only possible to calculate after 48 h exposure, since after 24 h embryo development was not completed in these conditions (Fig. 2B and D).

Comparisons between results obtained for *C. gigas* at 24 and 48 h further revealed a significant increase of EC50s at the highest salinity (33), and all combinations of temperature, with the increase of exposure time. These results illustrated that toxicity threshold determination can depend on time of exposure for toxicants that affect embryonic development by retarding development such as As.

4. Concluding remarks

The present study allowed to assess the embryotoxicity of As under different thermohaline conditions to *C. angulata* and to compare results with those from an important worldwide distributed species *C. gigas*. Results obtained showed marked differences on each species embryo-development capacity, namely concerning the tolerance range to varying salinity and temperature, which further reflected in the pattern of As toxicity observed. *C. angulata* presented a narrower range of salinity and temperature than *C. gigas* for which embryo-development successfully occurred, considering 24 and 48 h post fertilization. *C. angulata* presented better embryonic development at intermediate




Fig. 4. Median effect concentrations (EC50s) of As and relative 95% confidence intervals for *C. gigas* at different salinity and temperature conditions, after 24 and 48 h exposure. Different letters represent statistical differences between tested groups ($p \leq 0.05$): lower-case letters for comparisons between combinations of salinity and temperature conditions after 24 h exposure; upper-case letters for comparisons between combinations of salinity and temperature conditions after 48 h exposure; asterisks (*) for statistical differences between 24 and 48 h exposures at each combination of salinity and temperature.

salinity (26) and at temperatures above 20 °C, while *C. gigas* presented high frequencies of developing embryos at all combinations of salinity and temperatures tested. Overall, these results suggest that early ontogeny of *C. angulata* may be limited to a narrower range of abiotic factors (salinity and temperature) compared to *C. gigas*, with possible implications at the population level. Considering that early life stages generally constitute the most susceptible stage of oysters' life cycle, the thermohaline differences observed for embryo development may dictate species competitive advantages towards one another under the projected scenarios of climate driven alterations of temperature and salinity regimes in estuarine systems worldwide. Hence, *C. angulata* may be more susceptible to environmental change than *C. gigas*.

Concerning As toxicity, EC50 values for *C. angulata* showed that the effect of As was overall lower at the intermediate salinity 26, however temperature also showed to influence As toxicity in an antagonistic manner, with higher temperatures inducing lower toxicity effects. For *C. gigas* the effect of As was highly dependent on salinity, with sensitivity to As showing to decrease with the increase of salinity. The effect of temperature was only noticeable after 24 h exposure, when low temperatures induced higher sensitivity, while after 48 h development the effect of temperature was mitigated.

Comparisons on As toxicity measured as EC50 values between species, showed that *C. angulata* was at least 10 times more sensitive to As than *C. gigas* and these differences were likely species related. Our results suggest that the survival of *C. angulata* strongly depends on a narrower range of abiotic factors compared to its closely related congener, and therefore the future of this population may be endangered considering projections on climate change and pollution worldwide.

Other issues rise from the present study, namely the observations that the delay effects of As on *Crassostrea* embryo development is not permanent (noted by differences between 24 and 48 h exposures) and add to recent findings that showed that sea urchin embryos can recover from toxicity induced by different trace metals (Morrone et al., 2018). These results are also important for future studies considering differentiated effects under varying physico-chemical parameters that may consider the 48-h criterion. For future studies relying on bivalves collected in the environment we suggest trace elements analysis to be performed on oocyte suspensions. All together these issues should be considered for future ecotoxicological studies and water quality framework directives relying on embryotoxicity assays.

Acknowledgements

This study was supported by the Portuguese Science Foundation (FCT) through CESAM (UID/AMB/50017), by FCT/MEC through national funds, and the co-funding by FEDER, within the PT2020 Partnership Agreement and Compete 2020. Anthony Moreira benefited from PhD grant SFRH/BD/93107/2013. Rosa Freitas benefited from a research position funded by the Integrated Programme of SR&TD “Smart Valorization of Endogenous Marine Biological Resources Under a Changing Climate” (reference Centro-01-0145-FEDER-000018), co-funded by Centro 2020 program, Portugal 2020, European Union, through the European Regional Development Fund. The authors would like to thank Mr. Pedro Ferreira from Exporsado for providing *C. angulata* progenitors.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.marenvres.2018.06.003>.

References

Amemiya, I., 1926. Notes on experiments on the early developmental stages of the Portuguese, American and English native oysters, with special reference to the effect of varying salinity. *J. Mar. Biol. Assoc. U. K.* 14, 161–176.

- ASTM, 2004. Standard Guide for Conducting Static Acute Toxicity Tests Starting with Embryos of Four Species of Saltwater Bivalve molluscs E724. American Society for Testing and Materials, West Conshohocken.
- Batista, F.M., López-Sanmartín, M., Grade, A., Morgado, I., Valente, M., Navas, J.I., Power, D.M., Ruano, F., 2015. Sequence variation in ostreid herpesvirus 1 microvar isolates detected in dying and asymptomatic *Crassostrea angulata* adults in the Iberian Peninsula: insights into viral origin and spread. *Aquaculture* 435, 43–51. <https://doi.org/10.1016/j.aquaculture.2014.09.016>.
- Beck, M.W., Brumbaugh, R.D., Airoldi, L., Carranza, A., Coen, L.D., Crawford, C., Defeo, O., Edgar, G.J., Hancock, B., Kay, M.C., Lenihan, H.S., Luckenbach, M.W., Toropova, C.L., Zhang, G., Guo, X., 2011. Oyster reefs at risk and recommendations for conservation, restoration, and management. *Bioscience* 61, 107–116. <https://doi.org/10.1525/bio.2011.61.2.5>.
- Beck, M.W., Brumbaugh, R.D., Airoldi, L., Carranza, A., Coen, L.D., Crawford, C., Defeo, O., Edgar, G.J., Hancock, B., Kay, M., Lenihan, H., Luckenbach, M.W., Toropova, C.L., Zhang, G., 2009. Shellfish Reefs at Risk: a Global Analysis of Problems and Solutions. The Nature Conservancy, Arlington VA 52 pp.
- Beiras, R., Albentosa, M., 2004. Inhibition of embryo development of the commercial bivalves *Ruditapes decussatus* and *Mytilus galloprovincialis* by trace metals; implications for the implementation of seawater quality criteria. *Aquaculture* 230, 205–213. [https://doi.org/10.1016/S0044-8486\(03\)00432-0](https://doi.org/10.1016/S0044-8486(03)00432-0).
- Beiras, R., His, E., 1994. Effects of dissolved mercury on embryogenesis, survival, growth and metamorphosis of *Crassostrea gigas* oyster larvae. *Mar. Ecol. Prog. Ser.* 113, 95–103. <https://doi.org/10.3354/meps113095>.
- Bouchon-Brandely, M., 1882. On the sexuality of the common oyster (*Ostrea edulis*) and that of the Portuguese oyster (*O. angulata*). Artificial fecundation of the Portuguese oyster. *Ann. Mag. Nat. Hist.* 10, 328–330. <https://doi.org/10.1080/00222938209459717>.
- Boukadida, K., Banni, M., Gourves, P.-Y., Cachot, J., 2016. High sensitivity of embryolarval stage of the Mediterranean mussel, *Mytilus galloprovincialis* to metal pollution in combination with temperature increase. *Mar. Environ. Res.* 122, 59–66. <https://doi.org/10.1016/j.marenvres.2016.09.007>.
- Brander, K.M., 2007. Global fish production and climate change. *Proc. Natl. Acad. Sci. Unit. States Am.* 104, 19709–19714. <https://doi.org/10.1073/pnas.0702059104>.
- Buestel, D., Ropert, M., Prou, J., Goulletquer, P., 2009. History, status, and future of oyster culture in France. *J. Shellfish Res.* 28, 813–820.
- Byrne, M., 2012. Global Change Ecotoxicology: Identification of Early Life History Bottlenecks in Marine Invertebrates, Variable Species Responses and Variable Experimental Approaches. *Marine Environmental Research, Emerging and Persistent Impacts on Marine Organisms: Detection Methods and Action Mechanisms*, vol. 76. pp. 3–15. <https://doi.org/10.1016/j.marenvres.2011.10.004>.
- Calabrese, A., Collier, R.S., Nelson, D.A., MacInnes, J.R., 1973. The toxicity of heavy metals to embryos of the American oyster *Crassostrea virginica*. *Mar. Biol.* 18, 162–166. <https://doi.org/10.1007/BF00367984>.
- Coen, L.D., Brumbaugh, R.D., Bushek, D., Grizzle, R., Luckenbach, M.W., Posey, M.H., Powers, S.P., Tolley, S.G., 2007. Ecosystem services related to oyster restoration. *Mar. Ecol. Prog. Ser.* 341, 303–307. <https://doi.org/10.3354/meps341303>.
- Coglianesi, M.P., 1982. The effects of salinity on copper and silver toxicity to embryos of the Pacific Oyster. *Arch. Environ. Contam. Toxicol.* 11, 297–303. <https://doi.org/10.1007/BF01055206>.
- Comps, M., 1988. Epizootic diseases of oysters associated with viral infections. *Am. Fish. Soc. Spec. Publ.* 18, 23–37.
- Connell, D.W., 1989. Bioaccumulation of Xenobiotic Compounds. CRC Press.
- Costa, P.M., Diniz, M.S., Caeiro, S., Lobo, J., Martins, M., Ferreira, A.M., Caetano, M., Vale, C., DelValls, T.Á., Costa, M.H., 2009. Histological biomarkers in liver and gills of juvenile *Solea senegalensis* exposed to contaminated estuarine sediments: a weighted indices approach. *Aquat. Toxicol.* 92, 202–212. <https://doi.org/10.1016/j.aquatox.2008.12.009>.
- Cranfield, H.J., Michael, K.P., Doonan, L.J., 1999. Changes in the distribution of epifaunal reefs and oysters during 130 years of dredging for oysters in Foveaux Strait, southern New Zealand. *Aquatic Conserv. Mar. Freshw. Ecosyst.* 9, 461–483. [https://doi.org/10.1002/\(SICI\)1099-0755\(199909/10\)9:5 < 461::AID-AQC353 > 3.0.CO;2-Z](https://doi.org/10.1002/(SICI)1099-0755(199909/10)9:5 < 461::AID-AQC353 > 3.0.CO;2-Z).
- Cross, I., Merlo, M.A., Rodríguez, M.E., Portela-Bens, S., Rebordinos, L., 2014. Adaptation to abiotic stress in the oyster *Crassostrea angulata* relays on genetic polymorphisms. *Fish Shellfish Immunol.* 41, 618–624. <https://doi.org/10.1016/j.fsi.2014.10.011>.
- De Gieter, M., Elskens, M., Baeyens, W., 2005. Fluxes and major transport routes of Arsenic in the Scheldt estuary. *Mar. Chem.* 95, 15–30. <https://doi.org/10.1016/j.marchem.2004.06.012>.
- Dekshenieks, M.M., Hofmann, E.E., Klinck, J.M., Powell, E.N., 2000. Quantifying the effects of environmental change on an oyster population: a modeling study. *Estuaries* 23 (593). <https://doi.org/10.2307/1352887>.
- Dove, M.C., O'connor, W.A., 2007. Salinity and temperature tolerance of sydney rock oysters *saccostrea glomerata* during early ontogeny. *J. Shellfish Res.* 26, 939–947. [https://doi.org/10.2983/0730-8000\(2007\)26\[939:SATTOS\]2.0.CO;2](https://doi.org/10.2983/0730-8000(2007)26[939:SATTOS]2.0.CO;2).
- Ereira, T., Coelho, J.P., Duarte, A.C., Pardal, M.A., Pereira, M.E., 2015. Size-Dependent Arsenic Accumulation in *Scrobicularia plana* in a Temperate Coastal Lagoon (Ria de Aveiro, Portugal). *Water Air Soil Pollut.* 226, 1–7. <https://doi.org/10.1007/s11270-015-2484-5>.
- Fabioux, C., Huvet, A., Lapegue, S., Heurtebise, S., Boudry, P., 2002. Past and present geographical distribution of populations of Portuguese (*Crassostrea angulata*) and Pacific (*C. gigas*) oysters along the European and north African Atlantic coasts. *Haliotis* 31, 33–44.
- Food and Agriculture Organization (FAO), 2015. World aquaculture production of fish, crustaceans, molluscs, etc., by principal species in 2013. Available at: fao.org/fishery/docs/STAT/summary/a-6.pdf.
- Fodrie, F.J., Rodriguez, A.B., Gittman, R.K., Grabowski, J.H., Lindquist, N.L., Peterson,

- C.H., Piehler, M.F., Ridge, J.T., 2017. Oyster reefs as carbon sources and sinks. *Proc. R. Soc. B* 284 (20170891). <https://doi.org/10.1098/rspb.2017.0891>.
- Galloway, J.M., Swindles, G.T., Jamieson, H.E., Palmer, M., Parsons, M.B., Sanei, H., Macumber, A.L., Timothy Patterson, R., Falck, H., 2018. Organic matter control on the distribution of arsenic in lake sediments impacted by ~65years of gold ore processing in subarctic Canada. *Sci. Total Environ.* 622–623, 1668–1679. <https://doi.org/10.1016/j.scitotenv.2017.10.048>.
- Gamain, P., Gonzalez, P., Cachot, J., Clérandeau, C., Mazzella, N., Gourves, P.Y., Morin, B., 2017. Combined effects of temperature and copper and S-metolachlor on embryonic development of the Pacific oyster. *Crassostrea gigas*. *Marine pollution bulletin* 115, 201–210.
- Gamain, P., Gonzalez, P., Cachot, J., Pardon, P., Tapie, N., Gourves, P.Y., Budzinski, H., Morin, B., 2016. Combined effects of pollutants and salinity on embryo-larval development of the Pacific oyster. *Crassostrea gigas*. *Marine Environmental Research* 113, 31–38. <https://doi.org/10.1016/j.marenvres.2015.11.002>.
- Gouletquer, P., Wolowicz, M., Latala, A., Geairon, P., Huvet, A., Boudry, P., 1999. Comparative analysis of oxygen consumption rates between cupped oyster spat of *Crassostrea gigas* of French, Japanese, Spanish and Taiwanese origins. *Aquat. Living Resour.* 12, 271–277.
- Grabowski, J.H., Brumbaugh, R.D., Conrad, R.F., Keeler, A.G., Opaluch, J.J., Peterson, C.H., Piehler, M.F., Powers, S.P., Smyth, A.R., 2012. Economic valuation of ecosystem services provided by oyster reefs. *Bioscience* 62, 900–909. <https://doi.org/10.1525/bio.2012.62.10.10>.
- Grosell, M., Blanchard, J., Brix, K.V., Gerdes, R., 2007. Physiology is pivotal for interactions between salinity and acute copper toxicity to fish and invertebrates. *Aquat. Toxicol.* 84, 162–172.
- Harley, C.D.G., Randall Hughes, A., Hultgren, K.M., Miner, B.G., Sorte, C.J.B., Thornber, C.S., Rodriguez, L.F., Tomanek, L., Williams, S.L., 2006. The impacts of climate change in coastal marine systems. *Ecol. Lett.* 9, 228–241. <https://doi.org/10.1111/j.1461-0248.2005.00871.x>.
- Hazel, J.R., 1995. Thermal adaptation in biological membranes: is homeoviscous adaptation the explanation? *Annu. Rev. Physiol.* 57, 19–42. <https://doi.org/10.1146/annurev.ph.57.030195.000315>.
- Henke, K., 2009. *Arsenic: Environmental Chemistry, Health Threats and Waste Treatment*. John Wiley & Sons.
- Heral, M., 1989. Traditional oyster culture in France. In: *Barnabe Aquaculture*, pp. 342–387.
- Heral, M., Deslous-Paoli, J.-M., Prou, J., 1986. Dynamiques des productions et des biomasses des huîtres creuses cultivées (*Crassostrea angulata* et *Crassostrea gigas*) dans le bassin de Marennes-Oléron depuis un siècle. In: *CIEM Conseil International Pour l'Exploration de La Mer*.
- His, E., 1972. Premiers éléments de comparaison entre l'huître portugaise et l'huître japonaise. *Sci. Peche* 219, 1–9.
- His, E., Beiras, R., Seaman, M.N.L., 1999. The Assessment of marine pollution - bioassays with bivalve embryos and larvae. In: Southward, A.J., P.A.T., C.M.Y. (Eds.), *Advances in Marine Biology*. Academic Press, pp. 1–178.
- His, E., Robert, R., Dinot, A., 1989. Combined effects of temperature and salinity on fed and starved larvae of the mediterranean mussel *Mytilus galloprovincialis* and the Japanese oyster *Crassostrea gigas*. *Mar. Biol.* 100, 455–463. <https://doi.org/10.1007/BF00394822>.
- His, E., Seaman, M.N.L., Beiras, R., 1997. A simplification the bivalve embryogenesis and larval development bioassay method for water quality assessment. *Water Res.* 31, 351–355. [https://doi.org/10.1016/S0043-1354\(96\)00244-8](https://doi.org/10.1016/S0043-1354(96)00244-8).
- Hoegh-Guldberg, O., Bruno, J.F., 2010. The impact of climate change on the World's marine ecosystems. *Science* 328, 1523–1528. <https://doi.org/10.1126/science.1189930>.
- Hrs-Brenko, M., Claus, C., Bubic, S., 1977. Synergistic effects of lead, salinity and temperature on embryonic development of the mussel *Mytilus galloprovincialis*. *Mar. Biol.* 44, 109–115. <https://doi.org/10.1007/BF00386951>.
- Hsiao, S.-T., Chuang, S.-C., Chen, K.-S., Ho, P.-H., Wu, C.-L., Chen, C.A., 2016. DNA barcoding reveals that the common cupped oyster in Taiwan is the Portuguese oyster *Crassostrea angulata* (Ostreoida; Ostreidae), not *C. gigas*. *Sci. Rep.* 6 (34057). <https://doi.org/10.1038/srep34057>.
- Huvet, A., Gérard, A., Ledu, C., Phélipot, P., Heurtebise, S., Boudry, P., 2002. Is fertility of hybrids enough to conclude that the two oysters *Crassostrea gigas* and *Crassostrea angulata* are the same species? *Aquat. Living Resour.* 15, 45–52. [https://doi.org/10.1016/S0990-7440\(01\)01148-2](https://doi.org/10.1016/S0990-7440(01)01148-2).
- IPCC, 2014. *Core Writing Team* In: Pachauri, R.K., Meyer, L.A. (Eds.), *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. IPCC, Geneva, Switzerland, pp. 151.
- Knezovich, J.P., Harrison, F.L., Tucker, J.S., 1981. The influence of organic chelators on the toxicity of copper to embryos of the Pacific oyster, *Crassostrea gigas*. *Arch. Environ. Contam. Toxicol.* 10, 241–249. <https://doi.org/10.1007/BF01055625>.
- Knight, K., 2017. High and dry oysters at most risk from climate change. *J. Exp. Biol.* 220, 734–734. <https://doi.org/10.1242/jeb.157974>.
- Le Dantec, J., 1968. *Ecologie et reproduction de l'huître portugaise (Crassostrea Angulata LAMARCK) dans le bassin d'Arcachon et sur la rive gauche de la Gironde*. *Rev. Trav. Inst. Peche Marit.* 32, 237–362.
- Lejeune, C., Chevaldonné, P., Pergent-Martini, C., Boudouresque, C.F., Pérez, T., 2010. Climate change effects on a miniature ocean: the highly diverse, highly impacted Mediterranean Sea. *Trends Ecol. Evol.* 25, 250–260. <https://doi.org/10.1016/j.tree.2009.10.009>.
- Lenihan, H.S., Peterson, C.H., 1998. How habitat degradation through fishery disturbance enhances impacts of hypoxia on oyster reefs. *Ecol. Appl.* 8, 128–140. [https://doi.org/10.1890/1051-0761\(1998\)008\[0128:HHDTFD\]2.0.CO;2](https://doi.org/10.1890/1051-0761(1998)008[0128:HHDTFD]2.0.CO;2).
- Leverett, D., Thain, J., 2013. *ICES Techniques in Marine Environmental Sciences: Oyster Embryo-larval Bioassay (Revised)*.
- Levinton, J., Doall, M., Ralston, D., Starke, A., Allam, B., 2011. Climate change, precipitation and impacts on an estuarine refuge from disease. *PLoS One* 6, e18849. <https://doi.org/10.1371/journal.pone.0018849>.
- Libralato, G., Losso, C., Avezzi, F., Volpi Ghirardini, A., 2009. Influence of the salinity adjustment methods, salts and brine, on the toxicity of wastewater samples to mussel embryos. *Environ. Technol.* 30, 85–91. <https://doi.org/10.1080/09593330802505094>.
- Libralato, G., Losso, C., Ghirardini, A.V., 2007. Toxicity of untreated wood leachates towards two saltwater organisms (*Crassostrea gigas* and *Artemia franciscana*). *J. Hazard Mater.* 144, 590–593. <https://doi.org/10.1016/j.jhazmat.2006.10.082>.
- MacInnes, J.R., Calabrese, A., 1979. Combined effects of salinity, temperature, and copper on embryos and early larvae of the American oyster, *Crassostrea virginica*. *Arch. Environ. Contam. Toxicol.* 8, 553–562. <https://doi.org/10.1007/BF01055036>.
- Mamindy-Pajany, Y., Libralato, G., Roméo, M., Hurel, C., Losso, C., Volpi Ghirardini, A., Marmier, N., 2010. Ecotoxicological evaluation of Mediterranean dredged sediment ports based on elutriates with oyster embryotoxicity tests after composting process. *Water Res.* 44 (6), 1986–1994. <http://dx.doi.org/10.1016/j.watres.2009.11.056>.
- Mamindy-Pajany, Y., Hurel, C., Gêret, F., Galgani, F., Battaglia-Brunet, F., Marmier, N., Roméo, M., 2013. Arsenic in marine sediments from French Mediterranean ports: geochemical partitioning, bioavailability and ecotoxicology. *Chemosphere* 90, 2730–2736. <https://doi.org/10.1016/j.chemosphere.2012.11.056>.
- Martin, M., Osborn, K.E., Billig, P., Glickstein, N., 1981. Toxicities of ten metals to *Crassostrea gigas* and *Mytilus edulis* embryos and Cancer magister larvae. *Mar. Pollut. Bull.* 12, 305–308. [https://doi.org/10.1016/0025-326X\(81\)90081-3](https://doi.org/10.1016/0025-326X(81)90081-3).
- Masson, M., Schäfer, J., Blanc, G., Pierre, A., 2007. Seasonal variations and annual fluxes of arsenic in the garonne, dordogne and isle rivers, France. *Sci. Total Environ.* 373, 196–207. <https://doi.org/10.1016/j.scitotenv.2006.10.039>.
- Miossec, L., Le Deuff, R.-M., Gouletquer, P., 2009. *Alien Species Alert: Crassostrea gigas (Pacific Oyster)*. ICES Cooperative Research Report 299.
- Moreira, A., Figueira, E., Soares, A.M.V.M., Freitas, R., 2016a. The effects of arsenic and seawater acidification on antioxidant and biomineralization responses in two closely related *Crassostrea* species. *Sci. Total Environ.* 545–546, 569–581. <https://doi.org/10.1016/j.scitotenv.2015.12.029>.
- Moreira, A., Figueira, E., Soares, A.M.V.M., Freitas, R., 2016b. Salinity influences the biochemical response of *Crassostrea angulata* to Arsenic. *Environ. Pollut.* 214, 756–766. <https://doi.org/10.1016/j.envpol.2016.04.036>.
- Moreira, A., Freitas, R., Figueira, E., Volpi Ghirardini, A., Soares, A.M.V.M., Radaelli, M., Guida, M., Libralato, G., 2018. Combined effects of arsenic, salinity and temperature on *Crassostrea gigas* embryotoxicity. *Ecotoxicol. Environ. Saf.* 147, 251–259. <https://doi.org/10.1016/j.ecoenv.2017.08.043>.
- Morroni, L., Pinsino, A., Pellegrini, D., Regoli, F., 2018. Reversibility of trace metals effects on sea urchin embryonic development. *Ecotoxicol. Environ. Saf.* 148, 923–929. <https://doi.org/10.1016/j.ecoenv.2017.11.013>.
- Neff, J., 2002. *Arsenic in the Ocean*.
- O'Connor, W.A., Lawler, N.F., 2004. Salinity and temperature tolerance of embryos and juveniles of the pearl oyster, *Pinctada imbricata* Röding. *Aquaculture* 229, 493–506. [https://doi.org/10.1016/S0044-8486\(03\)00400-9](https://doi.org/10.1016/S0044-8486(03)00400-9).
- Pace, D., Marsh, A., Leong, K., Green, J., Hedgecock, D., Manahan, T., 2006. *Physiological Bases of Genetically Determined Variation in Growth of Marine Invertebrate Larvae: A Study of Growth Heterosis in the Bivalve Crassostrea gigas*.
- Parker, L.M., Ross, P.M., O'Connor, W.A., 2010. Comparing the effect of elevated pCO₂ and temperature on the fertilization and early development of two species of oysters. *Mar. Biol.* 157, 2435–2452. <https://doi.org/10.1007/s00227-010-1508-3>.
- Qin, J., Huang, Z., Chen, J., Zou, Q., You, W., Ke, C., 2012. Sequencing and de novo Analysis of *Crassostrea angulata* (Fujian Oyster) from 8 Different Developing Phases Using 454 GSFlx. *PLoS One* 7. <https://doi.org/10.1371/journal.pone.0043653>.
- Ridge, J.T., Rodriguez, A.B., Fodrie, F.J., 2017. Evidence of exceptional oyster-reef resilience to fluctuations in sea level. *Ecol. Evol. n/a–n/a*. n.d. <https://doi.org/10.1002/eec3.3473>.
- Ringwood, A.H., 1990. The relative sensitivities of different life stages of *Osogomon californicum* to cadmium toxicity. *Arch. Environ. Contam. Toxicol.* 19, 338–340.
- Robins, P.E., Skov, M.W., Lewis, M.J., Giménez, L., Davies, A.G., Malham, S.K., Neill, S.P., McDonald, J.E., Whitton, T.A., Jackson, S.E., Jago, C.F., 2016. Impact of climate change on UK estuaries: a review of past trends and potential projections. *Estuar. Coast Shelf Sci.* 169, 119–135. <https://doi.org/10.1016/j.ecss.2015.12.016>.
- Ruano, F., 1997. Fisheries and farming of important marine bivalves in Portugal. The history, present condition and future of the molluscan fisheries of North and Central America and Europe 3, 191–200.
- Ruesink, J.L., Lenihan, H.S., Trimble, A.C., Heiman, K.W., Micheli, F., Byers, J.E., Kay, M.C., 2005. Introduction of non-native oysters: ecosystem effects and restoration implications. *Annu. Rev. Ecol. Systemat.* 36, 643–689. <https://doi.org/10.1146/annurev.ecolsys.36.102003.152638>.
- Soletchnik, P., Huvet, A., Le Moine, O., Razet, D., Geairon, P., Faury, N., Gouletquer, P., Boudry, P., 2002. A comparative field study of growth, survival and reproduction of *Crassostrea gigas*, *C. angulata* and their hybrids. *Aquat. Living Resour.* 15, 243–250. [https://doi.org/10.1016/S0990-7440\(02\)01175-0](https://doi.org/10.1016/S0990-7440(02)01175-0).
- Taris, N., Ernande, B., McCombie, H., Boudry, P., 2006. Phenotypic and genetic consequences of size selection at the larval stage in the Pacific oyster (*Crassostrea gigas*). *J. Exp. Mar. Biol. Ecol.* 333, 147–158. <https://doi.org/10.1016/j.jembe.2005.12.007>.
- Thiyagarajan, V., Ko, G.W.K., 2012. Larval growth response of the Portuguese oyster (*Crassostrea angulata*) to multiple climate change stressors. *Aquaculture* 370–371, 90–95. <https://doi.org/10.1016/j.aquaculture.2012.09.025>.
- Urban, H.-J., 2000. Culture potential of the pearl oyster (*Pinctada imbricata*) from the Caribbean: I. Gametogenic activity, growth, mortality and production of a natural

- population. *Aquaculture* 189, 361–373. [https://doi.org/10.1016/S0044-8486\(00\)00393-8](https://doi.org/10.1016/S0044-8486(00)00393-8).
- Virvilis, C., Angelidis, P., 2006. Presence of the parasite *Marteilia* sp. in the flat oyster (*Ostrea edulis* L) in Greece. *Aquaculture* 259, 1–5. <https://doi.org/10.1016/j.aquaculture.2006.05.008>.
- Weng, N., Wang, W.-X., 2017. Dynamics of maternally transferred trace elements in oyster larvae and latent growth effects. *Sci. Rep.* 7 (3580). <https://doi.org/10.1038/s41598-017-03753-2>.
- Zanette, J., de Almeida, E.A., da Silva, A.Z., Guzinski, J., Ferreira, J.F., Di Mascio, P., Marques, M.R.F., Bainy, A.C.D., 2011. Salinity influences glutathione S-transferase activity and lipid peroxidation responses in the *Crassostrea gigas* oyster exposed to diesel oil. *Sci. Total Environ.* 409, 1976–1983. <https://doi.org/10.1016/j.scitotenv.2011.01.048>.