



Review Article

The lullaby of the sun: the role of vitamin D in sleep disturbance

Giovanna Muscogiuri ^{a,*}, Luigi Barrea ^{a,1}, Marianna Scannapieco ^a, Carolina Di Somma ^b, Massimo Scacchi ^c, Gianluca Aimaretti ^d, Silvia Savastano ^a, Annamaria Colao ^a, Paolo Marzullo ^{c,d}

^a Dipartimento di Medicina Clinica e Chirurgia, Sezione di Endocrinologia, Università Federico II di Napoli, Naples, Italy

^b IRCCS SDN, Napoli Via Gianturco 113, 80143, Naples, Italy

^c IRCCS, Istituto Auxologico Italiano, Div. General Medicine, Ospedale San Giuseppe, Piancavallo, Verbania, Italy

^d Università del Piemonte Orientale, Endocrinology, Dept. Translational Medicine, Novara



ARTICLE INFO

Article history:

Received 26 August 2018

Accepted 17 October 2018

Available online 24 November 2018

Keywords:

Sleep disturbance

Sleep

Vitamin D

Sun

Vitamin

ABSTRACT

Vitamin D is a pro-hormone belonging to the category of the fat-soluble group of vitamins; it is obtained more from solar exposure and in smaller quantities through feeding. Although vitamin D has traditionally been shown to be involved in calcium homeostasis and bone health, recent studies have found a positive association between vitamin D and sleep. In particular, clinical studies in humans indicate that low levels of vitamin D are correlated with poor quality and short sleep duration. The mechanism by which this association is explained is still unclear. However, vitamin D receptors have been found in the brain regions involved in sleep regulation, and vitamin D appears to be involved in regulating the sleep–wake cycle. The current review summarizes the available evidence regarding the association between vitamin D and sleep, focusing on both clinical and preclinical studies.

© 2018 Published by Elsevier B.V.

1. Introduction

Vitamin D is a pro-hormone belonging to the category of fat-soluble vitamins group; it is synthesized in vivo when solar ultraviolet B (UVB) radiation interacts with the precursor molecule, 7-dehydrocholesterol, in the skin [1–3]. Although endogenous production is estimated to account for 90% of total vitamin D in healthy individuals, a minor source of vitamin D comes from dietary intake and supplementation [1,4]. Vitamin D is subsequently transported in the blood (bound to vitamin D-binding protein) to the liver where it is hydroxylated to 25-hydroxyvitamin D (25-(OH)D) [3]. 25-(OH)D is further converted to the metabolically active form, 1 α , 25-dihydroxyvitamin D (1 α , 25-(OH)2D), primarily in the kidneys [5].

Calcitriol (1 α , 25-(OH) 2D) binds to the vitamin D receptor (VDR), a transcription factor that translocates to the cell nucleus, in which it is heterodimerized with the retinoid X receptor (RXR).

Then, the calcitriol-VDR-RXR complex couples to a specific sequence of DNA known as the.

Vitamin D response element (VDRE), which is associated with promoters of several target genes and co-activator molecules [4].

Many conditions interfere with the endogenous synthesis of vitamin D such as low sun exposure [6], use of sunscreen [7], age [1], race/ethnicity and skin color [8], season, altitude, and latitude [9].

Vitamin D deficiency has been historically defined and recently recommended by the Institute of Medicine (IOM) as a 25(OH)D of less than 20 ng/ml. Vitamin D insufficiency has been defined as a 25(OH)D of 21–29 ng/ml [9–15]. While vitamin D has traditionally been shown to be involved in calcium homeostasis [16] and bone health [1], recent evidence suggests extraskeletal effects. An inadequate level of vitamin D has been linked to a number of diseases including metabolic disorders, psychiatric, respiratory and cardiovascular disorders, autoimmune conditions, and cancers as well as osteoporosis and osteomalacia [1,17,18]. The widespread systemic effects of vitamin D have been attributed to the ubiquitous expression of vitamin D receptors in various organ systems [1,19].

Recently there is growing evidence that vitamin D could play a role in sleep regulation. Low vitamin D levels have been reported to be associated with shorter sleep duration in the National Health and Nutrition Examination Survey (NHANES) [20]. Furthermore, it

* Corresponding author. Department of Clinical Medicine and Surgery, University "Federico II" Via Sergio Pansini, 5 I-80131 Naples Italy. Fax: +815465443.

E-mail address: giovanna.muscogiuri@gmail.com (G. Muscogiuri).

¹ equally contributed to the manuscript.

has been reported that adequate levels of vitamin D are necessary for the maintenance of sleep, reducing the number of nocturnal awakenings [14].

The exact mechanism by which vitamin D regulates sleep is still far from known. Basic studies reported the presence of vitamin D receptor on areas of the brainstem that are known to be pacemaker cells playing an important role in the first stage of sleep and in maintaining sleep. Given the key role of vitamin D in sleep regulation, we aim to review observational studies that report the association between vitamin D status and sleep disturbances (both sleep quality and duration). Moreover, we provide molecular insights to explain this association to better understand if vitamin D supplementation could have a role in the improvement of sleep disturbances.

2. Vitamin D and sleep disturbance: observational studies and clinical trial

Massa et al., [21] found a positive association between low levels of vitamin D and short sleep duration in a cross-sectional study conducted between December 2003 and March 2005 on a large cohort of men aged ≥ 65 years. Similar findings were reported by Bertisch et al., [22] who found a modest association between 25(OH)D deficiency and short sleep duration measured in a cross-sectional study conducted between 2000 and 2002 in a large, multiracial/ethnic, community-based cohort. The authors performed a race/ethnicity-specific analysis suggesting that this association persisted in an African Americans subgroup in which 25(OH)D deficiency was associated with a mean reduction of 25 min in average nightly sleep duration. Data from the Korean National Health and Nutrition Examination [Survey_2013] reported an association between short sleep duration and vitamin D deficiency in Korean women [23].

As a consequence of this background, Wei Huang et al. [24] found an improvement in the sleep duration of about 45 min in a prospective study of a group of patients supplemented with vitamin D (dosage tailored according to the serum vitamin D of each patient) for three months.

However, vitamin D not only influences sleep duration, but it has been reported to have also an effect on sleep quality. Young Saeng Jung et al., [25] found a significant association between serum vitamin D deficiency and poor sleep quality in a cross-sectional study conducted between January 2015 and December 2015 on a group of workers in the manufacturing industry who worked indoors. Similar results were reported in a cross-sectional study conducted in community-resident men aged ≥ 65 years [21]. The association between low levels of vitamin D and sleep quality has also been confirmed in special categories of patients. An association between low levels of vitamin D and poor sleep quality has been found in hemodialysis patients [26]. Tuck Seng Cheng et al., [27] found that vitamin D deficiency correlated with poor sleep quality in a prospective cohort study conducted between June 2009 and September 2010 in Singapore, in women who were in the first trimester of pregnancy. Another study investigated the association of vitamin D and sleep quality in pregnancy finding no relationship [28]. However, the study performed by Suzan Gunduz et al., was performed in women in the last trimester of pregnancy in which several pregnancy-related factors could contribute to impair sleep quality [28]. Vitamin D supplementation for three months improved sleep quality in a prospective study by Wei Huang et al., [24]. Opposite results were reported by Caitlin Mason et al., [29] who found no association between vitamin D and sleep quality in a double-blind, placebo-controlled randomized clinical trial conducted in Seattle between 2010 and 2012 in overweight or obese postmenopausal women with vitamin D deficiency; which had

been supplemented with 2000 IU per day of vitamin D for 12 months. The negative results of this study could be explained by the fact that women enrolled in the study were overweight/obese, and the amount of vitamin D that was used to supplement them could be underestimated for the BMI.

3. Molecular mechanisms: animal and cell studies

The exact mechanism by which vitamin D affects sleep regulation is still unclear, although it has been suggested that some areas of the brainstem, which are involved in sleep regulation, could be the missing link [14]. This hypothesis is supported by evidence of the expression of vitamin D receptors in areas of the brainstem such as the anterior and posterior hypothalamus, substantia nigra, midbrain central grey matter, raphe nuclei, and in the nucleus reticularis pontis oralis and caudalis that are involved in sleep regulation. Indeed, in these areas there are pacemaker cells that play an important role in the first stages of sleep and in maintaining sleep. Using radiolabeled 1,25(OH)₂ vitamin D₃ and the sensitive receptor microautoradiography method, vitamin D target neurons (those with nuclear concentrations of the hormone) have been discovered in specific brain and spinal cord areas in several animals [30–33]. In rats, nuclear uptake and retention of radiolabeled 1,25(OH)₂ vitamin D₃ has been demonstrated in neurons located in the midbrain central gray, the nucleus raphes dorsalis, and the nucleus reticularis pontis oralis and caudalis [30]. Vitamin D target cells are also present in the basal forebrain, the hypothalamic periventricular region, and preoptic-septal regions. In the thalamus, the reticular nucleus is strongly labeled, as well [30–32]. Target neurons of vitamin D have been found in the bed nucleus of the stria terminalis that are linked to the nucleus of the central amygdala and contiguous with labeled neurons in the piriform and entorhinal cortex, as well as the ventral hippocampus [30–32]. In the hamster brain, vitamin D target neurons have been noted in the midbrain central gray nucleus of Darkschewitsch, the interstitial nucleus of Cajal, the nucleus tractus optici lateralis and medialis, and the substantia nigra [31]. In the human brain, immunohistochemical studies using antibodies that bind vitamin D receptor proteins have provided evidence for vitamin D target neurons in the substantia nigra [33].

Recent experimental studies [34–36] have suggested that the central circadian clock may also be regulated by vitamin D due to the presence of both 1,25-dihydroxycholecalciferol (VDR) receptor and 1 α -hydroxylase (1 α -OHase) in human brain [34].

Human behavioral rhythms involve rest and activity cycles such as sleep/wakefulness, which are synchronized by environmental signals, particularly dark/light intervals, across a 24-h period [37]. The synchronization process involves hierarchical mechanisms where the central circadian clock at the hypothalamic suprachiasmatic nucleus (SCN) is entrained to sunlight via the retina and subsequently influences peripheral oscillators through hormonal and neural pathways [38]. Given that sunlight drives both vitamin D synthesis [1] and circadian rhythms [38], it is possible that vitamin D is involved in the transduction of light signals that regulate circadian rhythms [39].

It has been shown that the VDR as well as 25-hydroxylase and 1-hydroxylase, the enzymes controlling vitamin D activation, along with 24-CYP24A1, the enzyme controlling vitamin D degradation, are expressed in the brain [34,40,41]. Thus, the central nervous system (CNS) can synthesize its own vitamin D, which yields auto- or paracrine neurosteroid actions at the local level [34]. Similar to other neuroactive steroids, Vitamin D modulates neuronal excitability by acting on spontaneous regular firing, action potential duration, intrinsic excitability, and sensitivity to neurotransmitters as well as to neurotransmitter receptors such as the GABA receptor and the N-




Fig. 1. Mechanism of action of vitamin D on sleep regulation. Vitamin D has been reported to bind in several areas in the brainstem such as the anterior and posterior hypothalamus, substantia nigra, midbrain central grey matter, raphe nuclei, and in the nucleus reticularis pontisoralis and caudalis that have a role in sleep regulation. Furthermore, vitamin D plays a pivotal role in the synthesis of melatonin.

methyl-D-aspartate (NMDA) receptor [42]. In addition, vitamin D upregulates the synthesis of neurotrophins such as neural growth factor (NGF), neurotrophin 3 (NT3), and glial cell line-derived neurotrophic factor (GDNF), whereas neurotrophin 4 (NT4) is down-regulated [43]. Therefore, through multiple loops, vitamin D can potentially affect the development, maintenance, and survival of neurons. Moreover, hypothetically, low levels of vitamin D during early life could be relevant for future development of several brain diseases, such as schizophrenia and multiple sclerosis [44]. Whether this link also extends to sleep control is currently unknown.

Nevertheless, previous studies showed VDR expression in both developing [45] and adult rat brain [30]. Receptor sites for vitamin D were found in forebrain nuclei of the Siberian hamster that are known to be affected by neurodegenerative processes such as Alzheimer's disease as well as in adult hamster brain [32]. In the human brain, the VDR distribution has been described as strikingly similar to that detected in rodents [34]. It is found expressed in cortical and subcortical areas involved in sleep control, such as [a] the prefrontal cortex, which mediates normal sleep physiology and sleep-deprivation phenomena during NREM and REM sleep [46]; [b] the cingulate gyrus, which is activated by breathing and blood pressure challenges that are affected by sleep apnea [47]; [c] the hippocampal dentate gyrus, where adult neurogenesis is significant [48] and is influenced by sleep deprivation [49]; [d] in caudate nucleus, which is downregulated in disturbed sleep and insomnia [50]; [e] the lateral geniculate nucleus, a relay center in the thalamus for the visual pathway which plays a major role in ponto-geniculo-occipital waves during REM sleep [51]; [f] substantia nigra, which is claimed to regulate the sleep–wake cycle [52] and is involved in idiopathic rapid-eye-movement sleep behaviour disorder [53]; as well as [g] the anterior hypothalamus, particularly the supraoptic and paraventricular nuclei, which are located in tight contiguity with and receive efferent neural pathways [54] from the ventrolateral pre-optic nucleus (VLPO), a hypothalamic site with primary functions in the regulation of NREM and REM sleep [55]. In rat supraoptic and

paraventricular neurons, immunostaining for vitamin D binding protein co-localized with Arg-vasopressin [56], which not only regulates fluid balance but also partakes in REM sleep regulation under physiological conditions [57]. Furthermore, there is evidence that vitamin D contributes to regulating the production of melatonin, the pineal hormone controlling human circadian rhythms and sleep. Through actions on tryptophan hydroxylases (TPH)-2, which expresses VDRE at the gene level [58], vitamin D can regulate tryptophan conversion into 5-hydroxytryptophan, which is metabolized to serotonin to produce the sleep hormone, melatonin [59]. It is perhaps worth noting here that, in patients with MS, melatonin has been hypothesized to act as a mediator neuro-immunomodulatory effects of vitamin D [60]. These associations notwithstanding, experimental probing of vitamin D's role in sleep regulation is still scarcely reconnoitered (Fig. 1).

4. Conclusion

Although several findings support the role of vitamin D in Sleep Disturbance, no guidelines are currently available for or against recommending vitamin D supplementation in the prevention or therapy of this disease. Due to the paucity of intervention studies, further clinical trials are needed before evidence-based recommendations can be made to investigate the potential clinical effectiveness of vitamin D supplementation on sleep regulation.

Conflict of interest

The ICMJE Uniform Disclosure Form for Potential Conflicts of Interest associated with this article can be viewed by clicking on the following link: <https://doi.org/10.1016/j.sleep.2018.10.033>.

References

- [1] Holick MF. Vitamin D deficiency. *N Engl J Med* 2007;357(3):266–81.

- [2] Fuleihan G, Bouillon R, Clarke B, et al. Serum 25-hydroxyvitamin D levels: variability, knowledge gaps, and the concept of a desirable range. *J Bone Miner Res* 2015;30(7):1119–33.
- [3] Awada Z, Ossaily S, Zgheib N. The nutrigenetics and pharmacogenetics of vitamin D pathways. *Curr Pharmacogenomics Personalized Med (CPPM)* 2014;12(2):89–103.
- [4] Holick MF, Binkley NC, Bischoff-Ferrari HA, et al. Endocrine Society. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2011;96(7):1911–30.
- [5] Lehmann B, Meurer M. Extrarenal sites of calcitriol synthesis: the particular role of the skin. *Recent Results Cancer Res* 2003;164:135–45.
- [6] Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem with health consequences. *Am J Clin Nutr* 2008;87(4):1080–6.
- [7] Nair R, Maseeh A. Vitamin D: the “sunshine” vitamin. *J Pharmacol Pharmacother* 2012;3(2):118–26.
- [8] McCarty DE, Reddy A, Keigley Q, et al. Vitamin D, race, and excessive daytime sleepiness. *J Clin Sleep Med* 2012;8(6):693–7.
- [9] Moan J, Porojnicu AC, Dahlback A, et al. Addressing the health benefits and risks, involving vitamin D or skin cancer, of increased sun exposure. *Proc Natl Acad Sci U S A* 2008;105(2):668–73.
- [10] Jung IK. Prevalence of vitamin deficiency in Korea: results from KNHANES 2010 to 2011. *J Nutr Health* 2013;46(6):540–51.
- [11] Renzaho AM, Halliday JA, Nowson C. Vitamin D, obesity, and obesity-related chronic disease among ethnic minorities: a systematic review. *Nutrition* 2011;27(9):868–79.
- [12] Chen TC, Holick MF. Vitamin D and prostate cancer prevention and treatment. *Trends Endocrinol Metab* 2003;14:423–30.
- [13] Kamen DL, Vin T. Vitamin D and molecular actions on the immune system: modulation of innate and autoimmunity. *J Mol Med* 2010;88(5):441–50.
- [14] Gominak SC, Stumpf WE. The world epidemic of sleep disorders is linked to vitamin D deficiency. *Med Hypotheses* 2012;79:132–5.
- [15] Skaer TL, Sclar DA. Economic implications of sleep disorders. *Pharmacoeconomics* 2010;28:1015–23.
- [16] Christakos S, Dhawan P, Liu Y, et al. New insights into the mechanisms of vitamin D action. *J Cell Biochem* 2003;88(4):695–705.
- [17] Grober U, Spitz J, Reichrath J, et al. Vitamin D: update 2013: from rickets prophylaxis to general preventive healthcare. *Dermatoendocrinol* 2013;5(3):331–47.
- [18] Straube S, Andrew Moore R, Derry S, et al. Vitamin D and chronic pain. *Pain* 2009;141(1–2):10–3.
- [19] Adams JS, Hewison M. Update in vitamin D. *J Clin Endocrinol Metab* 2010;95(2):471–8.
- [20] Pande RU, Chandrasekhar R, Kaplish N, et al. Low serum vitamin D concentration as a predictor of short sleep duration: a NHANES 2005–2006 analysis. *Sleep* 2009;32:A136.
- [21] Massa Jennifer, Stone Katie L, Wei Esther K, et al. Misti paudel, susan redline, sonia ancoli-Israel, eric orwoll, eva schernhammer; vitamin D and actigraphic sleep outcomes in older community-dwelling men: the MrOS sleep study. *Sleep* 1 February 2015;38(Issue 2):251–7.
- [22] Bertisch SM, Sillau S, de Boer IH, et al. 25-Hydroxyvitamin D concentration and sleep duration and continuity: multi-ethnic study of atherosclerosis. *Sleep* 2015;38(8):1305–11. <https://doi.org/10.5665/sleep.4914>.
- [23] Kim J, Lee W, Kweon H, et al. Associations between sleep duration and serum vitamin D level in Korean women. *Korean Journal of Family Practice* 2016;6(6):586–91.
- [24] Huang W, Shah S, Long Q, et al. Improvement of pain, sleep, and quality of life in chronic pain patients with vitamin D supplementation. *Clin J Pain* 2013;29(4):341–7.
- [25] Jung YS, Chae CH, Kim YO, et al. The relationship between serum vitamin D levels and sleep quality in fixed day indoor field workers in the electronics manufacturing industry in Korea. *Annals of Occupational and Environmental Medicine* 2017;29:25. <https://doi.org/10.1186/s40557-017-0187-7>.
- [26] Han B, Zhu F-X, Shi C, et al. Association between serum vitamin D levels and sleep disturbance in hemodialysis patients. *Nutrients* 2017;9(2):139. <https://doi.org/10.3390/nu9020139>.
- [27] Cheng TS, Loy SL, Cheung YB, et al. Plasma vitamin D deficiency is associated with poor sleep quality and night-time eating at mid-pregnancy in Singapore. *Nutrients* 2017;9(4):340. <https://doi.org/10.3390/nu9040340>.
- [28] Suzan Gunduz, Kosger H, Aldemir S, et al. Sleep deprivation in the last trimester of pregnancy and inadequate vitamin D: is there a relationship? *J Chin Med Assoc* 2016;79(1):34–8.
- [29] Mason C, De Dieu Tapsoba J, Duggan C, et al. Repletion of Vitamin D associated with deterioration of sleep quality among postmenopausal women. *Prev Med* 2016;93:166–70. <https://doi.org/10.1016/j.ypmed.2016.09.035>.
- [30] Stumpf WE, O'Brien LP. 1,25 (OH)₂ vitamin D₃ sites of action in the brain. An autoradiographic study. *Histochemistry* 1987;87:383–406.
- [31] Stumpf WE, Bidmon HJ, Li L, et al. Nuclear receptor sites for vitamin D-soltril in midbrain and hindbrain of Siberian hamster (*Phodopus sungorus*) assessed by autoradiography. *Histochemistry* 1992;98:155–64.
- [32] Musiol IM, Stumpf WE, Bidmon H-J, et al. Vitamin D-soltril nuclear binding to neurons of the septal, substriatal, and amygdaloid area in the Siberian hamster (*Phodopus sungorus*) brain. *Neuroscience* 1992;48:841–8.
- [33] Stumpf WE. Drug localization and targeting with receptor microscopic autoradiography. *J Pharmacol Toxicol Methods* 2005;51(1):25–40.
- [34] Eyles DW, Smith S, Kinobe R, et al. Distribution of the vitamin D receptor and 1 alpha-hydroxylase in human brain. *J Chem Neuroanat* 2005;29:21–30.
- [35] Gutierrez-Monreal MA, Duran RC-D, Moreno-Cuevas JE, et al. A role for 1; 25-dihydroxy vitamin D₃ in the expression of circadian genes. *J Biol Rhythm* 2014;29:384–8 [CrossRef] [PubMed].
- [36] Masood T, Kushwaha RS, Singh R, et al. Circadian rhythm of serum 25 (OH) vitamin D, calcium and phosphorus levels in the treatment and management of type-2 diabetic patients. *Drug Discov. Ther.* 2015;9:70–4 [CrossRef] [PubMed].
- [37] Vitaterna MH, Takahashi JS, Turek FW. Overview of circadian rhythms. *Alcohol Res Health* 2001;25:85–93 [PubMed].
- [38] Dibner C, Schibler U, Albrecht U. The mammalian circadian timing system: organization and coordination of central and peripheral clocks. *Annu Rev Physiol* 2010;72:517–49 [CrossRef] [PubMed].
- [39] Lucock M, Jones P, Martin C, et al. Vitamin D: beyond metabolism. *J. Evid. Based Complement. Altern. Med.* 2015;20:310–22 [CrossRef] [PubMed].
- [40] Garçon E, Wion-Barbot N, Montero-Menei CN, et al. New clues about vitamin D functions in the nervous system. *Trends Endocrinol Metab* 2002;13:100–5.
- [41] Balden R, Selvamani A, Sohrabji F. Vitamin D deficiency exacerbates experimental stroke injury and dysregulates ischemia-induced inflammation in adult rats. *Endocrinology* 2012;153:2420–35.
- [42] Zakon HH. The effects of steroid hormones on electrical activity of excitable cells. *Trends Neurosci* 1998;21(5):202–7.
- [43] Shipton EA, Shipton EE. Vitamin D and pain: vitamin D and its role in the aetiology and maintenance of chronic pain states and associated comorbidities. *Pain Res Treat* 2015;2015: 904967.
- [44] McGrath J, Feron F, Eyles D, Mackay-Sim A. Vitamin D: the neglected neurosteroid? *Trends Neurosci* 2001 Oct;24(10):570–2.
- [45] Veenstra TD, Prüfer K, Koeningberger C, et al. 1,25-Dihydroxyvitamin D₃ receptors in the central nervous system of the rat embryo. *Brain Res* 1998 Sep 7;804(2):193–205.
- [46] Muzur A, Pace-Schott EF, Hobson JA. The prefrontal cortex in sleep. *Trends Cognit Sci* 2002 Nov 1;6(11):475–81.
- [47] Macey PM, Henderson LA, Macey KE, et al. Brain morphology associated with obstructive sleep apnea. *Am J Respir Crit Care Med* 2002;166(10):1382–7.
- [48] Cameron HA, McKay RD. Adult neurogenesis produces a large pool of new granule cells in the dentate gyrus. *J Comp Neurol* 2001;435(4):406–17.
- [49] Guzmán-Marín R, Suntsova N, Stewart DR, et al. Sleep deprivation reduces proliferation of cells in the dentate gyrus of the hippocampus in rats. *J Physiol* 2003 Jun 1;549(Pt 2):563–71.
- [50] Stoffers D, Altena E, van der Werf YD, et al. The caudate: a key node in the neuronal network imbalance of insomnia? *Brain* 2014 Feb;137(Pt 2):610–20.
- [51] Marks GA, Roffwarg HP, Shaffery JP. Neuronal activity in the lateral geniculate nucleus associated with Ponto-geniculo-occipital waves lacks lamina specificity. *Brain Res* 1999 Jan 2;815(1):21–8.
- [52] Lima Marcelo MS, Andersen Monica L, Reksidler Angela B, et al. The role of the substantia nigra pars compacta in regulating sleep patterns in rats. *PLoS One* 2007;2(6):e513.
- [53] Iranzo A, Lomeña F, Stockner H, et al. Sleep Innsbruck Barcelona (SINBAR) group. Decreased striatal dopamine transporter uptake and substantia nigra hyperchogenicity as risk markers of synucleinopathy in patients with idiopathic rapid-eye-movement sleep behaviour disorder: a prospective study [corrected]. *Lancet Neurol* 2010 Nov;9(11):1070–7.
- [54] McKinley MJ, Yao ST, Uschakov A, et al. The median preoptic nucleus: front and centre for the regulation of body fluid, sodium, temperature, sleep and cardiovascular homeostasis. *Acta Physiol (Oxf)*. 2015 May;214(1):8–32.
- [55] Lu J, Greco MA, Shiromani P, et al. Effect of lesions of the ventrolateral preoptic nucleus on NREM and REM sleep. *J Neurosci* 2000 May 15;20(10):3830–42.
- [56] Jirikowski GF, Kaunzner UW, Dief Ael E, et al. Distribution of vitamin D binding protein expressing neurons in the rat hypothalamus. *Histochem Cell Biol* 2009 Mar;131(3):365–70.
- [57] Born J, Kellner C, Uthgenannt D, et al. Vasopressin regulates human sleep by reducing rapid-eye-movement sleep. *Am J Physiol* 1992 Mar;262(3 Pt1):E295–300.
- [58] Kaneko I, Sabir MS, Dussik CM, et al. 1,25-Dihydroxyvitamin D regulates expression of the tryptophan hydroxylase 2 and leptin genes: implication for behavioral influences of vitamin D. *Faseb J* 2015 Sep;29(9):4023–35.
- [59] Patrick RP, Ames BN. Vitamin D and the omega-3 fatty acids control serotonin synthesis and action, part 2: relevance for ADHD, bipolar disorder, schizophrenia, and impulsive behavior. *Faseb J* 2015 Jun;29(6):2207–22.
- [60] Golan D, Staun-Ram E, Glass-Marmor L, et al. The influence of vitamin D supplementation on melatonin status in patients with multiple sclerosis. *Brain Behav Immun* 2013 Aug;32:180–5.