



Original article

Association between Mediterranean diet and hand grip strength in older adult women



Luigi Barrea^{a,*}, Giovanna Muscogiuri^a, Carolina Di Somma^b, Giovanni Tramontano^c, Vincenzo De Luca^c, Maddalena Illario^d, Annamaria Colao^a, Silvia Savastano^a

^a Dipartimento di Medicina Clinica e Chirurgia, Unit of Endocrinology, Federico II University Medical School of Naples, Via Sergio Pansini 5, 80131 Naples, Italy

^b IRCCS SDN, Napoli Via Gianturco 113, 80143, Naples, Italy

^c Research and Development Unit, Federico II University Hospital, Via Sergio Pansini 5, 80131 Naples, Italy

^d Health Innovation Division, Campania Region, Italy

ARTICLE INFO

Article history:

Received 7 September 2017

Accepted 17 March 2018

Keywords:

Environmental factors
Mediterranean diet
Hand grip strength
Elderly
Nutritionist

SUMMARY

Background & aims: Mediterranean Diet (MD) is an eating pattern associated with multiple healthy benefits, including the conservation of skeletal muscle. Frailty is a major geriatric syndrome characterized by low muscle strength. The Hand Grip Strength (HGS) is the most frequently used indicator of muscle functional capacity for clinical purposes. The association between the adherence to the MD and HGS in elderly has not yet fully investigated. The goal of this study was to examine the association between the adherence to the MD and HGS in a not hospitalized elderly who participated in the project **PERSONALISED ict Supported Services for Independent Living and Active Ageing (PERSSILAA)**.

Methods: Eighty-four elderly women were consecutively enrolled (aged 60–85 years) in this cross-sectional observational study. Anthropometric measures were evaluated. A validated 14-item questionnaire PREDIMED (PREvención con Dieta MEDiterránea) was used for the assessment of adherence to the MD. Dietary data were collected by a 7-day food records. Muscle strength was measured by HGS using a grip strength dynamometer (KERN & SOHN GmbH).

Results: The majority of participants were overweight (46.4%). An average adherence to the MD was found in 52.4% of participants, while the minority of them showed a low adherence (21.4%). HGS > cut-point of 20 kg were found in 43 subjects (51.2%). According to the adherence to MD, 39% participants with HGS values higher than cut-point presented a high adherence score compared with 14% of those with lower values of HGS ($p = 0.018$). The participants with HGS > cut-point presented significantly higher PREDIMED score than those with HGS < cut-point ($p < 0.001$). Based on ROC curves, the most sensitive and specific cut-point for the PREDIMED score to predict HGS categories was ≥ 8 . No evident correlations were observed between HGS and age, while HGS was negatively correlated with hip circumference ($r = -0.233$, $p = 0.033$) and BMI ($r = -0.219$, $p = 0.045$), and positively correlated with PREDIMED score ($r = 0.598$, $p < 0.001$). At binomial logistic regression analysis almost all 14-items of PREDIMED questionnaire were significantly associated with HGS adjusted for BMI. At multinomial logistic regression analysis to assess the association of the three classes of adherence to the MD with the HGS, after adjusting for BMI the lowest adherence to MD was associated with the lowest Odds Ratio of HGS ($p < 0.001$).

Conclusions: This study evidenced a positive association between the adherence to the MD and muscle strength in a sample of active elderly women, stratified according to the HGS > cut-point of 20 kg. Our study highlights the usefulness of the developing health services to detect and prevent age-associated decline in physical performance in elderly subjects by addressing nutritional and physical intervention.

© 2018 Elsevier Ltd and European Society for Clinical Nutrition and Metabolism. All rights reserved.

Abbreviations: MD, Mediterranean Diet; HGS, Hand Grip Strength; PERSSILAA, PERSONALISED ict Supported Services for Independent Living and Active Ageing; BMI, Body Mass Index; WC, Waist Circumference; NCEP-ATP III, According to the National Cholesterol Education Program's Adult Treatment Panel III; HC, Hip circumference; WHR, Waist to Hip Ratio; PREDIMED, PREvención con Dieta MEDiterránea; FFQ, Food Frequency Questionnaire; LEP, Limb Extension Power; SPPB, Short Physical Performance Battery.

* Corresponding author. Fax +39 081 746 3668.

E-mail addresses: luigi.barrea@unina.it (L. Barrea), giovanna.muscogiuri@gmail.com (G. Muscogiuri), cdisomma@unina.it (C. Di Somma), giovanni.tramontano@libero.it (G. Tramontano), vinc.deluca@gmail.com (V. De Luca), illarior@unina.it (M. Illario), colao@unina.it (A. Colao), sisavast@unina.it (S. Savastano).

<https://doi.org/10.1016/j.clnu.2018.03.012>

0261-5614/© 2018 Elsevier Ltd and European Society for Clinical Nutrition and Metabolism. All rights reserved.

1. Introduction

Nutrition is one of the major modifiable environmental risk factors for several non-communicable diseases and represents a mainstay for their prevention and also their initial treatment [1]. In particular, in the elderly population nutrition is an important element of health and there is increasing scientific and clinical evidence showing the link between nutrition and health as part of aging [2,3].

The Mediterranean diet (MD) is a plant-based, antioxidant-rich diet known for its several health benefits. This dietary pattern is considered a nutritionally adequate, complete and easy to follow, as it is based on the traditional foods that people used to eat in their countries [4]. In particular, the MD features a high intake of whole grains, fruits, vegetables, tree nuts, legumes, and olive oil on a daily basis; a moderate intake of fish and poultry, low consumption of dairy products, red meat, processed meat and sweets, and moderate consumption of wine with meals [5,6]. Large observational prospective epidemiological studies support that the Mediterranean dietary pattern increases life expectancy, reduce the risk of major chronic diseases, and improve quality of life and well-being [7,8]. However, it should be considered that the Mediterranean dietary patterns may vary according to age, gender, ethnicity, culture and other lifestyle factors [9]. Indeed, previous studies have identified gender as a key determinant of food choices [10] with females having more motivation towards healthy eating and are more aware of healthy eating compared with males [11,12]. In elderly subjects numerous studies have reported that higher adherence to the MD is associated with lowest mortality [13–16], although most of these studies were not conducted specifically to determine the MD effects in elderly subjects. Higher adherence to the MD has been also proven to be effective in preserving the skeletal muscle mass in healthy women likely due to the potential anti-inflammatory and anti-oxidant properties of micronutrients (e.g. carotenoids and vitamins C and E) or through their direct role in muscle metabolism and physiology, such as with magnesium and potassium [17].

Nutrition plays a decisive role in the development of frailty [18,19]. Frailty or “the frailty syndrome” are commonly used terms to denote a clinical entity in elderly subjects. Frailty has been reported to be associated with increased risk for adverse outcomes, such as onset of disability, morbidity, institutionalization or mortality [20]. In particular, the presence of frailty has been associated with increased all-cause mortality [21–24] and increased incident cardiovascular disease [25], as well as poor survival after cardiac and surgical procedures. The effectiveness of the adherence to a Mediterranean dietary pattern on the risk of development of frailty in elderly adults has been demonstrated in different clinical settings. In particular, using the MD score, a high adherence to a MD was associated with a slower decline of mobility and to the development of frailty over time in community-dwelling elderly persons of both gender participating to the InCHIANTI Study [26,27].

The Hand Grip Strength (HGS) is an objective component of the frailty syndrome in the elderly subjects, thereby representing the most frequently used indicator of muscle functional capacity for clinical purposes [28]. HGS is a non-invasive and reliable method for assessment of muscle power and nutritional status and portable HGS devices are quick and easy to use [29]. The association between nutrition status and HGS is well documented [30] and HGS is currently considered as a marker of the nutrition status [30,31]. In particular, HGS reflects early nutrition deprivation and nutrition repletion also before changes in body composition parameters can be detected [30]. The Academy of Nutrition and Dietetics and the American Society for Parenteral and Enteral Nutrition has recently recommended reduced HGS values as a criterion for the

identification and documentation of undernutrition in clinical practice [31]. Besides nutrition, other parameters, such as sex and age, have also been identified to be associated with HGS in elderly subjects [32–36]. HGS is associated with increased all-cause mortality [37], disability and increased length of hospital stay [38]. Thus, it is essential for public health to implement screenings and multidisciplinary treatments of frailty, especially through indirect but reliable measures, such as HGS.

This cross-sectional observational study was designed to capture the association between a Mediterranean dietary pattern with HGS in elderly women participants in the PERSONALISED ICT SUPPORTED SERVICES FOR INDEPENDENT LIVING AND ACTIVE AGEING (PERS-SILAA), an European project developing health services to detect and prevent frailty in elderly subjects by addressing cognitive, physical and nutritional domains in Campania Region, Italy and Enschede, the Netherlands [39].

2. Subjects and methods

2.1. Design and setting

PERSILAA is an European project developing health services to detect and prevent frailty in elderly subjects by addressing cognitive, physical and nutritional domains in Campania Region, Italy and Enschede, the Netherlands [39]. In particular, PERSILAA is a Community-Based, Technology-Supported Service Model for Detecting and Preventing Frailty in elderly [40] and the evaluation of the association between MD and HGS is based only from the Italian population data. In Italy, elderly are invited for the first screening *via* the church, after which they can complete this screening at the church with the help of a trained volunteer or online. As elderly subjects in Italy frequently visit the church, where many activities take place, this appeared to be the best way to reach them. This community based healthcare service has been provided in Campania Region, in collaboration with the Department of Gastroenterology, Endocrinology and Surgery of Federico II University Hospital (Italy). The work has been carried out in accordance with the Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans, and it has been approved by the Ethical Committee of the University of Naples “Federico II” Medical School (n. 178/14). The purpose of the protocol was explained to all elderly subjects enrolled, and written informed consent was obtained.

2.2. Population study

Our study population consisted of 183 residents of the Campania Region, Italy, who were healthy elderly with active lifestyles (age ≥ 60 years), and who were enrolled at local catholic churches in the same district of Naples and were involved in the first screening of PERSILAA project, from October 2016 to March 2017. All potential subjects were invited to participate in a full medical history, including cognitive testing (Mini-Mental State Examination (MMSE) and AD8 Informant Questionnaire for Cognitive Impairment), physical performance (Short Physical Performance Battery), pharmacological history, physical activity, smoking habits, and a complete clinical examination. Criteria for exclusion from the study were:

- a. Male individuals;
- b. Absence of mild-severe cognitive impairment;
- c. Severe-moderate limitation of physical performance;
- d. Individuals who needed assistance with their daily activities;
- e. Subject unable to perform the HGS test, defined as situations that led to an inability in understanding verbal instructions or

having clinical conditions that could influence performing the HGS technique correctly, including osteoarticular or musculoskeletal diseases, pain, confusion, and moderate/severe neurological and/or cognitive impairment, stroke or hand injury, cancer, acute or chronic inflammatory diseases, endocrine diseases, in particular altered thyroid hormone function tests;

- f. Underweight (BMI \leq 20.0 kg/m²) or with obesity grade II/III (BMI \geq 35.0 kg/m²);
- g. Hypocaloric diet in the last three months or specific nutritional regimens, including vegan, vegetarian hyperproteic or ketogenic diets;
- h. Vitamin/mineral antioxidant or protein supplementation;
- i. Individuals occasionally or currently taking medications that could influence musculoskeletal system and hand grip strength (i.e. anticonvulsivants and psychotropic drugs, steroidal and non-steroidal anti-inflammatory drugs, hormone replacement therapy, anti-thyroid agents, weight-loss medications, dopaminergic drugs, Parkinson's disease drugs, or statins);
- j. Alcohol abuse according to the Diagnostic and Statistical Manual of Mental Disorders (DSM)-V diagnostic criteria.

On the basis of these criteria, 99 subjects were excluded and the final study population consisted of 84 women.

2.3. Anthropometric measurements

The measurements were performed between 8 and 11 AM and were made in a standard way by one operator (a nutritionist experienced in providing nutritional assessment). All anthropometric measurements were recorded with subjects wearing only light clothes and without shoes. In each subject, weight and height were assessed in order to calculate the Body Mass Index (BMI) [weight (kg) divided by height squared (m²), kg/m²]. Height was measured to the nearest 0.5 cm using a wall-mounted stadiometer (Seca 711; Seca, Hamburg, Germany). Body weight was determined to the nearest 0.1 kg using a calibrated balance beam scale (Seca 711; Seca, Hamburg, Germany). BMI was classified according to WHO's criteria with normal weight: 18.5–24.9 kg/m²; overweight, 25.0–29.9 kg/m² and grade I obesity, 30.0–34.9 kg/m². Waist Circumference (WC) was measured to the nearest 0.1 cm with a non-stretchable measuring tape, at the end of several consecutive natural breaths, at the level parallel to the floor, midpoint between the top of the iliac crest and the lower margin of the last palpable rib in midaxillary line [41]. According to the National Cholesterol Education Program's Adult Treatment Panel III (NCEP-ATP III) criteria, abdominal obesity was defined as WC \geq 88 cm in women [42]. Hip Circumference (HC) was measured as the maximum circumference around the buttocks posteriorly and the symphysis pubis anteriorly, and measured to the nearest 0.5 cm. Waist to Hip Ratio (WHR) was calculated by dividing WC (in cm) by HC (cm). The cutoffs points (Caucasian) of WHR used ($>$ 0.85 in women) [43] denote abdominal obesity. Average of two readings was used for analysis.

2.4. Adherence to the MD

The adherence to the MD was assessed using the previously validated 14-item questionnaire for the assessment of PREvencción con Dieta MEDiterránea (PREDIMED) [44]. A qualified Nutritionist provided the questionnaire during a face-to-face interview to all the subjects included in the study. Briefly, score 1 and 0 was assigned for each items; PREDIMED score was calculated as follows: 0–5, lowest adherence; score 6–9, average adherence; score \geq 10, highest adherence [44].

2.5. Dietary assessment

Data were obtained during a face-to-face interview between the patient and a qualified Nutritionist. The dietary interview was used in order to quantify foods and drinks by using a photographic food atlas (\approx 1000 photographs) of known portion sizes and to be sure of accurate completion of the records (21). Moreover, dietary data, including beverage intakes consumption, were collected by a 7-day food records. The subjects returned the records to the nutritionist who asked supplemental questions if necessary. Data were stored and processed using a commercial software (Terapia Alimentare Dietosystem[®] DS-Medica, <http://www.dsmedica.info>). Considering quantities and qualities of foods consumed, the software is able to calculate not only the daily caloric intake but also the quantities of macronutrients (protein; total, complex and simple carbohydrates; total fat, saturated fat (SFA)), unsaturated fat (Monounsaturated Fatty Acids, MUFA; Polyunsaturated Fatty Acids, PUFA: n-6 PUFA, n-3 PUFA and n-6/n-3 PUFAs ratio; cholesterol and fibers).

2.6. Hand grip strength

HGS (kg) was quantified by measuring the amount of static force that the hand and forearm can squeeze around a hand-held dynamometer [45] using a Collins dynamometer (KERN & SOHN GmbH). All measurements were performed under strictly standardized conditions by the operator, using the same device in order to avoid inter-observer and inter-device variability, after explaining the procedure to each patient. The participants were seated, their elbow by their side and flexed to right angles, and a neutral wrist position [46] for 3 s. The maximum value of 3 consecutive measurements in the non-dominant arm was registered. Patients used their dominant hand when they were unable to perform HGS with their non-dominant hand. Brief pauses were taken between measurements. The maximum value (kg) out of three trials using the dominant hand was recorded. Between two consecutive trials, a 1-min recovery was provided. The instrument was routinely checked with resistors and capacitors of known values. The coefficient of variation (CV) of repeated measurements of HGS was assessed in 10 patients and resulted 1.8%. Cut-point used for low HGS in women was $<$ 20 kg [47].

2.7. Statistical analysis

The minimum required total sample was calculated using the pooled standard deviation with the level test 0.05 and power 80% of means of the measurements of HGS. With a type I error of 0.05 and a type of II (β) error of 0.10, the resulting size was 84 women.

Results are expressed as mean \pm SD or as median plus range, according to the variable distributions evaluated by Kolmogorov–Smirnov test ($p <$ 0.01).

1. The chi square (χ^2) test was used to determine the significance of differences in response frequency of dietary components included in the PREDIMED questionnaire and in PREDIMED categories (low adherence, average adherence and high adherence to the MD) in the study population grouped according to HGS values dichotomized in two categories, above and below the cut-point (20 kg).
2. Receiver operator characteristic (ROC) curve analysis was performed to determine sensitivity and specificity, area under the curve (AUC), and Interval Confidence (CI), as well as cut-point for PREDIMED score in detecting HGS cut-point ($<$ 20 kg and $>$ 20 kg) in the participants. Test AUC for ROC analysis was also performed. We want show that AUC resulted 0.752 for a particular test is significant from the null hypothesis value 0.5

(meaning no discriminating power), than we enter 0.957 for AUC ROC and 0.5 for null hypothesis values. For α level we selected 0.05 type I error and for β level we selected 0.20 type II error.

- Differences in PREDIMED score, total energy and daily nutrients intake of participants, and HGS values dichotomized in two categories, above and below the cut-point (20 kg) among participants, were analyzed by unpaired Student's *t* test or Wilcoxon signed-rank test, when appropriate.
- Bivariate proportional odds ratio (OR) models, 95% IC, and R^2 , were performed to assess the association among quantitative variables (single items of PREDIMED questionnaire) with HGS, after adjusting for BMI.
- Multinomial logistic regression, 95% IC, and R^2 , was performed to model the relationship between the HGS and the three categories of PREDIMED score (low adherence, average adherence and high adherence to the MD).
- The correlations between study variables were performed using Pearson *r* or Spearman's *rho* correlation coefficients according to the variable's distribution.
- A multiple linear regression analysis model (stepwise method), expressed as R^2 , Beta (β) and *t*, with HGS as dependent variables was used to estimate the predictive value of food items included in the PREDIMED, the score of adherence to the MD, and the daily nutrients intake (protein, fat, unsaturated fat, PUFA, n-3 PUFA and cholesterol).

In these analyses, we entered only those variables that had a *p* value < 0.05 in the univariate analysis (partial correlation). To avoid multicollinearity, variables with a variance inflation factor (VIP) > 10 were excluded. Values $\leq 5\%$ were considered statistically significant. Data were stored and analyzed using the MedCalc® package (Version 12.3.0 1993–2012 MedCalc Software bvba – MedCalc Software, Mariakerke, Belgium).

3. Results

Study population consisted of 84 elderly women, aged 60–85 years (mean age 71.7 ± 5.5 yrs), with 20 participants (24%) with age ≥ 75 yrs. Concerning their marital status and education status, the vast majority of these subjects were married (n.43; 51%) or widowed (n.36; 43%) with middle school diploma (n.95; 94%), while the rest 5% of them were single or have higher school diploma. Eighteen subjects were normal weight (21.4%), 39 were overweight (46.4%), and 27 presented grade I obesity (32.1%). The PREDIMED questionnaires, the 7-day food records, and HGS were assessed in all participants. All participants were retired and declared to have never smoked. No participants engaged in any regular physical activity for more than once a week in the 6 months preceding the beginning of the study or in mild-moderate daily physical activities, like walking or gardening. WC and WHR > cut-off values were found in 51 subjects (60.7%) and 54 subjects (64.3%), respectively. According to PREDIMED score, the majority of participants reached an average adherence to the MD (52.4%), while the minority of them showed a low adherence (21.4%). HGS > cut-point values were found in 43 subjects (51.2%).

In [Supplementary file S1](#) we reported the responses of each item included in PREDIMED questionnaire. Extra-virgin olive oil was the most consumed food item, followed by vegetables. The participants were grouped according to the responses to each item included in PREDIMED questionnaire and to the HGS ([Table 1](#)). As showed in the table, the participants with HGS values higher than cut-point exhibited significant differences compared with those with higher values for the intake of the following dietary components: soda drinks ($p = 0.003$), legumes ($p = 0.045$) and commercial

Table 1

Response frequency of dietary components included in the PREDIMED questionnaire according to amount of cut-off points of HGS.

	Questions PREDIMED questionnaire		HGS < 20		HGS > 20		χ^2	<i>p</i> values
			n = 43		n = 41			
			n	%	n	%		
1	Use of extra-virgin olive oil as main culinary lipid		40	93.0	40	97.6	0.22	0.643
2	Extra virgin olive oil >4 tablespoons		34	79.1	35	85.4	0.22	0.639
3	Vegetables ≥ 2 servings/day		26	60.5	32	78.0	2.27	0.132
4	Fruits ≥ 3 servings/day		26	60.5	30	73.2	1.01	0.316
5	Red/processed meats <1/day		25	58.1	31	75.6	2.15	0.143
6	Butter, cream, margarine <1/day		20	46.5	27	65.9	2.45	0.118
7	Soda drinks <1/day		18	41.9	31	75.6	8.50	0.003
8	Wine glasses ≥ 7 /week		18	41.9	21	51.2	0.41	0.522
9	Legumes ≥ 3 /week		19	44.2	28	68.3	4.02	0.045
10	Fish/seafood ≥ 3 /week		18	41.9	23	56.1	1.18	0.277
11	Commercial sweets and confectionery ≤ 2 /week		15	34.9	25	61.0	4.73	0.029
12	Tree nuts ≥ 3 /week		7	16.3	11	26.8	0.83	0.362
13	Poultry more than red meats		24	55.8	24	58.5	0.01	0.975
14	Use of sofrito sauce ≥ 2 /week		11	25.6	15	36.6	0.73	0.393
PREDIMED score								
	Low – adherence		14	32.6	4	9.8	5.20	0.022
	Average – adherence		23	53.5	21	51.2	0.01	0.992
	High – adherence		6	14.0	16	39.0	5.59	0.018

Differences in the intake of single items of the PREDIMED score according to HGS values in the study participants. Results are expressed as number and percentage of responses obtained with PREDIMED questionnaire. The differences were analyzed by χ^2 test. A *p* value in bold type denotes a significant difference ($p < 0.05$). Cut-point used for low HGS in women was <20 kg [47].

PREDIMED, PREvención con DietaMEDiterránea; **HGS**, Hand Grip Strength.

sweets and confectionery ($p = 0.029$). Considering the adherence to MD, 39% participants with HGS values higher than cut-point presented a high adherence score compared with 14% of those with lower values of HGS ($p = 0.018$). Accordingly, participants with HGS > cut-point presented significantly PREDIMED score than those with HGS < cut-point ($p < 0.001$) ([Fig. 1](#)).

ROC analysis for predictive values of the adherence to MD in detecting the HGS categories, was reported in [Fig. 2](#). The number of cases required was set at 30, from the AUC 0.752. Based on ROC curves, the most sensitive and specific cut-point for the PREDIMED score to predict HGS categories was ≥ 8 (sensitivity 63.4, specificity 81.4; $p < 0.001$). In [Table 2](#) we reported the total energy and the daily nutrients intake obtained from the 7-day food records. As showed in table, the participants with HGS < cut-point had significantly lower percentage of energy from protein ($p < 0.001$),




Fig. 1. Differences in PREDIMED score between the two groups of participants stratified according to the HGS cut-point. Participants with HGS > cut-point presented significantly higher adherence to MD than those with HGS < cut-point. Results are expressed as mean \pm SD according to the variable distributions evaluated by Kolmogorov–Smirnov test. The unpaired *t*-test was used to test the significance of differences between the two groups. A *p* value in bold type denotes a significant difference ($p < 0.05$). Cut-point used for low HGS in women was <20 kg [47]. **PREDIMED**, PREvención con Dieta MEDiterránea; **MD**, Mediterranean Diet; **HGS**, Hand Grip Strength.




Fig. 2. ROC for predictive values of PREDIMED score in detecting HGS values. The most sensitive and specific cut-off of the PREDIMED score to predict HGS cut-off point (20 Kg), was ≥ 8 (sensitivity 63.4, specificity 81.4; $p < 0.001$). A p value in bold type denotes a significant difference ($p < 0.05$). Cut-point used for low HGS in women was < 20 kg [47]. **ROC**, Receiver operating characteristic analysis; **PREDIMED**, PREVENCIÓN con Dieta MEDiterránea; **HGS**, Hand Grip Strength; **AUC**, Area Under Curve.

total carbohydrate ($p < 0.001$), unsaturated fat ($p = 0.018$) and n-3 PUFA ($p = 0.031$), and a significantly higher total fat ($p < 0.001$) and cholesterol intake ($p = 0.006$) than HGS $>$ cut-point.

3.1. Correlation studies

No evident correlations were observed between HGS and age. Among anthropometric measures, HGS was negatively correlated with HC ($r = -0.233$, $p = 0.033$) and BMI ($r = -0.219$, $p = 0.045$); HGS was also positively correlated with PREDIMED score, independently of BMI (Fig. 3). The results of binomial logistic regression model performed to assess the association of dietary components included in the PREDIMED questionnaire with HGS, adjusted for

BMI, were reported in Table 3. The traditional foods of the MD, when consumed more frequently, were significantly associated with HGS.

The results of multinomial logistic regression model to assess the association of the three classes of PREDIMED score with the HGS, adjusted for BMI, were reported in Table 4. The lowest adherence to MD was associated with the lowest OR of HGS ($p < 0.001$).

In Supplementary file S2 we reported the correlations among HGS, total energy and daily nutrients intake, evaluated by using the 7-day food records. HGS correlated with some of the dietary nutrients intake, in particular protein ($p < 0.001$), fat ($p < 0.001$), unsaturated fat ($p = 0.008$), PUFA ($p = 0.015$), n-3 PUFA ($p = 0.027$), cholesterol ($p = 0.014$), also after adjusting for total energy intake and BMI. In addition, the correlation between PREDIMED score and HGS was independent of protein intake ($r = 0.472$; $p < 0.001$).

To compare the relative predictive power of the food items included in the PREDIMED, the score of adherence to the MD, and the daily nutrients intake associated with the HGS, we performed a multiple linear regression analysis using models that included as measures the consumption frequency of each food item of PREDIMED questionnaire, the PREDIMED score along with the daily nutrients intake (protein, fat, unsaturated fat, PUFA, n-3 PUFA and cholesterol). Using these model PREDIMED score entered at the first step ($p < 0.001$), followed by the protein intake ($p < 0.001$); results were reported in Table 5.

4. Discussion

In this cross-sectional observational study we evaluated the association between the adherence to the MD, using the PREDIMED score, and the HGS in a sample of active elderly women. The novel finding of this study is the association between adherence to the MD and its individual foods with the HGS. In particular, when grouping the study participants according to the mean reference value for HGS, defined as 20 kg in women, we observed that subjects with HGS above the cut-point presented lower intake of energy-dense nutrient-poor “junk food” and soda drinks, higher consumption of legumes, and higher adherence to the MD when compared to their counterpart with HGS below the cut-point. By using ROC analysis, a PREDIMED score ≥ 8 was found as the most sensitive and specific cut-point to predict value of HGS above the

Table 2
Total energy and daily nutrients intake of participants grouped on the basis of HGS cut-point.

Parameters	Total n = 84	HGS < 20 n = 43	HGS > 20 n = 41	p-value*
Total energy (kcal)	2032.52 \pm 257.00	2040.01 \pm 273.88	2024.68 \pm 241.19	0.787
Protein (% of total kcal)	13.47 \pm 2.17	12.24 \pm 2.04	14.75 \pm 1.45	<0.001
Carbohydrate (% of total kcal)	55.05 (50.90–61.90)	55.10 (50.91–60.0)	56.00 (51.00–61.90)	<0.001
Complex (% of total kcal)	30.80 (21.01–39.32)	30.61 (23.70–39.30)	30.60 (21.00–38.20)	0.788
Simple (% of total kcal)	25.60 (16.40–37.70)	25.70 (18.70–35.60)	25.40 (16.40–37.70)	0.957
Fat (% of total kcal)	30.95 \pm 3.61	32.34 \pm 3.38	29.50 \pm 3.27	<0.001
SFA (% of total kcal)	9.07 \pm 1.46	9.24 \pm 1.62	8.89 \pm 1.28	0.271
Unsaturated fat (% of total kcal)	21.89 \pm 3.65	20.98 \pm 3.96	22.83 \pm 3.05	0.018
MUFA (% of total kcal)	15.75 (10.01–22.14)	15.69 (10.21–22.14)	15.95 (10.01–20.70)	0.405
PUFA (% of total kcal)	13.41 \pm 8.96	5.21 \pm 4.25	6.67 \pm 2.99	0.070
n-6 PUFA (g/day)	8.52 \pm 8.31	7.77 \pm 9.77	9.30 \pm 6.47	0.398
n-3 PUFA (g/day)	4.90 \pm 2.71	4.28 \pm 2.85	5.54 \pm 2.42	0.031
n-6/n-3 PUFAs ratio	2.35 \pm 3.37	2.56 \pm 4.33	2.14 \pm 1.95	0.563
Cholesterol (mg/day)	321.37 \pm 38.10	332.42 \pm 34.91	309.78 \pm 38.24	0.006
Fiber (g/day)	18.54 \pm 5.36	18.00 \pm 5.69	19.09 \pm 4.99	0.357

Differences in total energy and daily nutrients intake of participants grouped on the basis of HGS cut-point. Results are expressed as mean \pm standard deviation or as median plus range according to variable distributions evaluated by Kolmogorov–Smirnov test. Differences between groups were analyzed by unpaired Student’s t test or Wilcoxon signed-rank test, when appropriate. * p -value indicates comparison between those with low HGS $<$ cut-point versus those with HGS $>$ cut-point. A p value in bold type denotes a significant difference ($p < 0.05$).

HGS, Hand Grip Strength; **SFA**, Saturated Fatty Acids; **MUFA**, MonoUnsaturated Fatty Acids; **PUFA**, PolyUnsaturated Fatty Acids.




Fig. 3. Correlation between PREDIMED score and HGS, adjusted for BMI. A significant positive correlation was observed between PREDIMED score and HGS, independently of BMI. A *p* value in bold type denotes a significant difference ($p < 0.05$). **PREDIMED**, PREvención con Dieta MEDiterránea; **HGS**, Hand Grip Strength, **BMI**, Body Mass Index.

Table 3
Binomial Logistic Regression model to assess the association of dietary components included in the PREDIMED questionnaire with the HGS, after adjusting for BMI.

Questions	OR	95% IC	R ²	<i>p</i> value
1 Use of extra virgin olive oil as main culinary lipid	1.59	1.11–2.26	0.13	0.011
2 Extra virgin olive oil >4 tablespoons	1.09	0.98–1.21	0.03	0.125
3 Vegetables ≥2 servings/day	1.12	1.02–1.23	0.08	0.020
4 Fruits ≥3 servings/day	1.11	1.01–1.21	0.07	0.024
5 Red/processed meats <1/day	1.17	1.07–1.30	0.13	0.003
6 Butter, cream, margarine <1/day	1.12	1.03–1.23	0.09	0.008
7 Soda drinks <1/day	1.16	1.06–1.28	0.14	0.002
8 Wine glasses ≥7/week	1.08	0.98–1.14	0.03	0.135
9 Legumes ≥3/week	1.07	0.99–1.16	0.04	0.077
10 Fish/seafood ≥3/week	1.01	0.94–1.08	0.01	0.756
11 Commercial sweets and confectionery ≤2/week	1.11	1.02–1.20	0.08	0.015
12 Tree nuts ≥3/week	1.09	1.00–1.19	0.05	0.041
13 Poultry more than red meats	1.06	0.98–1.14	0.03	0.149
14 Use of sofrito sauce ≥2/week	1.04	0.96–1.12	0.01	0.320

Higher values of HGS were positively associated with a higher consumption of almost all of the Mediterranean food items, after adjusting for BMI. A *p* value in bold type denotes a significant difference ($p < 0.05$).

PREDIMED, PREvención con Dieta MEDiterránea; **HGS**, Hand Grip Strength; **BMI**, Body Mass Index; **ODDs**, Odds; **IC**, Interval Confidence.

Table 4
Multinomial Logistic Regression model to assess the association of the three classes of PREDIMED score with the HGS, after adjusting for BMI.

Adherence of MD	OR	95% IC	R ²	<i>p</i> value
Low adherence of MD	0.73	0.61–0.86	0.24	<0.001
Average adherence of MD	1.02	0.95–1.09	0.01	0.611
High adherence	1.14	1.04–1.25	0.11	0.003

The lowest adherence to MD was associated with the lowest OR of HGS, after adjusting for BMI. A *p* value in bold type denotes a significant difference ($p < 0.05$). **PREDIMED**, PREvención con Dieta MEDiterránea; **HGS**, Hand Grip Strength; **BMI**, Body Mass Index; **MD**, Mediterranean Diet; **ODDs**, Odds; **IC**, Interval Confidence.

mean reference values. To the best of our knowledge, no previous studies provided a specific cut-point for the PREDIMED score to predict the highest HGS values.

In this study, we took advantage of two easy and reliable tools to evaluate the adherence to the MD, on the one side, and the muscle functional capacity on the other side. The quantitative score (14-item) of adherence to the MD PREDIMED score is a key element

Table 5

Multiple regression analysis models (stepwise method) with the HGS as dependent variable to estimate the predictive value of a food items of PREDIMED questionnaire, PREDIMED score and nutrients intake (protein, fat, unsaturated fat, PUFA, n-3 PUFA and cholesterol).

Parameters	Multiple Regression analysis			
	R ²	β	t	<i>p</i> value
PREDIMED SCORE	0.358	0.598	6.8	<0.001
Protein intake (% of total kcal)	0.474	0.432	4.8	<0.001

Variable excluded: Other items of PREDIMED score; fat, unsaturated fat, PUFA, n-3 PUFA and cholesterol.

Among adherence to the MD, single food items of PREDIMED questionnaire, and among dietary nutrients intake, HGS were well predicted by PREDIMED score and protein intake. A *p* value in bold type denotes a significant difference ($p < 0.05$).

HGS, Hand Grip Strength; **PREDIMED**, PREvención con Dieta MEDiterránea; **PUFA**, Polyunsaturated fatty acids.

in the interventions conducted in the PREDIMED trials and that has been previously validated against the full-length Food Frequency Questionnaire (FFQ). PREDIMED is less time-demanding, less expensive and requires less collaboration from participants than the usual FFQ or other more comprehensive methods [44], characteristics that make this questionnaire particularly suitable for elderly population. Moreover, PREDIMED questionnaire allows providing feedback to the participant immediately after the interview is completed under the control of an expert Nutritionist. HGS evaluates the maximum isometric strength of the hand and forearm muscles and is commonly used in clinical as well as research settings [48]. As the strength of the forearm muscles does not necessarily represent the strength of other muscle groups, the validity of this test as a measure of general strength has been questioned [45], particularly among obese women [49]. However, results of the InCHIANTI study demonstrated that HGS equally well identified subjects with poor mobility as did lower extremity muscle power and knee extension torque, and thresholds of 30 kg for men and 20 kg for women were recommended for use in clinical practice [50]. Thus, when standardized methods and calibrated equipment are used, HGS represents a reliable and inexpensive surrogate of overall muscle strength and a valid predictor of physical disability and mobility limitation [51], remaining an objective measure of physical performance, even when performed by different operators [52] or different brands of dynamometers [53]. Of interest, HGS is also negatively associated with physical frailty even when the effects of BMI and arm muscle circumference are removed [54].

Is it well-known that the muscle strength is largely affected by nutrition, and specific dietary food components have been suggested to play an important role in the age-associated decline in physical performance [55]. Previous studies have focused on either individual nutrients, such as protein [56–58] or individual food groups [59,60]. However, diet includes a complex combination of several foods from various groups and nutrients, and some nutrients have highly similar properties, thus it is very challenging, in free-living populations, to separate the effect of a single nutrient or food group from others [61]. Consequently, there has been an increasing interest in using dietary pattern to evaluate the effects of nutrition on age-associated decline in physical performance.

The health benefits of a Mediterranean-style diet have been well documented for chronic disease morbidity and mortality [62,63]. Nevertheless, the large body of existing literature concerning MD and body weight focused on young individuals, with the aim of preventing and reducing obesity and cardiovascular diseases [64]. However, the World's population is ageing individuals older than 65 yrs represent an important and growing proportion and pose a considerable healthcare expense [65]. Dietary pattern-based

interventions have the potential of being a cost-effective solution for preventing or potentially slowing age-associated declines, including physical performance [27]. Several studies have investigated the association between Mediterranean-style diet, muscle mass, muscle strength, and muscle quality of older population living in different geographic areas. In these studies, the adherence to the MD was assessed using different questionnaires, mainly the 9-item MD according to the method developed by Trichopoulou et al. [66], while the muscular fitness was measured by means of different methods, such as free-fat mass, lower Limb Extension Power (LEP) or Short Physical Performance Battery (SPPB: standing balance, walking speed, and chair stand tests), and HGS [26,27,67]. These studies revealed some discrepancies in the relationship between adherence to the MD and the measures of physical performance. Bollwein et al. detected an association between a high dietary quality and a lower risk of low walking speed and low physical activity in community-dwelling older adults of both sexes living in the North Europe, but not with HGS [68]. More recently, Fougère et al. examined the association between adherence to the MD and two physical performance measures (i.e. SPPB score and HGS) among older individuals of adults of both sexes aged 77 years and older from the TRELONG study in the North Italy. Accordingly, the Authors found a statistically significant association between higher adherence to the MD and higher performance lower limbs, but again not with HGS [69]. In line with these findings, Keilaiti et al. correlated the intake of the individual food components of the MD in relation to the arm muscle quality in a large sample of women distributed over a wide age range (18–79 yrs) from the Twins UK registry, and found that fruit and alcohol intake were significantly associated with arm muscle quality, without significant associations with intakes of legumes, fish, or dairy products. However, in this study, the Authors failed to find any correlation with HGS and MD [17].

Our study is the first to report a strong correlation between the adherence the MD and HGS. It is well known that the development of frailty and the age-associated declines in physical function and skeletal muscle strength have been partially due to inflammation and oxidative stress that invariably is associated to aging [27]. Therefore, in line with other clinical studies, the higher intakes of several antioxidant micronutrients, including β -carotene and vitamins C and E, and other food bioactives, such as polyphenolic compounds present in plant foods or n3-PUFA present in fish and nuts, the low to moderate intake of alcohol and oils, such as extra-virgin olive oil, associated with a Mediterranean-style diet may also be responsible for the positive association between the adherence to the MD and HGS in our group of elderly women. Conversely, our study did not evidence any association between age and HGS. A possible explanation of this discrepancy could be due to the very homogeneous characteristics of our study sample. It is well known that gender, age, and educational status represent strong determinants in cross-sectional studies; in particular, gender influences food choice behaviours and it is a consolidated assumption that women's dietary profiles are characterized by higher carbohydrates intake, including fruit and vegetables compared to males [70,71]. It is also known that marital status influences the dietary behaviours, as married women and those who shared food preparation with family members had higher MD scores, compared to those who were single or had sole responsibility for food preparation, respectively [72]. In order to improve the power of the study, we increased the homogeneity of the subjects sample by including females only to avoid the effect of gender-specific food choice in dietary intake and anthropometric measures. The age range of the community-dwelling elderly women was relatively low, since only 24% of them were ≥ 75 yrs. In addition, as all participants were recruited at the local church, most of them lived in the same

neighbourhood, were more likely to be in the same social network, the vast majority were married/widowed, the educational level was very similar, and were not engaged in any physical activities. The lack of daily physical activities was not surprising in our study population of elderly women, taking also into consideration that this study was carried out during the winter season. Of interest, the association between HGS and PREDIMED score was also independent of protein intake. This finding, the lack of association between HGS and total energy intake, and the lack of physical activity, let us to exclude that, in our study population of elderly women, the positive association between the adherence the MD and HGS could be the result of higher energy or protein intake or physical activity, which, in the elderly population is likely associated with better muscle strength, rather than any Mediterranean diet effect.

We are aware that there are some limitations in the current study. First, the cross-sectional nature of this study did not allow any statements on a causal relationship between MD and HGS. Second, the sample size is relatively small and dietary intake is related to a larger number of other lifestyle factors. Nevertheless, the sample size was calculated using 80% power; in addition, the association between MD and HGS was independent of the effects of other health-related behaviours (physical activity, BMI) and protein intake. Third, the PREDIMED score, although easy to be accepted by the participants, allows only relative but no absolute statements about the degree of adherence to the MD. Fourth, the study sample, stratified according to HGS, included different BMI ranges. It is well-known that obesity is associated to increased risk of functional limitation. Thus, HGS cut-points for increased risk for mobility limitation may need to be examined separately for normal-weight, overweight and obese subjects in old population. However, as also reported by Sallinen J et al., while among men the HGS cut-points for mobility increased along with BMI, in women a single HGS threshold seemed to be sufficient at any level of BMI was identified [73]. In any case, in our study, the association between the adherence to the MD or its single items with HGS was adjusted for BMI. In addition, not only the PREDIMED questionnaire was face-to-face administered and not self-reported but, to avoid the inter-operator variability, only one expert Nutritionist administered the questionnaires, evaluated the anthropometric measures and determined the HGS. Anyway, the suggested cut-point for the PREDIMED score for identify the highest HGS values should be viewed with caution until results of studies in larger population samples have become available to perform an appropriate cross-validation.

4.1. Conclusions

This is the first study to evidence a positive association between adherence to the MD evaluated by PREDIMED score and its individual foods with the HGS in elderly active women. These results would recommend the evaluation of adherence of MD in the nutrition assessment as good clinical practice in clinical setting, thereby supporting the beneficial effects of nutritional interventions promoting a Mediterranean food pattern as an inexpensive and safe adjuvant prevention and treatment for age-related decline in physical performance. In this respect, this study i) suggests that, also without specific HGS measurements, the determination of a specific cut-point for the score of adherence to MD might help identifying subjects at high-risk for low muscle strength, who could get benefit of personalized dietary interventions; ii) further supports the collaborative role of Nutritionists in the evaluation and management of elderly population, highlighting the usefulness of the development of health services to detect and prevent age-associated decline in physical performance in elderly subjects by carefully addressing nutritional.

Future studies on male elderly or dietary intervention trials on large series population of both sexes will be critical for unravelling the beneficial effects of the MD on the HGS in elderly population.

Conflict of interest

None of the authors had a conflict of interest.

Authors' contributions

The authors' responsibilities were as follows LB and SS: were responsible for the concept and design of the study and interpreted data and drafted the manuscript;

LB: conducted statistical analyses; CDS, GM, GT, VDL, MI and AC: provided a critical review of the manuscript.

All authors contributed to and agreed on the final version of the manuscript.

Funding sources

None of the authors had a conflict of interest. The authors declare no support from any commercial organization for the submitted work.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.clnu.2018.03.012>

References

- <http://www.wcrf.org/>.
- Millen BE, Abrams S, Adams-Campbell L, Anderson CA, Brenna JT, Campbell WW, et al. The 2015 dietary guidelines advisory committee scientific report: development and major conclusions. *Adv Nutr* 2016;7:438–44. <https://doi.org/10.3945/an.116.012120>.
- Kieft-de Jong JC, Mathers JC, Franco OH. Nutrition and healthy ageing: the key ingredients. *Proc Nutr Soc* 2014;73:249–59. <https://doi.org/10.1017/S0029665113003881>.
- Sofi F, Abbate R, Gensini GF, Casini A. Accruing evidence on benefits of adherence to the Mediterranean diet on health: an updated systematic review and meta-analysis. *Am J Clin Nutr* 2010;92:1189–96. <https://doi.org/10.3945/ajcn.2010.29673>.
- Estruch R, Ros E, Martínez-González MA. Mediterranean diet for primary prevention of cardiovascular disease. *N Engl J Med* 2013;369:676–7. <https://doi.org/10.1056/NEJMc1306659>.
- Calton EK, James AP, Pannu PK, Soares MJ. Certain dietary patterns are beneficial for the metabolic syndrome: reviewing the evidence. *Nutr Res* 2014;34:559–68. <https://doi.org/10.1016/j.nutres.2014.06.012>.
- Esposito K, Maiorino MI, Bellastella G, Chiodini P, Panagiotakos D, Giugliano D. A journey into a Mediterranean diet and type 2 diabetes: a systematic review with meta-analyses. *BMJ Open* 2015;5:e008222. <https://doi.org/10.1136/bmjopen-2015-008222>.
- Martinez-Gonzalez MA, Martin-Calvo N. Mediterranean diet and life expectancy; beyond olive oil, fruits, and vegetables. *Curr Opin Clin Nutr Metab Care* 2016;19:401–7. <https://doi.org/10.1097/MCO.0000000000000316>.
- Park SY, Murphy SP, Wilkens LR, Yamamoto JF, Sharma S, Hankin JH, et al. Dietary patterns using the Food Guide Pyramid groups are associated with sociodemographic and lifestyle factors: the multiethnic cohort study. *J Nutr* 2005;135:843–9.
- Leblanc V, Bégin C, Hudon AM, Royer MM, Corneau L, Dodin S, et al. Gender differences in the long-term effects of a nutritional intervention program promoting the Mediterranean diet: changes in dietary intakes, eating behaviors, anthropometric and metabolic variables. *Nutr J* 2014;13:107. <https://doi.org/10.1186/1475-2891-13-107>.
- Hearty AP, McCarthy SN, Kearney JM, Gibney MJ. Relationship between attitudes towards healthy eating and dietary behaviour, lifestyle and demographic factors in a representative sample of Irish adults. *Appetite* 2007;48:1–11. <https://doi.org/10.1016/j.appet.2006.03.329>.
- Wardle J, Haase AM, Steptoe A, Nillapun M, Jonwutiwes K, Bellisle F. Gender differences in food choice: the contribution of health beliefs and dieting. *Ann Behav Med* 2004;27:107–16. PMID: 15053018.
- Trichopoulos A, Orfanos P, Norat T, Bueno-de-Mesquita B, Ocké MC, Peeters PH, et al. Modified Mediterranean diet and survival: EPIC-elderly prospective cohort study. *BMJ* 2005;330:991. <https://doi.org/10.1136/bmj.38415.644155.8F>.
- Knoops KT, de Groot LC, Kromhout D, Perrin AE, Moreiras-Varela O, Menotti A, et al. Mediterranean diet, lifestyle factors, and 10-year mortality in elderly European men and women: the HALE project. *JAMA* 2004;292:1433–9. <https://doi.org/10.1001/jama.292.12.1433>.
- Martínez-Gómez D, Guallar-Castillón P, León-Muñoz LM, López-García E, Rodríguez-Artalejo F. Combined impact of traditional and non-traditional health behaviors on mortality: a national prospective cohort study in Spanish older adults. *BMC Med* 2013;11:47. <https://doi.org/10.1186/1741-7015-11-47>.
- Vasto S, Barera A, Rizzo C, Di Carlo M, Caruso C, Panotopoulos G. Mediterranean diet and longevity: an example of nutraceuticals? *Curr Vasc Pharmacol* 2014;12:735–8.
- Kelaiditi E, Jennings A, Steves CJ, Skinner J, Cassidy A, MacGregor AJ, et al. Measurements of skeletal muscle mass and power are positively related to a Mediterranean dietary pattern in women. *Osteoporos Int* 2016;27:3251–60. <https://doi.org/10.1007/s00198-016-3665-9>.
- Michel JP, Cruz-Jentoft AJ, Cederholm T. Frailty, exercise and nutrition. *Clin Geriatr Med* 2015;31:375–87. <https://doi.org/10.1016/j.cger.2015.04.006>.
- Cruz-Jentoft AJ, Kiesswetter E, Drey M, Sieber CC. Nutrition, frailty, and sarcopenia. *Aging Clin Exp Res* 2017;29:43–8. <https://doi.org/10.1007/s40520-016-0709-0>.
- Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001;56:M146–56. PMID: 11253156.
- Masel MC, Ostir GV, Ottenbacher KJ. Frailty, mortality, and health-related quality of life in older Mexican Americans. *J Am Geriatr Soc* 2010;58:2149–53. <https://doi.org/10.1111/j.1532-5415.2010.03146.x>.
- Mitnitski A, Song X, Skoog I, Broe GA, Cox JL, Grunfeld E, et al. Relative fitness and frailty of elderly men and women in developed countries and their relationship with mortality. *J Am Geriatr Soc* 2005;53:2184–9. <https://doi.org/10.1111/j.1532-5415.2005.00506.x>.
- Ensrud KE, Ewing SK, Taylor BC, Fink HA, Stone KL, Cauley JA, et al. Frailty and risk of falls, fracture, and mortality in older women: the study of osteoporotic fractures. *J Gerontol A Biol Sci Med Sci* 2007;62:744–51.
- Cooper R, Kuh D, Hardy R, Mortality Review Group; FALCon and HALCyon Study Teams. Objectively measured physical capability levels and mortality: systematic review and meta-analysis. *BMJ* 2010;341:c4467. <https://doi.org/10.1136/bmj.c4467>.
- Newman AB, Simonsick EM, Naydeck BL, Boudreau RM, Kritchevsky SB, Nevitt MC, et al. Association of long-distance corridor walk performance with mortality, cardiovascular disease, mobility limitation, and disability. *JAMA* 2006;295:2018–26. <https://doi.org/10.1001/jama.295.17.2018>.
- Milaneschi Y, Bandinelli S, Corsi AM, Lauretani F, Paolisso G, Dominguez LJ, et al. Mediterranean diet and mobility decline in older persons. *Exp Gerontol* 2011;46:303–8. <https://doi.org/10.1016/j.exger.2010.11.030>.
- Talegawkar SA, Bandinelli S, Bandeen-Roche K, Chen P, Milaneschi Y, Tanaka T, et al. A higher adherence to a Mediterranean-style diet is inversely associated with the development of frailty in community-dwelling elderly men and women. *J Nutr* 2012;142:2161–6. <https://doi.org/10.3945/jn.112.165498>.
- Kim SW, Lee HA, Cho EH. Low handgrip strength is associated with low bone mineral density and fragility fractures in postmenopausal healthy Korean women. *J Kor Med Sci* 2012;27:744–7. <https://doi.org/10.3346/jkms.2012.27.7.744>.
- Bohannon RW, Schaubert KL. Test-retest reliability of grip-strength measures obtained over a 12-week interval from community-dwelling elders. *J Hand Ther* 2005;18:426–7. <https://doi.org/10.1197/j.jht.2005.07.003>.
- Norman K, Stobäus N, Gonzalez MC, Schulzke JD, Pirlich M. Hand grip strength: outcome predictor and marker of nutritional status. *Clin Nutr* 2011;30:135–42. <https://doi.org/10.1016/j.clnu.2010.09.010>.
- White JV, Guenter P, Jensen G, Malone A, Schofield M, Academy Malnutrition Work Group; A.S.P.E.N. Malnutrition Task Force; A.S.P.E.N. Board of Directors. Consensus statement: academy of nutrition and Dietetics and american society for parenteral and enteral nutrition: characteristics recommended for the identification and documentation of adult malnutrition (undernutrition). *J Parenter Enter Nutr* 2012;36:275–83. <https://doi.org/10.1177/0148607112440285>.
- Günther CM, Bürger A, Rickert M, Crispin A, Schulz CU. Grip strength in healthy caucasian adults: reference values. *J Hand Surg Am* 2008;33:558–65. <https://doi.org/10.1016/j.jhbs.2008.01.008>.
- Budziareck MB, Puzos Duarte RR, Barbosa-Silva MC. Reference values and determinants for handgrip strength in healthy subjects. *Clin Nutr* 2008;27:357–62. <https://doi.org/10.1016/j.clnu.2008.03.008>.
- Chilima DM, Ismail SJ. Nutrition and handgrip strength of older adults in rural Malawi. *Publ Health Nutr* 2001;4:11–7. PMID: 11255491.
- Frederiksen H, Hjelmberg J, Mortensen J, McGue M, Vaupel JW, Christensen K. Age trajectories of grip strength: cross-sectional and longitudinal data among 8,342 Danes aged 46 to 102. *Ann Epidemiol* 2006;16:554–62. <https://doi.org/10.1016/j.annepidem.2005.10.006>.
- Peolsson A, Hedlund R, Öberg B. Intra- and inter-tester reliability and reference values for hand strength. *J Rehabil Med* 2001;33:36–41. PMID: 11480468.
- Chainani V, Shaharyar S, Dave K, Choksi V, Ravindranathan S, Hanno R, et al. Objective measures of the frailty syndrome (hand grip strength and gait

- speed) and cardiovascular mortality: a systematic review. *Int J Cardiol* 2016;215:487–93. <https://doi.org/10.1016/j.ijcard.2016.04.068>.
- [38] Bohannon RW. Hand-grip dynamometry predicts future outcomes in aging adults. *J Geriatr Phys Ther* 2008;31(1):3–10. PMID: 18489802.
- [39] van Velsen L, Illario M, Jansen-Kosterink S, Crola C, Di Somma C, Colao A, et al. A community-based, technology-supported health service for detecting and preventing frailty among older adults: a participatory design development process. *J Aging Res* 2015;2015:216084. <https://doi.org/10.1155/2015/216084>.
- [40] <https://perssilaa.com/>.
- [41] Geneva: World Health Organization. *Waist circumference and waist-hip ratio. Report of WHO expert consultation. World health organization (WHO); 2008.*
- [42] Expert Panel on detection evaluation, and treatment of high blood cholesterol in adults. Executive summary of the third report of the national cholesterol education program (NCEP) expert Panel on detection, evaluation, and treatment of high blood cholesterol in Adults (Adult Treatment Panel III). *JAMA* 2001;285:2486–97. <https://doi.org/10.1001/jama.285.19.2486>.
- [43] World health organization (WHO). The problem of obesity. 2000 [Last accessed on 15 August 2015]. Available from: [http://www.who.int/trs/WHO_TRS_894_\(part1\).pdf](http://www.who.int/trs/WHO_TRS_894_(part1).pdf).
- [44] Martínez-González Miguel Angel, García-Arellano Ana, Toledo Estefanía, Salas-Salvadó Jordi, Buil-Cosiales Pilar, Corella Dolores, et al. A 14-item mediterranean diet assessment tool and obesity indexes among high-risk subjects: the PREDIMED trial. *PLoS One* 2012;7:e43134. <https://doi.org/10.1371/journal.pone.0043134>.
- [45] Roberts HC, Denison HJ, Martin HJ, Patel HP, Syddall H, Cooper C, et al. A review of the measurement of grip strength in clinical and epidemiological studies: towards a standardised approach. *Age Ageing* 2011;40:423–9. <https://doi.org/10.1093/ageing/afr051>.
- [46] Massy-Westropp NM, Gill TK, Taylor AW, Bohannon RW, Hill CL. Hand Grip Strength: age and gender stratified normative data in a population-based study. *BMC Res Notes* 2011;4:127. <https://doi.org/10.1186/1756-0500-4-127>.
- [47] Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, et al. Sarcopenia: european consensus on definition and diagnosis: report of the european working group on sarcopenia in older people. *Age Ageing* 2010;39:412–23. <https://doi.org/10.1093/ageing/afq034>.
- [48] Roman B, Carta L, Martínez-González MA, Serra-Majem L. Effectiveness of the Mediterranean diet in the elderly. *Clin Interv Aging* 2008;3:97–109. PMID: PMC2544374.
- [49] Hulens M, Vansant G, Lysens R, Claessens AL, Muls E, Brumagne S. Study of differences in peripheral muscle strength of lean versus obese women: an allometric approach. *Int J Obes Relat Metab Disord* 2001;25(5):676–81.
- [50] Lauretani F, Russo CR, Bandinelli S, Bartali B, Cavazzini C, Di Iorio A, et al. Age-associated changes in skeletal muscles and their effect on mobility: an operational diagnosis of sarcopenia. *J Appl Physiol* 1985;2003(95):1851–60. <https://doi.org/10.1152/jappphysiol.00246.2003>.
- [51] Rantanen T, Guralnik JM, Foley D, Masaki K, Leveille S, Curb JD, et al. Midlife hand grip strength as a predictor of old age disability. *JAMA* 1999;281:558–60. PMID: 10022113.
- [52] Mathiowetz V. Comparison of Rolyan and Jamar dynamometers for measuring grip strength. *Occup Ther Int* 2002;9:201–9. PMID: 12374997.
- [53] Smidt N, van der Windt DA, Assendelft WJ, Mourits AJ, Devillé WL, de Winter AF, et al. Interobserver reproducibility of the assessment of severity of complaints, grip strength, and pressure pain threshold in patients with lateral epicondylitis. *Arch Phys Med Rehabil* 2002;83:1145–50. PMID: 12161838.
- [54] Syddall H, Cooper C, Martin F, Briggs R, Aihie Sayer A. Is grip strength a useful single marker of frailty? *Age Ageing* 2003;32:650–6. PMID: 14600007.
- [55] Granic A, Jagger C, Davies K, Adamson A, Kirkwood T, Hill TR, et al. Effect of dietary patterns on muscle strength and physical performance in the very old: findings from the newcastle 85+ study. *PLoS One* 2016;11:e0149699. <https://doi.org/10.1371/journal.pone.0149699>.
- [56] Komar B, Schwingshackl L, Hoffmann G. Effects of leucine-rich protein supplements on anthropometric parameter and muscle strength in the elderly: a systematic review and meta-analysis. *J Nutr Health Aging* 2015;19:437–46. <https://doi.org/10.1007/s12603-014-0559-4>.
- [57] Rondanelli M, Faliva M, Monteferrario F, Peroni G, Repaci E, Allieri F, et al. Novel insights on nutrient management of sarcopenia in elderly. *BioMed Res Int* 2015;2015:524948. <https://doi.org/10.1155/2015/524948>.
- [58] Murphy CH, Oikawa SY, Phillips SM. Dietary protein to maintain muscle mass in aging: a case for per-meal protein recommendations. *J Frailty Aging* 2016;5:49–58. <https://doi.org/10.14283/jfa.2016.80>.
- [59] Kim J, Lee Y, Kye S, Chung YS, Kim KM. Association of vegetables and fruits consumption with sarcopenia in older adults: the fourth Korea national health and nutrition examination survey. *Age Ageing* 2015;44(1):96–102. <https://doi.org/10.1093/ageing/afu028>.
- [60] Beaudart C, Dawson A, Shaw SC, Harvey NC, Kanis JA, Binkley N, et al. Nutrition and physical activity in the prevention and treatment of sarcopenia: systematic review. *Osteoporos Int* 2017;28:1817–33. <https://doi.org/10.1007/s00198-017-3980-9>.
- [61] Hu FB. *Curr Opin Lipidol* 2002;13:3–9. PMID: 11790957.
- [62] Sofi F, Macchi C, Abbate R, Gensini GF, Casini A. Mediterranean diet and health status: an updated meta-analysis and a proposal for a literature-based adherence score. *Publ Health Nutr* 2014;17:2769–82. <https://doi.org/10.1017/S1368980013003169>.
- [63] Dinu M, Pagliai G, Casini A, Sofi F. Mediterranean diet and multiple health outcomes: an umbrella review of meta-analyses of observational studies and randomised trials. *Eur J Clin Nutr* 2017. <https://doi.org/10.1038/ejcn.2017.58>.
- [64] Buckland G, Bach A, Serra-Majem L. Obesity and the Mediterranean diet: a systematic review of observational and intervention studies. *Obes Rev* 2008;9:582–93. <https://doi.org/10.1111/j.1467-789X.2008.00503.x>.
- [65] http://www.un.org/en/development/desa/population/publications/pdf/ageing/WPA2015_Report.pdf.
- [66] Trichopoulos A, Costacou T, Bamia C, Trichopoulos D. Adherence to a Mediterranean diet and survival in a Greek population. *N Engl J Med* 2003;348:2599–608. <https://doi.org/10.1056/NEJMoa025039>.
- [67] Hashemi R, Motlagh AD, Heshmat R, Esmailzadeh A, Payab M, Yousefinia M, et al. Diet and its relationship to sarcopenia in community dwelling Iranian elderly: a cross sectional study. *Nutrition* 2015;31:97–104. <https://doi.org/10.1016/j.nut.2014.05.003>.
- [68] Bollwein J, Diekmann R, Kaiser MJ, Bauer JM, Uter W, Sieber CC, et al. Dietary quality is related to frailty in community-dwelling older adults. *J Gerontol A Biol Sci Med Sci* 2013;68:483–9. <https://doi.org/10.1093/geron/gis204>.
- [69] Fougère B, Mazzucco S, Spagnolo P, Guyonnet S, Vellas B, Cesari M, et al. Association between the mediterranean-style dietary pattern score and physical performance: results from TRELONG study. *J Nutr Health Aging* 2016;20:415–9. <https://doi.org/10.1007/s12603-015-0588-7>.
- [70] Ares G, Gábaro A. Influence of gender, age and motives underlying food choice on perceived healthiness and willingness to try functional foods. *Appetite* 2007;49:148–58. <https://doi.org/10.1016/j.appet.2007.01.006>.
- [71] Westenhoefer J. Age and gender dependent profile of food choice. *Forum Nutr* 2005;57:44–51. PMID: 15702587.
- [72] Papadaki A, Wood L, Sebire SJ, Jago R. Adherence to the Mediterranean diet among employees in South West England: formative research to inform a web-based, work-place nutrition intervention. *Prev Med Rep* 2015;2:223–8. <https://doi.org/10.1016/j.pmedr.2015.03.009>.
- [73] Sallinen J, Stenholm S, Rantanen T, Heliövaara M, Sainio P, Koskinen S. Hand-grip strength cut points to screen older persons at risk for mobility limitation. *J Am Geriatr Soc* 2010;58:1721–6. <https://doi.org/10.1111/j.1532-5415.2010.03035.x>.