

Resolvin E1 for reducing vascular calcification

Neil MacRitchie^{1*} and Pasquale Maffia^{1,2,3*}

¹Centre for Immunobiology, Institute of Infection, Immunity and Inflammation, College of Medical, Veterinary and Life Sciences, University of Glasgow, Sir Graeme Davies Building, 120 University Place, Glasgow G12 8TA, UK; ²Institute of Cardiovascular and Medical Sciences, College of Medical, Veterinary and Life Sciences, University of Glasgow, 120 University Place, Glasgow G12 8TA, UK; ³Department of Pharmacy, University of Naples Federico II, Via Domenico Montesano, 49, 80131 Naples, Italy

This editorial refers to ‘The G-protein coupled receptor ChemR23 determines smooth muscle cell phenotypic switching to enhance high phosphate-induced vascular calcification’, by M. Carracedo et al., doi: 10.1093/cvr/cvy316.

Vascular calcification—the deposition of excess mineral within the vessel wall—is now recognized as a complex, actively controlled process, principally mediated at the level of the vascular smooth muscle cell (VSMC). The extent of calcium deposition within the vessel wall can be used to predict the risk of adverse cardiac events (ACEs).¹

In the absence of vascular pathology, VSMCs exist in a quiescent state characterized by a set of contractile proteins such as CNN1 (calponin 1) and ACTA2 (alpha smooth muscle actin) which allow VSMCs to carry out their primary function, namely to provide structural support and to mediate vasoreactivity. In response to vascular injury/pathology, VSMCs can undergo a phenotypic transition towards a more synthetic phenotype. This change is characterized by down-regulation of contractile proteins, increased proliferation and migration and enhanced extracellular matrix production. It is now widely thought that in disease, VSMCs form a heterogeneous population with some VSMCs gaining an osteochondrogenic phenotype associated with osteogenic markers such as runt-related transcription factor 2 (RUNX2), type 1 collagen (Col1A1) and enhanced response to bone morphogenetic protein-2 (BMP-2).² The accumulative effect of this phenotypic switch is a cell type much more prone to accumulating calcium deposits within the vessel wall; however, the molecular pathways that promote this event are still not well defined.

In the study by Carracedo and colleagues³ recently published in *Cardiovascular Research*, the authors investigated the role of the ChemR23 receptor in calcification of epigastric arteries derived from kidney transplant patients. In chronic kidney disease, vascular calcification is an independent predictor of cardiovascular morbidity and mortality and the primary inducer of vascular calcification in these patients is thought to be hyperphosphatemia.⁴ ChemR23 is a G-protein coupled receptor expressed on multiple cell types including VSMCs,⁵ and is up-regulated during bone development. Chemerin was the first ChemR23 ligand discovered and induces pro-inflammatory properties via ChemR23 agonism.⁶ Subsequently, resolvin E1 (RvE1), an inflammatory resolving lipid mediator derived from the omega-3 fatty acid eicosapentaenoic acid (EPA), was also described in the literature as being a ligand for ChemR23⁷ (Figure 1); however, a direct interaction between RvE1

and ChemR23 is somewhat controversial with other receptors such as the leukotriene B4 receptor BLT1 also a candidate for mediating the effects of RvE1.⁸ In contrast to chemerin, the RvE1 interaction with ChemR23 has been reported to produce anti-inflammatory activity. RvE1 and other omega-3 derived specialized proresolvin mediators (SPM) have shown promise in resolving inflammation in several diseases in humans.⁹ Moreover, data from animal models indicates that administration of RvE1 results in reduced atherosclerosis,¹⁰ and that ERV1/ChemR23 signalling protects against atherosclerosis.¹¹ However, the role of ChemR23 and RvE1 in VSMC calcification and osteogenesis has been unexplored until now.

Gene expression analysis of patient arteries revealed ChemR23 to be an independent predictor of Col1A1. To obtain more insight into the effect of ChemR23 on VSMC phenotype, the authors next cultured VSMCs from ChemR23^{-/-} and wild-type mice. Under cell culture conditions, VSMCs spontaneously de-differentiate yet this process was greatly attenuated in ChemR23^{-/-} VSMCs. Furthermore, in response to elevated phosphate levels, calcification of cells was also attenuated in ChemR23-deficient VSMCs, as was activation of BMP-2 signalling, a known inducer of calcification in VSMCs under high phosphate conditions.² Concomitant with reduced calcification was a reduction in pro-calcific RUNX2 and an increase in anti-calcific osteoprotegerin. To test the hypothesis *in vivo*, mice were injected with vitamin D3, which mimics phosphate-induced calcification. ChemR23^{-/-} mice had reduced calcification in the medial layer of the aortic root and carotid artery. These were accompanied by biochemical changes indicative of a less calcific, more contractile VSMC phenotype, despite ChemR23^{-/-} mice displaying more pronounced hyperphosphatemia. This strengthens the view that the loss of ChemR23 is acting directly via VSMCs and not due to systemic metabolic changes, thus supporting the *in vitro* observations. RvE1-treated wild-type VSMCs showed reduced calcification and BMP-2 expression, effects not observed in ChemR23^{-/-} VSMCs. The reduction in calcification was not associated with other changes in VSMC phenotype indicating RvE1 has calcification specific effects mediated by ChemR23. Finally, the authors introduced the *Caenorhabditis elegans* Fat 1 transgene into wild-type and ChemR23^{-/-} mice. Fat 1 encodes a fatty acid desaturase that converts omega-6 to omega-3 fatty acids. The rationale being that enrichment of EPA results in greater production of RvE1 as previously observed.¹² In support of RvE1-induced inhibition of ChemR23 mediated vascular calcification, the authors noted that ChemR23




Figure 1 ChemR23 as a mediator of VSMC dysfunction. RvE1 is derived from the Omega-3 fatty acid, EPA (box). By acting as an inhibitory ligand on ChemR23, RvE1 causes a reduction in BMP-2 gene expression. BMP-2 promotes phosphate uptake via sodium-dependent phosphate cotransporter PIT-1, and through this mechanism, enhances VSMC calcification. Therefore, the reduction in BMP-2 expression observed after RvE1 treatment in ChemR23^{+/+} cells likely results in reduced calcification due to reducing the accumulation of intracellular phosphate. This effect was mediated by ChemR23 since ChemR23^{-/-} VSMCs did not show changes in calcification following RvE1 treatment. This study also reveals ChemR23 to be a positive modulator of VSMC proliferation/differentiation. A possible candidate for ChemR23 activation is chemerin, a ChemR23 agonist. Activation of ChemR23 can induce aberrant VSMC phenotypic changes through enhancement of reactive oxygen species (ROS) and MAPK signalling. BLACK arrows show observed phenotypic changes and RED arrows show pathways inhibited after RvE1 treatment under elevated phosphate (PO_4^{3-}) conditions.

dependent effects were attenuated in the presence of increased endogenous omega-3 production, potentially due to RvE1 inhibitory action.³

Derivatives of the omega-3 fatty acids EPA and docosahexaenoic acid (DHA) are thought to possess anti-inflammatory properties, acting to oppose the pro-inflammatory actions of cyclo- and lipoxygenase arachidonic acid products. The current work indicates that EPA derived RvE1 may also help in preventing the adverse phenotypic changes in VSMCs, such as accumulation of calcium, that run in parallel with vascular inflammation across a range of vascular pathologies. A multitude of studies looking at the cardioprotective effects of omega-3 supplementation have failed to reach a consensus on whether there is a beneficial effect on CVD morbidity or mortality. Recently, attention has focused on the EPA content in omega-3 formulations. A relatively low daily EPA dose (469 mg; 1-g total omega-3) formulation proved ineffective in reducing ACEs in diabetic patients.¹³ A different trial involving more than 8000 participants across 11 countries employed a total daily dose of 4 g of a purified EPA ethyl ester formulation in patients at risk of CVD and with elevated triglycerides. Despite the concurrent use of statins, patients taking the EPA treatment had a 25% reduction in risk of ischaemic events, including cardiovascular death, compared with those who received placebo.¹⁴ A similar trial (STRENGTH—Statin Residual Risk

Reduction With Epanova in High Cardiovascular Risk Patients with Hypertriglyceridaemia; ClinicalTrials.gov number, NCT02104817) is currently looking at the use of an EPA enriched omega-3 formulation (Epanova) in modifying ACEs in approximately 13 000 statin treated patients. A positive outcome from both these trials would increase the therapeutic focus onto EPA and its derivatives.

The understanding of endogenous SPM production as well as blood and tissue concentrations is an emerging area of research and the modulating role of these compounds in human disease is still unknown. Results between human subject studies to date are mixed regarding detection of circulating SPMs¹⁵ and it is currently unclear what the dose-relationship is to dietary intake of EPA. Translating *in vivo*-based studies to human disease will require sensitive mass spectrometry profiling of SPMs in biological fluids from patient and control subjects. This could determine if circulating SPMs are biomarkers for CVD and may offer insight into the relationship between omega-3 intake and cardiovascular protection. Such personalized information based on SPM production could enable clinicians to better tailor omega-3/EPA doses in future trials and may shed light on why some patients show a therapeutic response to omega-3 intake and others do not. A key future goal would also be to determine if a failure to resolve inflammation is related to depleted local SPM levels

and if administered EPA/DHA could reverse this trend. Furthermore, the identification of ligands and receptors such as RvE1 and ChemR23, respectively, could allow the use of more refined lipid preparations and also inform pharmacological approaches to design higher potency SPM analogues with a favourable anti-inflammatory/VSMC preserving profile.

In summary, the current work highlights a novel role for the ChemR23 receptor in promoting VSMC differentiation towards an osteoblastic, synthetic phenotype and highlights RvE1 as an inhibitory ligand against ChemR23 mediated VSMC calcification (Figure 1), a hallmark feature of human vascular pathology. Given the current lack of treatments for reversing or halting VSMC calcification, the discovery that RvE1 as an inhibitor of this process is an important development and may contribute to the cardiovascular benefits of EPA.

Conflict of interest: none declared.

Funding

Our laboratory is supported by the British Heart Foundation grants [PG/12/81/29897 to P.M., RE/13/5/30177]; the Engineering and Physical Sciences Research Council (EPSRC) grant [EP/L014165/1 to P.M.]; and Wellcome Trust Institutional Strategic Support Fund to P.M.

References

- Detrano R, Guerci AD, Carr JJ, Bild DE, Burke G, Folsom AR, Liu K, Shea S, Szklo M, Bluemke DA, O'Leary DH, Tracy R, Watson K, Wong ND, Kronmal RA. Coronary calcium as a predictor of coronary events in four racial or ethnic groups. *N Engl J Med* 2008;**358**:1336–1345.
- Li X, Yang HY, Giachelli CM. BMP-2 promotes phosphate uptake, phenotypic modulation, and calcification of human vascular smooth muscle cells. *Atherosclerosis* 2008;**199**:271–277.
- Carracedo M, Artiach G, Witasz A, Clària J, Carlström M, Laguna-Fernandez A, Stenvinkel P, Bäck M. The G-protein coupled receptor ChemR23 determines smooth muscle cell phenotypic switching to enhance high phosphate-induced vascular calcification. *Cardiovasc Res* 2018. doi:10.1093/cvr/cvy316.
- Yamada S, Giachelli CM. Vascular calcification in CKD-MBD: roles for phosphate, FGF23, and Klotho. *Bone* 2017;**100**:87–93.
- Ho KJ, Spite M, Owens CD, Lancero H, Kroemer AH, Pande R, Creager MA, Serhan CN, Conte MS. Aspirin-triggered lipoxin and resolvin E1 modulate vascular smooth muscle phenotype and correlate with peripheral atherosclerosis. *Am J Pathol* 2010;**177**:2116–2123.
- Davenport AP, Alexander SP, Sharman JL, Pawson AJ, Benson HE, Monaghan AE, Liew WC, Mpamhanga CP, Bonner TI, Neubig RR, Pin JP, Spedding M, Harmar AJ. International Union of Basic and Clinical Pharmacology. LXXXVIII. G protein-coupled receptor list: recommendations for new pairings with cognate ligands. *Pharmacol Rev* 2013;**65**:967–986.
- Arita M, Ohira T, Sun YP, Elangovan S, Chiang N, Serhan CN. Resolvin E1 selectively interacts with leukotriene B4 receptor BLT1 and ChemR23 to regulate inflammation. *J Immunol* 2007;**178**:3912–3917.
- Bondue B, Wittamer V, Parmentier M. Chemerin and its receptors in leukocyte trafficking, inflammation and metabolism. *Cytokine Growth Factor Rev* 2011;**22**:331–338.
- Moro K, Nagahashi M, Ramanathan R, Takabe K, Wakai T. Resolvins and omega three polyunsaturated fatty acids: clinical implications in inflammatory diseases and cancer. *World J Clin Cases* 2016;**4**:155–164.
- Hasturk H, Abdallah R, Kantarci A, Nguyen D, Giordano N, Hamilton J, Van Dyke TE. Resolvin E1 (RvE1) attenuates atherosclerotic plaque formation in diet and inflammation-induced atherogenesis. *Arterioscler Thromb Vasc Biol* 2015;**35**:1123–1133.
- Laguna-Fernandez A, Checa A, Carracedo M, Artiach G, Petri MH, Baumgartner R, Forteza MJ, Jiang X, Andonova T, Walker ME, Dall J, Arnardottir H, Gisterå A, Thul S, Wheelock CE, Paulsson-Berne G, Ketelhuth DFJ, Hansson GK, Bäck M. ERV1/ChemR23 signaling protects against atherosclerosis by modifying oxidized low-density lipoprotein uptake and phagocytosis in macrophages. *Circulation* 2018;**138**:1693–1705.
- Bilal S, Haworth O, Wu L, Weylandt KH, Levy BD, Kang JX. Fat-1 transgenic mice with elevated omega-3 fatty acids are protected from allergic airway responses. *Biochim Biophys Acta* 2011;**1812**:1164–1169.
- ASCEND Study Collaborative Group; Bowman L, Mafham M, Wallendszus K, Stevens W, Buck G, Barton J, Murphy K, Aung T, Haynes R, Cox J, Murawska A, Young A, Lay M, Chen F, Sammons E, Waters E, Adler A, Bodansky J, Farmer A, McPherson R, Neil A, Simpson D, Peto R, Baigent C, Collins R, Parish S, Armitage J. Effects of n-3 fatty acid supplements in diabetes mellitus. *N Engl J Med* 2018;**379**:1540–1550.
- Bhatt DL, Steg PG, Miller M, Brinton EA, Jacobson TA, Ketchum SB, Doyle RT Jr, Juliano RA, Jiao L, Granowitz C, Tardif JC, Ballantyne CM; REDUCE-IT Investigators. Cardiovascular risk reduction with icosapent ethyl for hypertriglyceridemia. *N Engl J Med* 2019;**380**:11–22.
- Murphy RC. Specialized pro-resolving mediators: do they circulate in plasma? *J Lipid Res* 2015;**56**:1641–1642.